

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

FOR USE IN SPRAYERS, MISTERS, AND OTHER FOGGING DEVICES:

DIRECTIONS FOR USE (Cont'd.)

To kill accessible stages of insects* infesting dried fruit products in storage or being processed: Close room or warehouse and shut off all ventilating systems. Dispense this product as a fine mist in the air above trays or shelves of dried fruit. Do not apply to fruit directly. If overhead sprayers are used, be sure that the spray is discharged from a position of at least 6 feet over the top trays of fruit. Use a dosage of 1 gal. of this spray per 50,000 cu. ft. of room space. Repeat applications as necessary for insect control, but do not exceed more than ten applications at this dosage in any given month.

*Indian Meal Moth, Dried Fruit Beetle, Confused Flour Beetle, Saw-Toothed Grain Beetle, Roaches.

To kill the adult and accessible forms of insects pests** of stored food (stored in multwall paper bags or in cloth bags) in Warehouses, Storage Rooms and similar locations:

Close room or warehouse and shut off all ventilating systems. Use a spraying device to give a fine mist, or use fixed ceiling sprayers that produce a very fine spray. Use a dosage of 1 fl. oz. of this spray per 1,000 cu. ft. of room space. Repeat applications as necessary for insect control.

**Rice Weevil, Yellow Meal Worm, Cadelle, Confused Flour Beetle, Saw-Toothed Grain Beetle, Angoumois Grain Moth, Mediterranean Flour Moth and Indian Meal Moth.

PRENTOX[®]

PRENTOX MIST SPRAY

CONCENTRATE #562

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY ESTABLISHMENTS. For control of Roaches, Waterbugs, Silverfish and Listed accessible forms of insects pests of stored fruit.

ACTIVE INGREDIENTS:

Pyrethrins	0.62%
Piperonyl Butoxide, Technical*	5.03%
Petroleum Distillates	94.35%
TOTAL:	100.00%

*Equivalent to 4.02% (Butylcarbityl)(6-propylpiperonyl) ether and 1.01% related compounds
PRENTOX[®]—Registered Trademark of Prentiss Drug & Chemical Co., Inc.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.

If inhaled — Remove person to fresh air. Apply artificial respiration if indicated.

If on skin — Remove contaminated clothing and wash affected areas with soap and water.

If in eyes — Flush eyes with plenty of water. Call a physician immediately.

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. If swallowed, do not induce vomiting. Call a physician immediately.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying in meat packing plants and other food processing plants, wash with potable water all equipment, benches, shelving, etc., where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use, or store near heat or open flame. Exposure to temperatures above 130°F. may cause bursting.

EPA REG. NO. 655-482

EPA EST. NO. 655-GA-1

NET CONTENTS _____

GALLONS

LOT NO. _____

Manufactured by



prentiss

Drug & Chemical Co., Inc.

Plant: Kaolin Road
Sandersville, GA. 31082

Office: C.B. 2000
Floral Park, N.Y. 11001

To kill flies, mosquito
Close room and disc
Use a dosage of 1/2 to
Leave room closed f
destroy fallen insect

As a surface spray k
insects, Weevils, Fl
Adults, Cadellae an

Apply as a coarse w
insects. Use a dose
particular attention
and ceilings. Spray
drains, conveyors ar
may be found (For E
Flies, Mosquitoes ar
surfaces that may c
residues in food mar

To kill roaches, silver
situations:

Apply this spray w
wetting droplets into
floors, underneath s
places where these i
for control. Do not
Direct spray into ar
crawling insects, hit

**CAUTION — KEEP C
DOMESTIC ANIMAL**

**HARMFUL IF SWALL
MIST. AVOID CON
SPILLED ON SKIN,
WHEN SPACE SF
TREATED AREAS
TREATMENT IS CO
FOODS SHOULD BE
TREATMENT. ALL
SHOULD BE COV
THOROUGHLY CL
USING THE PRODU
WHEN THE FACILI**

Do not use more than
cu. ft. of treated area.
not be used in the pr
pour or spill near h
closed. For control c
thorough and repe
working the into c
recommended. Empl
insects directly and c