

A PRENTISS PEST-TESTED PRODUCT

MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. DO NOT BREATHE SPRAY MIST. DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. WASH THOROUGHLY AFTER HANDLING. AVOID CONTAMINATION OF FOOD AND FEED. KEEP OUT OF REACH OF DOMESTIC ANIMALS. DO NOT USE ON HUMANS, HOUSEHOLD PETS, OR LIVESTOCK. DO NOT ALLOW CHILDREN IN TREATED AREAS UNTIL SURFACES ARE DRY.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT REUSE EMPTY DRUM. RETURN TO DRUM RECONDITIONER OR DESTROY BY PERFORATING OR CRUSHING, AND BURYING IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

Mix 7½ oz. of Concentrate with water to yield 1 gal. of finished solution containing 1% Diazinon, 0.1% Pyrethrins, and 0.5% Piperonyl Butoxide Technical. For half strength, use in maintenance work, dilute 3¾ oz. per gal. of water. Mix enough material for use on job. Do not store finished solution overnight.

IN NON FOOD AREAS OF FOOD PROCESSING ESTABLISHMENTS

COCKROACHES, (WATERBUGS), ANTS AND SILVERFISH — Make spot applications of ½% Diazinon as a coarse spray, or apply with a paint brush to baseboards, areas around water pipes, cracks and crevices, surfaces behind and beneath cabinets, refrigerators, sinks, stoves and similar areas where these insects hide. For heavy cockroach infestations, make the first application at a strength of 1% Diazinon, followed by monthly applications of ½% Diazinon.

CARPET BEETLES — Apply a ½% Diazinon spray in spot applications along baseboards and edges of carpeting, under carpeting and rugs, and furniture, in closets and on shelving, and wherever these insects are seen or suspected.

SPIDERS AND SCORPIONS — Apply a ½% Diazinon spray around window frames, doors, and along baseboards. Make spot applications to other areas over which these pests may crawl.



SYNPY-DIAZINON® EMULSIFIABLE CONCENTRATE

For Professional use only, for the control of cockroaches, ants, silverfish, carpetbeetles, spiders, and scorpions.

ACTIVE INGREDIENTS:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	20.0%
Pyrethrins	2.0%
Piperonyl Butoxide, Technical*	10.0%
Petroleum Distillates	55.0%
INERT INGREDIENTS:	13.0%

TOTAL 100.0%

*Equivalent to 8.0% (Butylcarbityl) (6-propylpiperonyl) Ether and 2.0% related compounds.

WARNING

KEEP OUT OF REACH OF CHILDREN

DIAZINON — Registered trademark of CIBA-GEIGY CORPORATION.

E. P. A. Est. No. 655 - NJ - 1

E. P. A. Reg. No. 655-487

NET CONTENTS

ACCEPTED

APR 17 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 655-487

PANTRY PESTS — Exposed stages (SAW-TOOTHED GRAIN BEETLE, CONFUSED FLOUR BEETLE, RED FLOUR BEETLE, RICE WEEVIL, CIGARETTE BEETLE, DRUGSTORE BEETLE, INDIAN-MEAL MOTH) — Apply a ½% spray to shelves and cupboards where food containers suspected of being infested are stored. Remove all food and shelving paper before spraying. Do not replace the food packages until spray has dried. Cover shelves with shelving paper. This treatment is supplemental to good housekeeping, since it will have no effect on insects breeding within food packages. Infested food packages should be destroyed.

BROWN DOG TICKS IN BUILDINGS — Make thorough application of a ½% spray to infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of household pets, and localized areas of floors and floor coverings. Fresh bedding should be placed in animal quarters following treatment. Do not treat animals with this formulation.

FOOD HANDLING ESTABLISHMENTS

Places other than private residences, in which food is held, processed, prepared and/or served.

FOOD AREAS — APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY:

Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing) preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) also would be considered a food area. Mix 7½ oz. of Concentrate with water to yield 1 gallon of finished solution. Apply in small amounts directly into cracks and crevices using equipment capable of delivering a pin stream of insecticide in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases where roaches, ants, silverfish, spider's hide and to aid in the control of pantry pests — exposed stages — (saw-toothed grain beetle, confused flour beetle, red flour beetle, rice weevil, cigarette beetle, drugstore beetle, Indian meal-mouth). **DO NOT** use this product in conduits, motor housings and electrical switch boxes. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN CRACK AND CREVICE TREATMENT ARE NOT PERMITTED

NON-FOOD AREAS — Includes garbage rooms, lavatories, floor drains (to sewers) entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

Apply ½-1% Diazinon as a coarse spray or apply with a paint brush to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, pallets and similar areas where insects hide or through which they may enter.

Prentiss Drug & Chemical Co., Inc.

PLANT: 358 WILSON AVENUE, NEWARK, N. J. 07105

OFFICE: 363 SEVENTH AVENUE, NEW YORK, N. Y. 10001