


PYRONYL CROP SPRAY

ACTIVE INGREDIENTS:

Pyrethrins .....	6.00%
Piperonyl Butoxide, Technical * .....	60.00%
Petroleum Distillates .....	14.00%
INERT INGREDIENTS: .....	20.00%
TOTAL:	<u>100.00%</u>



\* Equivalent to 48.0% (Butylcarbityl) (6-propylpiperonyl) Ether and 12.0% related compounds.



**CAUTION**

KEEP OUT OF REACH OF CHILDREN  
HARMFUL IF SWALLOWED.

CAUTION - DO NOT USE IN UNDILUTED FORM.

This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.


DO NOT REUSE EMPTY CONTAINERS. DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE, OR RETURN TO DRUM RECONDITIONER.



NET CONTENTS \_\_\_\_\_ GAL.

E.P.A. Reg. No. 655-489

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY -



TERMS OF SALE OR USE: On purchase or use of this product, buyer and user agree to the following conditions:

WARRANTY: Prentox warrants that this product, as of the time of sale by Prentiss Drug & Chemical Co. (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. PRENTISS DRUG & CHEMICAL CO. MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Prentiss Drug & Chemical Co. within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user, or over which Prentiss has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Prentiss from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

DIRECTIONS FOR USE -

PRENTOX<sup>(R)</sup> PYRONYL<sup>(R)</sup> CROP SPRAY is designed for use alone as a clean-up spray or as a pre-harvest spray where other materials cannot be used because of residue restrictions. When applied according to accepted use cautions and directions, PRENTOX PYRONYL CROP SPRAY may also be used in combination with other insecticides where resistant insects are a problem, or as an "exciter", where insects are flushed out of hiding and into contact with spray residues.

Apply the listed amount per acre and repeat as required to maintain effective control. Use in sufficient water for thorough coverage (usually 100 gallons) unless otherwise noted for listed crops. Coverage of upper and lower leaf surfaces is essential for good control.

ACCEPTED  
653-485  
SEP 12 1973  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER.

DIRECTIONS FOR USE: (Cont.)

When used alone, PRETOX PYRONYL CROP SPRAY may be used up to and including day of harvest.

ASPARAGUS: Asparagus Beetle - 2 to 6 oz.

BEANS: Aphids, Leafhoppers, White Fly, Mexican Bean Beetle, 12-Spotted Cucumber Beetle - 2 to 6 oz.

BROCCOLI, CABBAGE, BRUSSELS SPROUTS, CAULIFLOWER: Aphids, Cabbage Looper, Cross-striped Cabbageworm, Diamondback Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug - 2 to 6 oz.

CELERY: Green Peach Aphid, Cabbage Looper, Leaf Tier - 2 to 6 oz.

CRANBERRIES: Fireworms, Leafhopper - 2 to 6 oz.

EGGPLANTS: Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid - 2 to 6 oz.

LETTUCE: Cabbage Looper, Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm - 2 to 6 oz.

PEPPERS: Green Peach Aphid - 2 to 6 oz.

POTATOES: Aphids, Colorado Potato Beetle, Flea Beetle, Leafhopper - 2 to 6 oz.

RADISH: Aphids, Flea Beetle - 2 to 6 oz.

SPINACH: Aphids, Cabbage Looper, Webworm - 2 to 6 oz.

TOMATOES: Green Peach Aphid, Colorado Potato Beetle, Stink Bug, Flea Beetle - 2 to 6 oz.

MOSQUITOES: May be used on croplands as an adulticide to protect agricultural workers during harvesting. Use at 1 oz. per 4 gals. water per acre when flown on by fixed wing aircraft.

DROSOPHILA: Fruit Fly Control: Dilute at the rate of 1 part PRETOX PYRONYL CROP SPRAY with 1200 parts water (1 pint per 150 gallons or 1 tablespoonful with 4 gallons water). Thoroughly mix the emulsion in the spray tank. Make treatments as follows:


1. Apply liberally to tomatoes and fruits in baskets, on trucks and in plants. Use sprayers at a high pressure for applying at the rate of five or six pints of the diluted spray to a two-ton load of tomatoes. Direct the spray for a maximum coverage of the baskets or hampers. ~~It is important to spray between and beneath the containers.~~ This spray not only kills the flies, but ~~the emulsion kills~~ any dead flies so that they are readily washed from the fruit.

ACCEPTED

655-489

SEP 12 1973

DIRECTIONS FOR USE: (Cont.)



2. Spray the raw stock stacked in the yard.
3. Dip baskets in the diluted spray after dumping the fruit to kill adhering larvae and pupae.
4. After washing and cleaning up the inside of the processing plant and just prior to bringing fruit into it, the entire space inside the cannery should be sprayed at a dilution of 1 part PRENTOX PYRONYL CROP SPRAY to 59 parts water (1 pint with 7 gallons 3 pints water or 2 tablespoonfuls with 3-1/4 pints water). At the rate of 1 gallon to 750 square feet (1 quart to 187 sq. ft.) direct spray on walls, ceilings, and floors, paying special attention to forcing the spray into all cracks and crevices for the control of ants, roaches, silverfish, crickets, spiders and cheese mites.

For use as a space spray in the processing plants, this dilution will give excellent control of flying insects, i.e. fruit flies, house flies, hornets, grain moths, gnats, mosquitoes and skipper flies. Apply at the rate of one-half ounce per 1000 cubic feet of space. Do not spray while the plant is in operation, as dead flies may fall into containers or the product being processed. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleared before using.

SPACE SPRAY: To control flying insects such as fruit flies, hornets, wasps, grain moths, gnats, mosquitoes and skipper flies, dilute one part PRENTOX PYRONYL CROP SPRAY with 59 parts water (2 tablespoonfuls with 3-1/4 pints water). Use at the rate of one-half ounce per 1000 cubic feet of space. Direct the space treatment upward and whenever practical keep doors and windows closed for at least ten minutes after application. Do not remain in treated areas. Ventilate the area after treatment is completed.

VAPORIZER APPLICATORS: Where oil residues are not undesirable, PRENTOX PYRONYL CROP SPRAY can be diluted in deodorized base oil instead of water, at the dilution rate of 1-59, and applied by means of any good vaporizer type of applicator.

**ACCEPTED**