


A PRENTISS PEST-TESTED PRODUCT

Prentox

VAPOR ?

EMULSIFIABLE CONCENTRATE INSECTICIDE

For Professional Use Only.



ACTIVE INGREDIENTS:	
Aromatic Petroleum Derivatives	65.10%
2,2-Dichlorovinyl dimethyl phosphate	22.69%
Related compounds	1.71%
INERT INGREDIENTS:	7.50%
	<hr/>
TOTAL:	100.00%

Contains 2 lbs. DDVP per gallon.

DANGER

POISON


KEEP OUT OF REACH OF CHILDREN

SEE OTHER PRECAUTIONS AND ANTIDOTE
ON SIDE PANELS.

NET CONTENTS

GALLONS

655-491



EPA Reg. No. 105-491

A PRENTISS PEST-TESTED PRODUCT

Prentox

PRECAUTIONS IN USING:


PRENTOX VAPON 2 EMULSIFIABLE CONCENTRATE and the diluted emulsion are poisonous. Therefore, precautions must be observed when handling them. (See antidote and warnings below). Store the Concentrate in a safe place. Children or animals must not be able to reach or handle the Concentrate or diluted emulsion. It is recommended that only the amount of diluted emulsion that will be used immediately be prepared from the Concentrate. Mix the Concentrate into water outdoors or in a well ventilated room. Wash off the outside of the bottle of Concentrate in running water before storing away. Do not contaminate feed, water or foodstuffs, milk or milking utensils.

THIS PRODUCT IS TOXIC TO FISH, BIRDS AND OTHER WILDLIFE. KEEP OUT OF LAKES, STREAMS, OR PONDS. DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREAS TREATED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DANGER - PRENTOX VAPON 2 EMULSIFIABLE CONCENTRATE IS POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN OR EYES. DO NOT GET ON SKIN OR EYES. DO NOT BREATHE FUMES. AVOID EXPOSURE TO FUMES. WEAR CLEAN RUBBER GLOVES, WATERPROOF PROTECTIVE CLOTHING AND GOGGLES WHEN HANDLING THE CONCENTRATE TO PREPARE EMULSIONS. WEAR A MASK OR RESPIRATOR OF A TYPE PASSED BY THE U.S. BUREAU OF MINES FOR DDVP (VAPONA INSECTICIDE) PROTECTION. KEEP ALL UNPROTECTED PERSONS OUT OF OPERATIONAL AREA. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. WASH THOROUGHLY WITH SOAP AND WATER BEFORE EATING OR SMOKING. WASH ALL CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE REUSE.

DECONTAMINATE AND DESTROY USED CONTAINERS.

DECONTAMINATE CONTAINERS BY WASHING THOROUGHLY WITH A STRONG SOLUTION OF ALKALI AND DETERGENT, AND RINSING WITH WATER. DESTROY CONTAINERS BY CRUSHING OR PUNCTURING; THEN BURY OR BURN. DISPOSE OF RINSINGS IN A WAY AS NOT TO CONSTITUTE A HAZARD. DO NOT CONTAMINATE WATER SUPPLIES.



1974

A PRENTISS TEST-TESTED PRODUCT



ANTIDOTE & FIRST AID TREATMENT:

Atropine is the emergency antidote for VAPON Insecticide poisoning. Consult your Physician about obtaining an adequate supply of 1/100-grain Atropine tablets for emergency use. **CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING.**

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100-grain Atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** (See WARNING SYMPTOMS below).

Keep patient prone and quiet. Start artificial respiration immediately, if patient is not breathing. Transport the patient immediately to the nearest Physician.



NOTE TO PHYSICIAN - WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, Atropine Sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete Atropinization has occurred. Morphine is contraindicated. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

Clear chest by postural drainage.

Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor.

Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.



A PRENTISS PEST-TESTED PRODUCT

Prentox

DIRECTIONS FOR USE - TO BE APPLIED BY PEST CONTROL OPERATORS OR TRAINED PERSONNEL ONLY -

DO NOT USE IN EDIBLE PRODUCTS AREAS OF FOOD PROCESSING PLANTS, RESTAURANTS OR OTHER AREAS WHERE FOOD IS COMMERCIALY PREPARED OR PROCESSED. DO NOT USE IN SERVING AREAS WHILE FOOD IS EXPOSED.

OUTDOOR LIVING AREAS, PICNIC GROUNDS, BACKYARD AREA, PATIOS, LOADING DOCKS, OUTDOOR LATRINES, PARKING AREA, REFUSE AREAS AROUND SERVICE STATIONS, OPEN-AIR DRIVE-INS, ICE CREAM STANDS, GARBAGE COLLECTION AND DISPOSAL AREAS.

Apply VAPON 2 Insecticide as a 0.5% spray to control flies, gnats, mosquitoes, roaches, fleas, ants, sowbugs, and brown dog ticks.

To prepare a 0.5% spray, dilute 1 pint of VAPON 2 Emulsifiable Concentrate with 6 gallons of water. Apply as a coarse wet spray at the rate of 1 quart per 1,000 sq. ft. Do not treat animals or humans. Do not contaminate water, food, food containers or cooking utensils. Repeat as necessary.

Apply VAPON 2 Insecticide as a 1% to 3% fog or mist solution to control flies, mosquitoes, wasps, gnats, and other small flying insects in outdoor areas.


To prepare a 1% fogging or misting solution, dilute 1 quart of VAPON 2 Emulsifiable Concentrate with 6 gallons of water. For a 1% oil solution, dilute 1 quart with 7 gallons of oil.

MANURE PILES, GARBAGE DUMPS - (PCO)

For the control of fly maggots in maggot breeding areas, apply at the rate of 1-2 quarts of 0.5% spray per 100 sq. ft. (1 quart of 1%).

DOG RUNWAYS -

Apply to outside dog runways, window sills, and ledges at a rate of 1 quart of 0.5% spray per 1,000 sq. ft. For the control of flea and brown dog ticks, increase dosage to 2 quarts of 0.5% spray per 1,000 sq. ft. (PCO)



A PRENTISS POST-TESTED PRODUCT

Prentox

APR 1974
DIVISION DIVISION
S. P.

OUTDOOR AREAS - CONTROL OF FLIES, MOSQUITOES, GNATS -

Apply with fogging apparatus or mist blowers at the rate of 5-10 pints of 1% fogging solution per acre (equivalent to 0.05-0.1 pound VAPON Insecticide per acre). Do not treat areas where food or feed crops are growing.

FOOD PROCESSING PLANTS, INDUSTRIAL PLANTS, THEATERS, WAREHOUSES - FOR THE CONTROL OF FLIES, ROACHES, ANTS, SILVERFISH

Apply with fogging or misting equipment at the rate of 1 gallon of 1% fogging solution per 64,000 cubic feet, for control of flies. When fogging for control of roaches, ants and silverfish, the fog should be directed into cracks and crevices and hiding places. Apply when plants are not in operation. Occupants should vacate before indoor treatments and should not reoccupy the premises until they have been ventilated. After application, open several windows or doors, to create air currents to insure reduction of VAPON Insecticide vapors. Food should be removed and food handling equipment covered with draping paper during application, or washed after application is completed. During fogging or mist spraying indoors, use a mask or respirator of a type passed by the USDA for DDVP (VAPONA Insecticide) protection.

HOUSEHOLD PEST CONTROL

Apply VAPON Insecticide as a 0.5% spray to control flies, mosquitoes, wasps, gnats, flying moths, and other small flying insects, cockroaches, ants, spiders, silverfish. To prepare a 0.5% spray, dilute 1 pint of VAPON 2 Emul. Conc. with 6 gallons of water.

For Households, Restaurants, Theaters, Food Processing Plants, Industrial Plants, Warehouses, apply 0.5% spray with low pressure sprayer to localized areas insects may infest. Treat infested areas around baseboards, cracks, walls, door and window frames, and localized areas of floors. Do not apply to areas which will be contacted frequently by children and do not allow them in treated areas until surfaces are dry. Do not contaminate food, water, food containers, or cooking utensils. Do not treat animals.

WARRANTY

NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE EXPLICIT OR IMPLIED, concerning this product or its uses, which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

