

A PRENTISS PEST-TESTED PRODUCT

HOUSEHOLD PEST CONTROL Apply as a 0.5% Spray to control flies, mosquitoes, wasps, gnats, flying moths and other small flying insect, cockroaches, ants, spiders, silverfish. To prepare a 0.5% spray, dilute 1 pint of Concentrate with 6 gal. of water.

For household, restaurants, theaters, food processing plants, industrial plants, warehouses, apply 0.5% spray with low pressure sprayer into all cracks and crevices. Do not apply to areas which will be contacted frequently by children, and do not allow them in treated areas until surfaces are dry. Do not contaminate food, water, food containers, or cooking utensils. Do not treat animals.

ANTIDOTE AND FIRST AID TREATMENT: Atropine is the emergency antidote for DDVP insecticide poisoning. Consult your Physician about obtaining an adequate supply of 1/100 grain Atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat, or by giving warm, salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove the person from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If material gets into eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100 grain Atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See WARNING SYMPTOMS).

KEEP PATIENT PRONE AND QUIET.

START ARTIFICIAL RESPIRATION IMMEDIATELY, IF A PATIENT IS NOT BREATHING.

TRANSPORT THE PATIENT IMMEDIATELY TO THE NEAREST PHYSICIAN.

NOTE TO PHYSICIAN:

WARNING SYMPTOMS Symptoms include weakness, headaches, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT Atropine is the specific therapeutic antagonist of cholinergic autonomic nervous stimulation. There are no signs of anticholinergic stimulation. Atropine sulfate should be injected in 1-2 minute intervals, in doses of 1-2 mgms. until complete atropinization is achieved. 4-8 AM for atropine. It may be administered in conjunction with Atropine. Morphine is contraindicated.

Clear chest by postural drainage.

Repeated exposure to cholinergic stimulants may, with a warning, cause prolonged convulsions. Convulsions will subside if chlorpromazine is used.

Allow further exposure until a fatal dose has been administered, as determined by breath test.

E.P.A. REG. NO. 655-492

E.P.A. EST. NO. 655-NJ-1



NET CONTENTS


GALLONS



VAPON 20% EMULSIFIABLE CONCENTRATE

FOR PROFESSIONAL USE ONLY

ACTIVE INGREDIENTS:	
Aromatic Petroleum Derivatives	73.0%
2,2-Dichlorovinyl dimethyl phosphate	18.6%
Related compounds	1.4%
INERT INGREDIENTS:	7.0%
TOTAL	100.0%



DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

PRECAUTIONS IN USING. PRENTOX VAPON 20% CONCENTRATE and the diluted emulsion are poisonous. Therefore, precautions must be observed when handling them (see Antidotes and Warnings). Store the Concentrate in a safe place. Children or animals must not be able to reach or handle the Concentrate or diluted emulsion. It is recommended that only the amount of diluted emulsion that will be used immediately be prepared from the Concentrate.

Mix the Concentrate into water outdoors or in a well ventilated room. Wash off outside of bottle of Concentrate in running water before storing away.

ENVIRONMENTAL CAUTIONS This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

SEE ANTIDOTE STATEMENT AND OTHER PRECAUTIONS ON BACK SIDE PANEL.

PRENTOX VAPON 20% is poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not breathe fumes. Avoid exposure to fumes. Wear clean rubber gloves, waterproof protective clothing and goggles when handling this concentrate to prepare emulsion. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for DDVP protection, keep all unprotected persons out of operational area. In case of contact, wash immediately with soap and water. Wash thoroughly with soap and water after use and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

DECONTAMINATE AND DESTROY USED CONTAINERS - Decontaminate used containers by washing thoroughly with a strong solution of alkali and detergent and rinsing with water.

DESTROY CONTAINERS BY CRUSHING OR PUNCTURING THEN BURY OR BURN. DISPOSE OF RINSINGS IN A WAY AS NOT TO CONSTITUTE A HAZARD, OR CONTAMINATE WATER SUPPLIES.

DIRECTIONS FOR USE - TO BE APPLIED BY PEST CONTROL OPERATORS OR TRAINED PERSONNEL ONLY.

Outdoor living areas, picnic grounds, backyard area, patios, loading docks, outdoor latrines, parking area, refuse areas around Service Stations, open-air Drive-ins, Ice Cream Stands, garbage collection and disposal areas.

Apply as a 0.5% spray to control flies, gnats, mosquitoes and other small flying insects. To prepare a 0.5% spray, dilute 1 pint of this Concentrate with 6 gal. of water. Apply as a coarse wet spray at the rate of 1 quart per 1,000 sq. ft. Do not treat animals (at this rate) or humans. Do not contaminate water, food, food containers or cooking utensils. Repeat as necessary.

Apply as a 1% to 3% fog or mist solution to control flies, mosquitoes, wasps, gnats and other small flying insects in outdoor areas.

To prepare a 1% fogging or misting solution, dilute 1 qt. of Concentrate with 6 gal. of water. For a 1% oil solution, dilute 1 qt. with 7 gal. of oil.

Manure piles, Garbage dumps: For the control of fly maggots in maggot breeding areas, apply at the rate of 1-2 quarts of 0.5% spray per 100 sq. ft. (1 quart of 1%).

DOG KENNELS: Apply to outside dog runways, window sills and ledges at a rate of 1 qt. or 0.5% spray per 1,000 sq. ft. For the control of fleas and brown dog ticks, increase dosage to 2 quarts of a 0.5% spray per 1,000 sq. ft.

Outdoor areas: For the control of flies, gnats, mosquitoes, ants, fleas, brown dog ticks, apply with fogging apparatus or mist blowers at the rate of 5-10 pints of 1% fogging solution per acre. Do not treat areas where food or feed crops are growing.

FOOD PROCESSING PLANTS, INDUSTRIAL PLANTS, THEATERS, WAREHOUSES, NON-FOOD AREAS: For the control of flies, gnats, mosquitoes, apply with fogging or misting equipment at the rate of 1 gal. of 1% fogging solution per 84,000 cu. ft. Apply when plants are not in operation. Occupants should vacate before indoor treatments, and should not reoccupy the premises until they have been ventilated. After application, open several windows or doors, to create air current, to insure reduction of DDVP insecticide vapors. During fogging or mist spraying, always use a mask or respirator of a type passed by the U.S. Bureau of Mines for DDVP insecticide protection.

FOOD HANDLING ESTABLISHMENTS Places where food is prepared and served should be protected from insecticide spray, fogging or misting.

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY:

Crack and crevice areas for application include parking areas, restrooms, and other areas where people frequent. Do not apply to food, food containers, or food processing equipment. Do not apply to food, food containers, or food processing equipment. Do not apply to food, food containers, or food processing equipment.

Apply 0.5% spray in small amounts directly into cracks and crevices. Some equipment capable of delivering a fine spray may be used to apply to different openings. Do not spray between equipment and floor openings leading to walls and follow cracks in walls, equipment legs and bases, etc. Do not apply to ants and silverfish hide.

Warning: Shock hazard exists if this product is used in radiators, motor housings and electrical switch boxes.

Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATION OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.

