

prentox

BEST DOCUMENT AVAILABLE



PYRONYL PET AND LIVESTOCK SPRAY CONCENTRATE

ACTIVE INGREDIENTS:	
Petroleum Distillates	78.80%
Piperonyl Butoxide, Technical*	12.00%
Pyrethrins	1.20%
INERT INGREDIENTS:	8.00%
TOTAL: 100.00%	

*Equivalent to 9.6% (Butylcarbityl)(6-propylpiperonyl) ether and 2.4% related compounds

PRENTOX® Registered Trademark of Prentiss Drug & Chemical Co., Inc.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist. Avoid skin and eye contact. Wash thoroughly after handling and before eating or smoking.

Do not allow children or pets to contact treated surfaces until spray has dried. Do not use on plants. Do not allow spray to contact food, feedstuffs or water supplies. Thoroughly wash dishes and food handling utensils contaminated with this product. Remove pets and cover fish bowls before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to lakes, streams and ponds. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use, or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticides and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required as to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician or Poison Control Center immediately. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia.

If inhaled — Remove victim to fresh air.

If on skin — Remove contaminated clothing and wash affected areas with soap and water.

If in eyes — Flush eyes with plenty of water. Call a physician immediately.

SEE SIDE PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 845-494 EPA EST. NO. 845-GA-1

NET CONTENTS _____

LOT NO. _____

DIRECTIONS FOR USE (Cont'd.)

To Control Insects in Kennels and Animal Quarters
Dilute at the rate of 13 fluid ounces in 1 gallon with water or 2 fluid ounces in 18 fluid ounces of water.

Remove pets, birds, and cover fish aquariums before spraying.

For rapid cleanup of fleas and ticks in kennels apply as a surface spray in a good machine adjusted to deliver coarse droplets. Apply to the point of run-off with special attention to cracks and crevices and other insect hiding places. Bedding should be removed and sprayed thoroughly.

To control flies, mosquitoes and gnats in kennels, apply this dilution in a good sprayer adjusted to deliver a fine mist. Use at the rate of approximately 1 ounce for each 1000 cubic feet of space. Fill the enclosed areas with mist. Wherever practical, keep doors and windows closed for 10 minutes or more after application. Do not remain in treated areas and ventilate after treatment is completed.

To Control Insects on Animals

FLEAS — To kill fleas on cats and dogs, dilute at 1 to 100, and thoroughly wet the animal by dipping or spraying. For best results against fleas and ticks on pets, kennels, animals quarters and bedding should also be treated.

Dilute this concentrate at the rate of 7 fluid ounces in 1 gallon with water or 1 fluid ounce in 18 fluid ounces of water.

For rapid control of ticks and lice on dogs and cats, apply this dilution as a mist spray. To insure thorough coverage, part the fur by stroking against the lay of the hair as spray is applied. Spray directly on ticks. Apply to moisture the fur lightly.

To control ear mites on cats and ear mites, syringe the ears with a few drops of this dilution so that it can penetrate the wax down to the surrounding tissues. Repeat treatment once or twice at 2 to 3 days intervals as necessary.

For temporary protection from mosquitoes, eye gnats, biting flies and chiggers apply to cover the surface of the hair. Repeat treatment as necessary when insects are troublesome.

HORSE, SHEEP, GOAT & HOG — Dilute 1 gal. of Concentrate with 100 gallons of water for control of active louse stages. Apply for thorough body coverage, using 1 to 3 gallons on horses, and 1 to 3 quarts on sheep, goats and hogs.

POULTRY — Dilute 1 gallon with 100 gallons of water for control of poultry lice and treat birds by submerging in a barrel, tank or tub, holding the wings away from the body, to insure thorough wetting of all infected areas. Dip on a warm day or in a warm building, to avoid chilling of fowls.

CATTLE — 1 gallon of Concentrate in 100 gallons of water gives effective control of the active stages of both sucking and biting lice, on beef and dairy cattle. Apply to thoroughly wet all portions of the body, using 1 to 3 gallons per head, depending on size of the animal. Repeated treatments are recommended due to hatching of eggs.

SHEEP "TICK" OR KED — Use 1 gallon of Concentrate in 100 gallons of water and thoroughly wet all portions of the body by dipping, or by spraying with sufficient pressure and a nozzle adjustment to give penetration into the wool. Treat at the rate of 1 to 4 pints per head, as determined by size of the animals and condition of fleece.

ACC
JUN 6 1984
655-444



Manufactured by
prentiss
Drug & Chemical Co., Inc

Plant: Kaolin Road
Sandersville, GA. 31082
Office: C.B. 2000
Floral Park, N.Y. 11001