

Prentox

**CAUTION
KEEP OUT OF REACH
OF CHILDREN**

**PYRONYL
OIL
CONCENTRATE
#1530**


ACCEPTED
655-496
NOV 2 1973
FOR THE FEDERAL REGISTER
INFRACONCENTRATE
ED UNDER NO
TO ATTACHED CO

CAUTION: HARMFUL IF SWALLOWED
DO NOT POUR OR SPILL NEAR OPEN FLAME.
MIX WELL IN CONTAINER BEFORE EACH USE.
KEEP CONTAINER CLOSED AND STORE IN A COOL PLACE.
DO NOT REUSE EMPTY DRUM. RETURN TO DRUM RECONDITIONER OR DESTROY BY PERFORATING OR CRUSHING, AND BURYING IN A SAFE PLACE.
APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DO NOT APPLY DIRECTLY TO FOOD. COVER OR REMOVE ANY FOOD AND FOOD PROCESSING EQUIPMENT DURING APPLICATION. DO NOT APPLY WHILE FOOD PROCESSING IS UNDERWAY. AFTER SPRAYING IN MEAT PACKING PLANTS, BAKERIES AND OTHER FOOD PROCESSING PLANTS, WASH ALL EQUIPMENT, BENCHES, SHELVING, ETC. WHERE EXPOSED FOOD WILL BE HANDLED, WITH POTABLE WATER.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

E. P. A. Reg. No. 655-496



ACTIVE INGREDIENTS:

Pyrethrins.....	1.50%
Piperonyl Butoxide, Technical*.....	3.00%
N-octyl bicycloheptene dicarboximide**.....	5.50%
Petroleum Distillates	90.00%
TOTAL:	100.00%

*Equivalent to 2.40% (Butylcarbityl) (6-propylpiperonyl) Ether and 0.60% related compounds.

**MGK-264 Insecticide Synergist.

NET CONTENTS _____

DIRECTIONS FOR USE

SPACE SPRAY - Use undiluted to kill accessible stages of insects* infesting dried fruit products in storage or being processed. Close room or warehouse and shut off all ventilating systems. Use a ULV spraying device to disperse this product as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use one ounce per 1000 cubic feet of space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-entry.

*Indian Meal Moth, Dried Fruit Beetle, Confused Flour Beetle, Saw-toothed Grain Beetle and Roaches.

SPACE SPRAY - Use undiluted to kill the adult and accessible forms of insect pests* of stored food (stored in multiwall paper bags or in cloth bags) in warehouses, storage rooms & similar locations. Close room or warehouse and shut off all ventilating systems. Use a ULV spraying device to disperse this product. First direct spray into corners under pallets and around stacks of packaged goods hitting as many insects as possible and then treated as a space spray. Infested stored products should be fumigated, or treated by other effective methods using an approved product intended for this purpose. Use a dosage of one fluid ounce of this spray per 1000 cubic feet of room space. Repeat applications as necessary for insect control. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-entry.

*Rice Weevil, Yellow Meal Worm, Cadelle, Confused Flour Beetle, Saw-toothed Grain Beetle, Angoumois, Grain Moth, Mediterranean Flour Moth and Indian Meal Moth.

SPACE SPRAY - Use diluted to kill flies, mosquitoes and gnats in open areas. Dilute concentrate with Petroleum Distillate such as Klearol N.F. at the rate of 1 part concentrate to 2 parts oil and mix well. Use at a time when the temperature is cool (75° or less) and the wind velocity is approximately 5 MPH, with a ULV spraying device, spray at a dosage rate of 15 ounces per minute up-wind of the area to be controlled. If the equipment is truck mounted, keep the speed to approximately 5 MPH with a swath width of no more than 300 feet. Repeat application as required.

SPACE SPRAY AND CONTACT SPRAY - Use undiluted to kill roaches, silverfish, waterbugs, earwigs, ants, spiders, in sheltered situations. With a ULV spraying device, apply the spray into all cracks and crevices in wood-work, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Do not allow spray to contact stored food. Use at a dosage rate of approximately one-half fluid ounce per 1000 cubic feet or room space. Repeat dosage as necessary for control. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-entry. Remove Pets, Birds and cover Fish Aquariums before spraying.