

DIRECTIONS FOR USE

STORAGE AND DISPOSAL

Do not contaminate water, food, feed, stock, or other animals. Do not dispose of this product in any way that is prohibited.

STORAGE: If container is damaged, stop any further use. If the container or by patching or other a resealing the case. Take care to avoid contact with pesticide and wear protective gear. One can of this product wears protective equipment as required to prevent contact with the insecticide vapors. Cover the spilled areas with general absorbent material such as clay, diatomaceous earth, sand, or sawdust. Transfer the contaminated absorbent to a shovel and put the waste in a plastic drum. Dispose of wastes as below. Pa. many taking into account the material drum or glass container. Do not store use of this insecticide in any place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this insecticide should be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse, if required, to avoid contamination or recycling or reconditioning or puncture and disposal in accordance with or by other procedures approved by State and local authorities.

DIRECTIONS FOR USE (Cont'd)

INDOOR USE AS A SPACE SPRAY UNDILUTED: The use of this insecticide for food processing or food handling establishments should be limited to short periods when the plants or facilities are not in operation. The insecticide should be removed or covered during treatment. Air conditioning systems should be covered during treatment. Clean up spillage before use.

Close windows and doors and shut off ventilating systems. Heat-treated vaporizer adjusted to deliver aerosol spray, nozzle opening 1/8 inch diameter and 80% less than 100 cubic feet. Apply at rate of 1/2 ounce per 1,000 cubic feet of space. Direct the spray toward ceiling and cover all areas of the area behind obstacles. This insecticide is not for use in areas where the treated area is occupied by people or animals. Repeat application as necessary. Sweep up and destroy fallen insects.


This method of application is not for use in areas where people are present, including FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, SMALL FLYING MOTHS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE BEETLES, SKIPPERS, BOXELDER BUGS, EARWIGS, ANTS, and other insects. Do not use in areas where people are present, including GRANARY WEEVILS, RICE WEEVILS, RICE WEEVILS, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANIMAL MEALWORMS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEALMOTH, TOBACCO MOTHS, YELLOW MEALWORMS, DARK MEALWORMS, MITE, CADELLES and RED FLOUR BEETLES.

CONTACT AND SURFACE SPRAY: For use in areas where people are present, adjust to deliver a coarse wet spray. Direct the spray toward cracks, crevices and devices, under pallets and containers, inside of bases of machinery, behind shelves and racks. Spray in the direction whenever possible. Repeat application as necessary.

DRIED FRUIT PRODUCTS: For storage or being processed, apply to CONFUSED FLOUR BEETLES, DRIED FRUIT BEETLES, INDIAN MEAL MOTHS, AND SAW-TOOTHED GRAIN BEETLES. Remove and destroy infested products. As noted above, disperse the product directly above the trays and shelves. Do not apply to food directly. Use one-half ounce per 1,000 cubic feet of space. Repeat application at this dosage every 6 months. Leave the room closed for one hour. Do not remain in treated area and ventilate the area before re-entry. Sweep up and destroy fallen insects.

STORED FOOD: (Stored in multi-wall bags or cloth bags in warehouses, storage rooms, flour mills, grain elevators, granaries, for control of ANGOUMOIS GRAIN MOTHS, CADELLES, CONFUSED FLOUR BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RICE WEEVILS, SAW-TOOTHED GRAIN BEETLES, YELLOW MEALWORMS, CHEESE MITES, CHEESE SKIPPERS, CIGARETTE BEETLES, GRAIN MITES, GRANARY WEEVILS, DARK MEALWORMS, RED FLOUR BEETLES. Remove and destroy infested products. As noted above, disperse the product directly, directing spray into corners, under pallets and around stacks of packages. Avoid hitting as many insects as possible and then treat as a space spray. Infested stored products should be fumigated or treated by other effective methods using an approved product. Intended for the purpose use a dosage of one-half fluid ounce per 1,000 cubic feet of space. Do not remain in treated areas and ventilate area before re-entry. This application will address the exposed stages of those stored product pests listed above. Sweep up and destroy fallen insects.

INDOOR USE AS A SPACE SPRAY, DILUTED: For use in conversion of mechanical fogging equipment to KILL FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 25 parts of refined kerosene and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before re-entry. Repeat treatment when reinfestation occurs.



Pyronyl Oil Concentrate OR-3610A

ONLY FOR SALE TO USE AND STORAGE BY SERVICE PERSONS

Authorized by U.S.D.A. for use in Federally inspected meat and poultry plants

ACTIVE INGREDIENTS:

Pyrethrins	3.00%
Piperonyl Butoxide Technical*	6.00%
N-octyl bicycloheptene dicarboximide**	10.00%
Petroleum Distillates	81.00%
TOTAL	100.00%

Consists of 4.50 fluid ounces per gallon. For use in Federally inspected meat and poultry plants.

PRENTOX® Registered Trademark of Prentiss, Inc.

E.P.A. REG. NO. 655-501 **E.P.A. EST. NO. 655-GA-1**

NET CONTENTS

LOT #

This product is effective in the control of the pests listed in the table of the application follows the directions for use as set forth in this table.

- ANGOUMOIS GRAIN MOTHS — ANTS — BOXELDER BUGS — CADELLES — CHEESE MITES — CHEESE SKIPPERS — CIGARETTE BEETLES — CLOVER MITES — CONFUSED FLOUR BEETLES — DARK MEALWORMS — DEER FLIES — DRIED FRUIT BEETLES — DRUGSTORE BEETLES — EARWIGS — FRUIT FLIES — (DROSOPHILA) — FUNGUS GNATS — GNATS — GRAIN MITES — GRANARY WEEVILS — HORN FLIES — HORNETS — HORSE FLIES — HOUSE FLIES — INDIAN MEAL MOTHS — LICE — MEDITERRANEAN FLOUR MOTHS — MOSQUITOES — MUSHROOM FLIES — RED FLOUR BEETLES — RICE WEEVILS — ROACHES — SAW-TOOTHED GRAIN BEETLES — SMALL FLYING MOTHS — SPIDER BEETLES — STABLE FLIES — TOBACCO MOTHS — WASPS — YELLOW MEALWORMS**

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Wash out with water. Do not induce vomiting. If not vomiting, give 10 to 15 ounces of water.

If inhaled: Remove to fresh air. If on skin: Remove contaminated clothing and wash affected areas with soap and water.

If in eyes: Flush eyes with plenty of water for at least 15 minutes immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful to water and aquatic life. Do not discharge into streams, rivers, or other bodies of water. Wash thoroughly after handling. Do not breathe dusts or mists. Avoid contact with eyes. Wash thoroughly after handling. Do not use if pregnant or nursing. Do not use if you are allergic to any of the ingredients. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you are taking any medication.

ENVIRONMENTAL HAZARDS

This product is highly toxic to fish and other aquatic life. Do not discharge into streams, rivers, or other bodies of water. Do not use if you are allergic to any of the ingredients. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you are taking any medication.

PHYSICAL HAZARDS

This product is highly flammable. Do not use near open flames or other sources of ignition. Do not use in confined spaces. Do not use if you are allergic to any of the ingredients. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you are taking any medication.



Manufactured by
Prentiss
Drug & Chemical Co. Inc

Plant: Kaolin Road
Sandersville, GA. 31082
Office: C.B. 2000
Floral Park, N.Y. 11001

SURFACE SPRAY
...
ANIMAL QUARTERS
...
IN MUSHROOM
...
FOR USE IN FEED
...
OUTDOOR GROUND
...
TEMPORARY
...
FOR USE ON ANIMALS
...

FEB 7 1984

Prentiss Drug & Chemical Co., Inc.
C. B. 2000
Floral Park, NY 11001

Attention: Mr. Nick Sanchez

Gentlemen:

Subject: Pretox Pyronyl Oil Concentrate
EPA Registration No. 655-501
Your Letter of January 13, 1984

The labeling submitted to fulfill the requirements of PP Notice 83-3
is acceptable. A stamped copy of the label is enclosed for your records.

Sincerely yours,

Timothy A. Gardner
Insecticide-Rodenticide Branch
Registration Division (20-767)

Enclosure

DCI-12118:Taxano. 1/26/84. D-91. C-2. M. 217. 157-20. 0:dc1. 2/11/84.pac

CONCURRENCES								
SYMBOL ▶	TS 76							
SURNAME ▶	Tavano							
DATE ▶	2/3/84							

EPA Form 1320-1 (4-81) OFFICIAL FILE COPY