

A PRENTISS PEST-TESTED PRODUCT

FISH AND WILDLIFE CAUTION:

This product is toxic to fish, shrimp, crabs, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT REUSE EMPTY CONTAINER. DESTROY BY PERFORATING OR CRUSHING, AND BURYING IN A SAFE PLACE.

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY —

IMPORTANT: Read the entire DIRECTIONS FOR USE and the CONDITIONS OF SALE AND WARRANTY before using this product.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Prentiss Drug & Chemical Co., Inc. or the Seller. All such risks shall be assumed by Buyer.

Prentiss Drug & Chemical Co., Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the DIRECTIONS FOR USE, subject to the inherent risks referred to above. Prentiss Drug & Chemical Co., Inc. makes no other express or implied warranty. In no case shall Prentiss Drug & Chemical Co., Inc. or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Prentiss Drug & Chemical Co., Inc. and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY which may be varied only by agreement in writing signed by a duly authorized representative of Prentiss Drug & Chemical Co., Inc.



DIAZINON® AG-600 INSECTICIDE

FOR CONTROL OF CERTAIN INSECTS ON FRUITS, NUTS, VEGETABLES, FIELD CROPS AND ORNAMENTALS.

ACTIVE INGREDIENTS:

0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate68.0%
Xylene6.6%

INERT INGREDIENTS:25.4%

TOTAL: 100.0%

DIAZINON® - Registered trademark of Ciba-Geigy Corp.
DIAZINON AG-600 contains 6 lbs. of Diazinon per gallon.

WARNING
KEEP OUT OF REACH OF CHILDREN

MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. DO NOT BREATHE SPRAY MIST. DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. WASH THOROUGHLY AFTER HANDLING. AVOID CONTAMINATION OF FOOD. DO NOT USE ON HUMANS OR PERMIT CHILDREN TO GO ON TO SPRAYED GRASS UNTIL SPRAY HAS COMPLETELY DRIED.

CAUTION:
FLAMMABLE MIXTURE.
Do not use, pour, spill or store near heat or open flame.

KEEP OUT OF REACH OF DOMESTIC ANIMALS, DO NOT CONTAMINATE FEED OR USE ON HOUSEHOLD PETS OR LIVESTOCK. DO NOT PERMIT PETS TO GO ON TO SPRAYED GRASS UNTIL SPRAY HAS COMPLETELY DRIED.

NET CONTENTS _____

E.P.A. Reg. No. 655-518
E.P.A. Est. No. 655-NJ-1

ACCEPTED
AUG 26 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 655-518

Prentiss Drug & Chemical Co., Inc.

PLANT 338 WILSON AVENUE, NEWARK, N. J. 07105

OFFICE 363 W. 42ND AVENUE, NEW YORK, N. Y. 10018

GENERAL INFORMATION

Diazinon AG-600 is an emulsifiable solution which when diluted with water according to the directions below is intended for the control of insect pests of agricultural crops. Do not use in barns or dwellings. Do not use on livestock.

In the directions for use given below, best control is obtained when application is made at first sign of insects. Application should be repeated as necessary to maintain control.

NOTE: The California Department of Agriculture has set a 5-day re-entry for Diazinon on citrus, grapes, peaches and nectarines. Five days should lapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organic phosphate pesticides are applied in combination, the interval shall be prolonged by adding to the longest applicable interval an additional 50% of that interval.

TOBACCO — Aphids, Flea Beetles: Apply 1/3-1/2 pt. per acre. Do not apply within 3 days of harvest. Grasshoppers: Apply 7/10 pt. per acre. Do not apply within 14 days of harvest. Cutworms: Broadcast 1½-2 qts. per acre before transplanting and work into the soil 2-4 inches immediately. Green June Beetle Larvae: Mix 1/6 pt. in 50 gals. of water and apply as a drench (100 gals. per 100 sq. yds.) to uprooted areas of plant beds with sprinkling can. Mole Crickets: Broadcast 7/10 qt. per acre, when insects are present. Make application to beds or fields 1-2 days prior to sowing of seed or transplanting, and work into soil 1-2 inches immediately. Wireworms: Broadcast 1½-2 qts. per acre, 3-4 weeks before transplanting. Work into soil 2 inches immediately.

PEANUTS — Cutworms: Apply 1½ qts. per acre in a 12-inch band over the row. Thorough coverage of the foliage is essential. Begin application when small larvae appear. 2 or 3 additional applications at 3 week intervals may be necessary. Do not apply within 21 days of peanut harvest. Livestock (1) may be fed peanut hulls (2) may be fed peanut forage 7 days following treatment and (3) may be fed peanut 21 days following treatment.

SUGARCANE — Aphids: Apply 1/3-7/10 pt. per acre. Allow 1 day between last application and harvest. Bagasse and forage may be fed to livestock. Wireworms: Apply 7/10 gal. per acre in a 12 inch band centered over the furrow. Make application directly to seed pieces in open furrow at covering time.

PINEAPPLES: Scales: Apply 3½ qts. of Diazinon AG-600 plus 2 gallons of light or medium horticultural oil in 500 gal. of water per acre to the vegetable slips on the plants following harvest of pineapple. Mealybugs: Apply 1½ qts. per acre in a minimum of 100 gal. of water. May be applied on day of harvest. Pineapple forage and refuse may be fed to livestock 7 days following last application.