DIRECTIONS FOR USE:



CONTACT SPRAY - Diluted for Use with Sprayers that give large, wetting Spray Particles - To kill Cockroaches and accessible stages of Granary Insects; Rice & Granary Weevil, Red Flour Beetle, Indian Meeal Moth Larvae, Cadelles, Grain Mites and Saw-Toothed Grain Beetles. Cover or remove exposed food and close room. Dilute Concentrate with Oil at the rate of 1 part Concentrate plus 9 parts Oil, and mix well. Spray outside walls of grain bins, flooring, about base of milling machinery, walls and baseboards, areas behind cabinets, around pipes and other insect harbourages using 1 gal. of Spray per 1,000 sq.ft. of surface. Force spray into all cracks and crevices. Repeat applications as necessary. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

SPACE SPRAY - Undiluted in Aerosol Generators - To kill Flies, Mosquitoes, Small Flying Moths & Gnats:

Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating equipment. Apply at the rate of 2 to 6 oz. of undiluted Concentrate per 1,000 cu.ft. of space. Direct output from aerosol generator upward towards upper corners of room and ceiling. Leave room closed for one half hour. Do not remain in treated ares. Ventilate the area after treatment is completed. Sweep up and destroy fallen insects. Repeat as necessary.

CPACE SPRAY - Dilute 1 plus 4 with Base Oil for Use in Barns & Poultry Houses, Warehouses, Storage Bins & Storage Rooms - To control House Flies, Stable Flies & Horn Flies, apply as a fine mist over the livestock in stanchions or holding pens or over the poultry in cages or in pens. Use a fogger or mist generator, so spray is directed and applied above the animals and direct spray to center of the area where unimals are housed. Apply 1 oz. per 1,000 cu.ft., so that the spray mist is distributed uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in Dairy Barns, aerate the barn thoroughly before milking. Repeat daily, morning and afternoon, or as required. Keep sprayer unit reservoir and nozzle clean, to ensure proper delivery of spray mist or fog. Product to be used in mechanical sprayers, which give particle size aerosol range or smaller, such as ULV type.

SPACE SPRAY - For Use in Mechanical Fogging Equipment - Kills Flies, Mosquitoes, Small Flying Moths & Gnats Cover or remove exposed food and food handling surfaces. Close room, making sure to shut off all air conditioning or ventilation equipment. Dilute Concentrate with Deodorized Kerosene at the rate of 1 part Concentrate with 9 parts Deodorized Kerosene and mix well. Apply at the rate of 50 fl. oz. of diluted spray per 25,000 cu.ft. of space. Make sure to direct the spray upward towards upper corners of room & ceiling.

Keep room closed for one half hour after spraying. Do not remain in treated area. Ventilate after treatment is completed.

Pyrethrins	2.00%
Piperonyl Butoxide, Technical *	4.00%
N-octyl bicycloheptene dicarboximide **	6.66%
Petroleum Distillates	87.34%
TOTAL:	100.00%

* Equivalent to 3.20% (Bucylcarbityl) (6-propylpiperonyl) Ether and 0.80% related compounds. ** MGK-26- Insecticide Synergist.

DO NOT REUSE EMPTY DRUM. RETURN TO DRUM RECONDITIONER OR DESTROY BY PERFORATING OR CRUSHING AND BURY-ING IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Remove birds and cover fish aquariums before spraying. NET CONTENTS

CAUTION KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. IN CASE OF CONTACT, WASH IMME-DIATELY WITH SOAP AND WATER. WASH ARMS, HANDS AND FACE THOROUGHLY WITH SOAP AND WATER BEFORE EATING OR SMOKING. WASH ALL CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE REUSE. WEAR NATURAL RUBBER GLOVES, WATER-FROOF PROTECTIVE CLOTHING AND GOGGLES. KEEP ALL UNPROTECTED PERSONS OUT OF OPERATIONAL AREA WHILE AP-PLYING THE INSECTICIDE, OR WORKING IN TREATED AREAS DURING POSTED PERIOD, WEAR A MASK OR RESPIRATOR OF A TYPE PASSED BY THE U.S. BUREAU OF MINES, AVOID CON-TAMINATION OF FEED AND FOODSTUFFS.

Apply this product only as specified on this label.

E.P.A. EST. NO 653 113 E.P.A. Reg. Nov 655-532



Prentiss Drug & Chemical Co. Inc.

PLANT: 338 WILSON AVENUE, NEWARK, N. J. 07105

CIFCE 301 SEVENTA AVENE NO WYORK N Y 1000