

A PRENTISS PEST-TESTED PRODUCT

ACCEPTED
 JUL 14 1975
 Under the Federal Insecticide, Fungicide, and Rodenticide Act
 Registration No. 100-100000-0000



DPBM CONCENTRATE #4425

FOR USE IN ULV TYPE MACHINES BY TRAINED PERSONNEL ONLY IN HOMES, RESTAURANTS, FOOD PROCESSING PLANTS, WAREHOUSES AND INDUSTRIAL INSTALLATIONS.

ACTIVE INGREDIENTS:

2,2-dichlorovinyl dimethyl phosphate ---	0.46%
Other related compounds -----	0.04%
Pyrethrins -----	2.00%
Piperonyl Butoxide, Technical * -----	4.00%
N-butyl bicycloheptene dicarboximide ---	4.00%
Petroleum Distillates -----	89.50%
TOTAL:	100.00%

NOTE TO PHYSICIAN: Atropine is antidotal.

DO NOT REUSE EMPTY DRUM. RETURN TO DRUM RECONDITIONER OR DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

* Equivalent to 3.2% (Butylcarbityl) (6-propylpiperonyl) Ether and 0.8% related compounds.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

WARNING KEEP OUT OF REACH OF CHILDREN

DO NOT POUR OR SPILL NEAR OPEN FLAME.

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. AVOID BREATHING OF SPRAY MIST. WEAR MASK OR RESPIRATOR OF A TYPE RECOMMENDED BY THE U.S. BUREAU OF MINES TO GIVE ADEQUATE PROTECTION AGAINST THIS MATERIAL. WEAR PROTECTIVE CLOTHING. WASH HANDS, ARMS AND FACE WITH SOAP AND WATER AFTER HANDLING AND BEFORE SMOKING. WASH ALL CONTAMINATED CLOTHING WITH SOAP AND WATER BEFORE RE-USE.

DO NOT USE IN EDIBLE PRODUCTS AREAS OF FOOD PROCESSING PLANTS, RESTAURANTS OR OTHER AREAS WHERE FOOD IS COMMERCIALY PREPARED OR PROCESSED. DO NOT USE IN SERVING AREAS WHILE FOOD IS EXPOSURE.

REMOVE FISH, BIRDS, AND OTHER FINE AQUARIUM BEFORE SPRAYING.



Mix well in container before each use. Keep container closed and store in cool place.

NET CONTENTS _____ GALLONS

E.P.A. EST. NO. 650-NJ-1

E.P.A. Reg. No. 656-513

A PRENTISS PEST-TESTED PRODUCT

Prentiss

DIRECTIONS FOR USE

GENERAL: This Concentrate is designed for use undiluted in mechanical sprayers which give particle size of aerosol size. It can be diluted with refined Kerosenes or Mineral Oil and can be applied in fogging equipment that gives larger than aerosol particle size, such as the Microsol or Challenger machines. It can also be diluted with oil specified above and applied as a contact spray to kill crawling insects shown later on this label.

Kills the Accessible Stages of INDIAN MEAL MOTH, DRIED FRUIT BEETLE AND CONFUSED FLOUR BEETLE in dried fruit products storage areas or processing areas. Close room or warehouse and shut off all ventilating systems. Use a ULV spraying device. Do not apply to dried fruit directly. Direct spray into areas of the room frequented by these insects, hitting as many insects as possible. Then spray into the air above the trays at a rate of 1/2 gal. of spray per 50,000 cu. ft. of space. Repeat application every 30 days. Leave room closed for one hour after application. Do not remain in treated areas and ventilate before re-entry. Infested stored dried fruit should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Kills the Accessible Stages of RICE WEEVIL, YELLOW MEALWORM, CADELLE, CONFUSED FLOUR BEETLE, SAW-TOOTHED GRAIN BEETLE, ANGOUMOIS GRAIN MOTH, MEDITERRANEAN FLOUR MOTH AND INDIAN MEAL MOTH in food warehouses, storage rooms and similar areas. Close room or warehouse and shut off all ventilating systems. Use a ULV spraying device. Direct spray into corners, under pallets and around stacks of packaged goods, hitting as many insects as possible. Then spray at the rate of 1 fl. oz. of spray per 1,000 cu. ft. of space. Repeat applications as necessary. Leave room closed for one hour after application. Do not remain in treated areas and ventilate prior to re-entry. Infested package goods should be fumigated or treated by other effective means using an approved product intended for this purpose.

To Kill LEACHES, SILVERFISH, EARWIGS, ANTS AND SPIDERS - Apply in ULV-type machine, or similar equipment adjusted to coarse spray. Apply the spray into all cracks and crevices in and on all walls, floors, and beneath sinks, behind pipes, and in all places where the insects are known to be present. Use 1/4 gal. per 1,000 cu. ft. of wall space or point of spray.

