

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

PRENTOX DDVP FIVE is specially formulated Insecticide designed for use in the control of exposed stages of stored-product insects infesting Warehouses containing non-perishable, packaged or bagged raw or processed food commodities. The term "non-perishable" means any raw or processed food not subject to rapid decay or deterioration that would render it unfit for consumption.

Raw commodities would include: cocoa beans, field-dried beans, field-dried peas, grains and nuts. Not included are: eggs, milk, meat, poultry, fresh fruits and vegetables such as onions, parsnips, potatoes and carrots.

Processed foods would include: flour, sugar, cereal, packaged cookies and crackers. Not included are hermetically sealed foods or manufactured dairy products and other processed foods requiring refrigeration.

USE & APPLICATION DIRECTION:

For the control of the exposed stages and the prevention of migration of most species of stored-product insects, rice weevils, confused flour beetles, saw-toothed grain beetles, cigarette beetles, Indian-meal moths and cocoa bean moths in Warehouse storage areas and in food processing areas of flour and grain mills, bakeries, canning and bottling plants and brewery warehouses during times when the plant is not in operation and no food products are exposed. If bulk, unpackaged foods are exposed, remove or cover before spraying begins. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas apply only when the facility is not in operation. Use DDVP FIVE in a ULV type machine, adjusted to deliver small particles. The rate of application should be between 1-2 grams of actual DDVP per 1,000 cu. ft. or 1/2 to 1 oz of liquid per 1,000 cu. ft. Check amount in insecticide tank, or auxiliary tank before and after treatment to assure proper dosage has been applied.

Treatment should be made during the hours when buildings are unoccupied. Buildings treated should be closed and ventilation kept at a minimum during application of the insecticide. Lock all entrances and do not allow unprotected workers to enter buildings while being treated.

Place Caution Signs at all entrances stating "This building should be opened and aired prior to re-entry of unprotected workers, or remain unoccupied by unprotected workers for at least 6 hours after treatment. Do not enter prior to _____ o'clock _____ (date) without protection." Repeat treatment at weekly intervals during season of peak insect activity for optimum results.

SPACE SPRAY AND CONTACT SPRAY — Dilute to 1-3% and apply at the rate of 1 gallon of spray per 64,000 cu. ft. to kill roaches, silverfish, fleas, waterbugs, earwigs, ants, spiders, in sheltered situation. Adjust a ULV spraying device to a coarse spray for control of the above listed crawling insects. Contact as many of the insects with the spray as possible. Thoroughly spray all parts of the room suspected of harboring these insects. Special attention should be paid to cracks, hidden surfaces under equipment, and food storage areas. Then adjust the ULV device to deliver a fine spray for use as a space spray. Sweep up and destroy dead insects. Repeat treatment as necessary.

FOR USE AS A CONTACT SPRAY — Dilute to 1% and spray 1 pint per 10,000 cubic feet or 1,000 square feet of floor space. Do not use in edible product areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in sewing areas while food is exposed.

prentox[®] D.D.V.P. FIVE

FOR USE BY PEST CONTROL OPERATIONS OR TRAINED PERSONNEL ONLY

FOR USE IN WAREHOUSES, RAILROAD CARS AND STATIONERY TRUCKS, FLOUR MILLS, GRAIN MILLS, RICE MILLS.

AUTHORIZED BY U.S.D.A. FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY ESTABLISHMENTS.

ACTIVE INGREDIENTS:

2,2-dichlorovinyl dimethyl phosphate* . . . 4.65%

Related Compounds 0.35%

Petroleum Distillates 15.00%

INERT INGREDIENTS: 80.00%

TOTAL: 100.00%

*Equivalent to 5% Vapona Insecticide

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

**KEEP OUT OF REACH OF CHILDREN
WARNING**

**PRECAUTIONARY STATEMENTS
HAZARDOUS TO HUMANS**

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN OR EYES. DO NOT GET ON SKIN OR IN EYES. DO NOT BREATHE FUMES. WEAR NATURAL RUBBER GLOVES, WATER-PROOF PROTECTIVE CLOTHING AND GOGGLES WHEN HANDLING. WEAR A MASK OR RESPIRATOR OF A TYPE PASSED BY THE U.S. BUREAU OF MINES FOR PROTECTION AGAINST THIS MATERIAL. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. WASH HANDS, ARMS, AND FACE THOROUGHLY WITH SOAP AND WATER BEFORE EATING OR SMOKING. WASH ALL CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE REUSE. NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

ANTIDOTE & FIRST AID TREATMENT:
ATROPINE is the emergency antidote. Consult your Physician about obtaining an adequate supply of 1/100-gram Atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

E.P.A. EST. NO. 655-1057 E.P.A. Reg. No. 655-536

ACCEPTED

NET CONTENTS _____ LOT # _____

Manufactured by

prentiss Plant: 338 Wilson Avenue
Newark, N.J. 07105
Drug & Chemical Co., Inc. 65-576 Office: 363 7th Avenue
New York, N.Y. 10001

GENERAL OPERATING INSTRUCTIONS:

(For specific operation instructions, refer to Instruction Booklet furnished with machine)

1 Position machine at farthest point from machine is to be moved during treatment, or end of the warehouse where volume is 100

2 Direct nozzle(s) upward and pointed directly toward any nearby object

3 Set output by adjusting flow to desired rate

4 Start machine and run for calculated time to be moved around area during treatment, pre-calculated path at pre-determined speed

5 When treatment time has been completed, flow by shutting off valve. Wait until spray stops machine from area

6 Wait six hours after completion of treatment

ANTIDOTE & FIRST AID TREATMENT: (CONTINUED)

INTERNAL: If the material has been swallowed, immediately. This may be done by introducing throat or by giving warm salt water (1 tablet of water) NEVER GIVE ANYTHING BY UNCONSCIOUS PERSON

EXTERNAL: If the material has been spilled, immediately remove patient from the vicinity. Remove all contaminated clothing, and wash with running water. If the material gets into the eyes, with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer Atropine tablets immediately. NEVER ADMINISTER UNLESS WARNING SYMPTOMS APPEAR (SEE SYMPTOMS below).

Keep patient prone and quiet. Start immediately if patient is not breathing. Immediately to the nearest Physician.

NOTE TO PHYSICIAN —

WARNING SYMPTOMS: Symptoms include tightness in chest, blurred vision, non-responsiveness, sweating, nausea, vomiting, and cramps

TREATMENT: Atropine is the specific choice against parasympathetic nervous system signs of parasympathetic stimulation. Atropine injected at 10-minute intervals in doses until complete Atropinization has occurred. Atropine may be administered in conjunction with Morphine is contraindicated. Clear chest. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors, warning, cause prolonged susceptibility to any cholinesterase inhibitor. Allow no further use of cholinesterase regeneration has been by blood test

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide Spray cannot be used or chemically processed in a landfill approved for pesticides or buried from water supplies.

CONTAINER DISPOSAL: 30 & 55 gallon container and offer for reconditioning, equivalent) and offer for recycling, recycling in approved landfill, or bury in a safe place.

SMALLER CONTAINERS: as in (b) above.

GENERAL: Consult federal, state or local for approved alternative procedures.



prentox[®] D.D.V.P. FIVE

**FOR USE BY PEST CONTROL OPERATIONS OR
TRAINED PERSONNEL ONLY
FOR USE IN WAREHOUSES, RAILROAD CARS
AND STATIONERY TRUCKS, FLOUR MILLS,
GRAIN MILLS, RICE MILLS.**

**AUTHORIZED BY U.S.D.A. FOR USE IN
FEDERALLY INSPECTED MEAT AND POULTRY
ESTABLISHMENTS.**

ACTIVE INGREDIENTS:

2,2-dichlorovinyl dimethyl phosphate*	4.65%
Related Compounds	0.35%
Petroleum Distillates	15.00%
INERT INGREDIENTS:	80.00%
TOTAL:	100.00%

*Equivalent to 5% Vapona Insecticide

**KEEP OUT OF REACH OF CHILDREN
WARNING**

**PRECAUTIONARY STATEMENTS
HAZARDOUS TO HUMANS**

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN OR EYES. DO NOT GET ON SKIN OR IN EYES. DO NOT BREATHE FUMES. WEAR NATURAL RUBBER GLOVES, WATER-PROOF PROTECTIVE CLOTHING AND GOGGLES WHEN HANDLING. WEAR A MASK OR RESPIRATOR OF A TYPE PASSED BY THE U.S. BUREAU OF MINES FOR PROTECTION AGAINST THIS MATERIAL. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. WASH HANDS, ARMS, AND FACE THOROUGHLY WITH SOAP AND WATER BEFORE EATING OR SMOKING. WASH ALL CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE REUSE. NOT FOR USE OR STORAGE IN OR AROUND THE HOME.
ANTIDOTE & FIRST AID TREATMENT:
ATROPINE is the emergency antidote. Consult your Physician about obtaining an adequate supply of 1/100-grain Atropine tablets for emergency use. **CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING**

E.P.A. EST. NO. 655-1001 E.P.A. Reg. NO. 655-536

ACCEPTED

NET CONTENTS LOT #

Manufactured by

prentiss Plant 338 Wilson Avenue
Newark, N.J. 07105
Drug & Chemical Co., Inc. Office: 363 7th Avenue
New York, N.Y. 10001

GENERAL OPERATING INSTRUCTIONS:

(For specific operation instructions, refer to Manufacturer's Instruction Booklet furnished with machine)

- 1 Position machine at farthest point from intended exit, if machinery to be moved during treatment, or in center aisle at one end of the warehouse where volume is 100,000 cu ft or less
- 2 Direct nozzle(s) upward and pointed down the aisle, not directly toward any nearby object
- 3 Set output by adjusting flow to desired level
- 4 Start machine and run for calculated time period. If machine is to be moved around area during treatment, begin backing along pre-calculated path at pre-determined speed
- 5 When treatment time has been completed, stop insecticide flow by shutting off valve. Wait until spray stops and then remove machine from area
- 6 Wait six hours after completion of treatment before ventilating

ANTIDOTE & FIRST AID TREATMENT: (Cont'd.)

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON**

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes

IF WARNING SYMPTOMS APPEAR Administer two 1/100-grain Atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR** (See WARNING SYMPTOMS below)

Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest Physician

NOTE TO PHYSICIAN -

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, Atropine Sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete Atropinization has occurred. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

Morphine is contraindicated. Clear chest by postural drainage. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test

STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited

PESTICIDE DISPOSAL: Pesticide, Spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies

CONTAINER DISPOSAL: 30 & 55 gallon drums: (a) Reseal container and offer for reconditioning or (b) triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place

SMALLER CONTAINERS: as in (b) above

GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures

2 80

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

Law to use this product in a manner

ng specially formulated Insecticide control of exposed stages of stored-grain Warehouses containing non-perishable means any raw or to rapid decay or deterioration that consumption.

include cocoa beans, field-dried beans, and nuts. Not included are eggs, milk, and vegetables such as onions, carrots.

include flour, sugar, cereal, packaged and included are hermetically sealed foods products and other processed foods

DIRECTION:

exposed stages and the prevention of stored-product insects, rice weevils, saw-toothed grain beetles, cigarette moths and cocoa bean moths in Warehouse processing areas of flour and grain and bottling plants and brewery when the plant is not in operation and posed. If bulk, unpackaged foods are before spraying begins. Food should during treatment. All food processing during treatment or thoroughly when using the product in these areas is not in operation. Use DDVP FIVE in to deliver small particles. The rate between 1-2 grams of actual DDVP per of liquid per 1,000 cu. ft. Check amount in tank before and after treatment to been applied.

se during the hours when buildings are sealed should be closed and ventilation application of the Insecticide. Lock all allowed unprotected workers to enter


entrances stating "This building should to re-entry of unprotected workers, or protected workers for at least 6 hours enter prior to o'clock

Repeat treatment at weekly intervals insect activity for optimum results

CONTACT SPRAY - Dilute to 1-3% and of spray per 64,000 cu ft to kill waterbugs, earwigs, ants, spiders, in st a ULV spraying device to a coarse above listed crawling insects. Contact as the spray as possible. Thoroughly spray of harboring these insects. Special to cracks, hidden surfaces under age areas. Then adjust the ULV device to as a space spray. Sweep up and destroy ment as necessary

CONTACT SPRAY - Dilute to 1% and spray 1 pint 1,000 square feet of floor space

duct areas of food processing plants. where food is commercially prepared in serving area while food is exposed



133