

ACCEPTED

prentox

® - FEB 10 1978

MALATHION W-25

25% WETTABLE POWDER

COMPOSITION

ACTIVE INGREDIENT:	
* Malathion	25%
INERT INGREDIENTS	75%
TOTAL	100%
*O,O — diethyl dithiophosphate of diethyl mercaptosuccinate	

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED.
 AVOID PROLONGED BREATHING OF DUST OR SPRAY MIST.
 AVOID REPEATED OR PROLONGED CONTACT WITH SKIN: WASH THOROUGHLY AFTER USING.
 AVOID CONTAMINATION OF FEED AND FOODSTUFFS.
 HIGHLY TOXIC TO FISH. DO NOT CONTAMINATE ANY BODY OF WATER, BY DIRECT APPLICATION, CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES AND CONTAINERS.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Burn or bury containers; never re-use. Stay out of smoke and fumes.

SEE BACK PANEL FOR ADDITIONAL DIRECTIONS FOR USE.

LBS. NET WEIGHT

EPA REGISTRATION NO. 655-549



Manufactured by
prentiss
 Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
 Newark, N.J. 07105
 Office: 363 7th Avenue
 New York, N.Y. 10001

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PEST CONTROL REGISTERED
 UNDER NO. 655-549

prentox[®]

MALATHION W-25

DIRECTIONS FOR USE

CROP	PESTS	DOSAGE PER 100 Gal. Water	REMARKS
APPLES	Aphids	2 to 2½ lbs.	Make one or more applications as needed. Use at least 2 applications 10 to 12 days apart during summer months. Do not apply later than 3 days before harvest.
	* Spider Mites	2 to 2½ lbs.	
PEARS	* Pear Psylla	2 to 2½ lbs.	Make one or more applications as needed. Use at least 2 applications 10 to 12 days apart during summer months. Do not apply later than 1 day before harvest.
	* Spider Mites	2 to 2½ lbs.	
ORNA- MENTALS	Black Scale Crawlers	6 lbs.	Make a thorough full coverage spray after the crawlers have settled on the foliage.
	Soft Brown Scale	6 lbs.	
POTATOES	Monterey Pine Scale	6 lbs.	Apply as needed.
	Juniper Scale	6 lbs.	
	Fletcher Scale	3 lbs.	
	Aphids, Thrips, Lygus, Whitefly * Spider Mites, Mealy Bugs, Leafhopper	2½ lbs.	
	Aphids	2½ lbs.	
CABBAGE, BROCCOLI, KALE, MUSTARD, TURNIPS	Aphids	2 lbs.	Make no application within 7 days of harvest on cabbage, kale and mustard. Make no application within 3 days of harvest on broccoli or turnips.
	Cabbage Looper Cabbage Worm		
TOMATOES	Aphids	2 lbs.	Apply as needed. Make no applications on tomatoes later than 1 day before harvest.
	* Spider Mites		
BEEF CATTLE	Hom Fly	16 to 24 lbs.	Apply as a complete coverage spray. Repeat applications at two-week intervals if needed. Do not apply to dairy animals. Do not treat calves under one month of age. Do not contaminate feed or drinking water.

*MALATHION W-25 is not suggested for control of phosphate-resistant mites or pear psylla.

SEE FRONT PANEL FOR PRECAUTIONARY STATEMENTS.



Manufactured by
prentiss
 Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
 Newark, N.J. 07105
 Office: 363 7th Avenue
 New York, N.Y. 10001

655-549

02/10/1978

11