

# prentox

5% MALATHION DUST

|                    |        |        |
|--------------------|--------|--------|
| ACTIVE INGREDIENT: |        |        |
| Malathion *        | -----  | 5.0%   |
| INERT INGREDIENTS: | -----  | 95.0%  |
|                    | TOTAL: | 100.0% |

**ACCEPTED**

FEB 10 1978

UNDER FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 655-551

655-551

\* 0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

## DIRECTIONS FOR USE:

5% MALATHION DUST may be applied with air and ground equipment when air is calm. Application should be made covering the underside of the leaves as well as the top. The usual rate of application for the following crops is between 20 to 30 pounds per acre.

**BEANS** - For the control of Mexican bean beetle, aphids, leaf hoppers - make 2 or more applications as needed with no application later than 1 day before harvest.

**PEAS** - For the control of pea aphid, pea weevil on peas - apply to foliage during early bloom, repeat as necessary with no application later than 3 days before harvest.

**POTATOES** - For the control of aphids, leafhoppers - apply on foliage of infested plants and repeat as necessary.

**BROCCOLI, CABBAGE, KALE, TURNIP, AND MUSTARD GREENS** - For the control of aphids, imported cabbage worms, cabbage looper - apply on foliage and repeat as necessary with no application later than 3 days before harvest for broccoli and turnips and no later than 7 days before harvest for cabbage, kale and mustard greens.

**CATTLE** - For the control of horn flies - apply on back and neck of the animal 3 tablespoons of 5% MALATHION DUST per animal. Repeat at 10-14 day intervals. Do not apply to dairy cattle later than 5 hours before milking or during milking.

DO NOT CONTAMINATE FEED, FOOD CONTAINERS OR WATER TROUGHS.

FURN OR BURY CONTAINERS; NEVER REUSE. STAY OUT OF SMOKE AND FUMES.

## CAUTION

**KEEP OUT OF REACH OF CHILDREN**

AVOID PROLONGED BREATHING OF DUST. AVOID PROLONGED OR REPEATED CONTACT WITH SKIN; WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

HIGHLY TOXIC TO FISH. DO NOT CONTAMINATE ANY BODY OF WATER NOR APPLY WHEN CONDITIONS FAVOR DRIFT FROM THE CROP AREA TREATED. THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT OR RESIDUES ON CROPS. PROTECTION INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE.

NET CONTENTS \_\_\_\_\_ LBS.

E.P.A. Reg. No. 655-551

02/10/1978



Manufactured by

# prentiss

Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue  
Newark, N.J. 07105

Office: 363 7th Avenue  
New York, N.Y. 10001

1/1