

PRECAUTIONARY STATEMENTS
Hazards to Humans
(and Domestic Animals)
CAUTION

AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN, EYES AND CLOTHING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. WASH THOROUGHLY AFTER HANDLING. HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. TREAT LAWNS IN LATE AFTERNOON. KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL THE FOLLOWING AFTERNOON.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply where runoff is likely to occur. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE—Keep out of reach of children, pets and other animals. Avoid moisture. Avoid contamination of feed or foodstuffs by storage or disposal. Open burning and dumping prohibited. Do not reuse empty container.

DISPOSAL—Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, state, or local disposal authorities.

CONTAINER DISPOSAL—Do not reuse empty container. Wrap container and put in trash collection.

OVERALL LAWN APPLICATION:

Prentox Diazinon 5G controls the following lawn insects on both grass and St. Augustine lawns.

CHINCH BUGS, EARWIGS, ANTS, LAWN MOTHS (SOD WEBWORMS), BERMUDA MITES, CRICKETS, ARMYWORMS, BROWN DOG TICKS, CHIGGERS, CLOVER MITES, CUTWORMS, FLEAS, LEAFHOPPERS, SOWBUGS, MILLIPEDES, AND SPRINGTAILS (COLLEMBOLA).

Apply Prentox Diazinon 5G to your lawn with a fertilizer spreader. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary.

For application with other types of equipment, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20 by 50) or 3.2 oz. per 100 sq. ft. (an area 10 by 10). For heavy lawns such as St. Augustine lawns, application should be increased to 3 lbs. per 1000 sq. ft. (or 4.8 oz. per 100 sq. ft.)

PRENTOX®
Diazinon® 5G

For control of: Ants, armyworms, bermuda mites, billbugs, brown dog ticks, chigger, chinch bugs, clover mites, crickets, cutworms, digger wasps, earwigs, fleas, lawn moths, leafhoppers, millipedes, springtails, sod webworms, sowbugs, white grubs of: Japanese beetle, european chafer, southern chafer.

ACTIVE INGREDIENT:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 5.0%

INERT INGREDIENTS: 95.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED—Call a physician immediately.
IF INHALED—Remove victim to fresh air and apply respiration if indicated.
IF ON SKIN—Wash promptly with soap and water. Rinse thoroughly.
IF IN EYES—Close eyes for at least 15 minutes with water and call a physician immediately.
NOTE TO PHYSICIAN: 2-PAM is antidotal and may be administered in conjunction with atropine sulfate.

E.P.A. Reg. No.: 655-556
 E.P.A. Est. No.: 655 NJ-1

NET CONTENTS: _____ Lbs.

PRENTOX Registered trademark of Prentiss Drug & Chemical Co., Inc.
 Diazinon Registered trademark of CIBA-GEIGY

Manufactured by

Drug & Chemical Co., Inc.

**Plant: 338 Wilson Avenue
 Newark, N.J. 07105
 Office: 363 7th Avenue
 New York, N.Y. 10001**

FOR CONTROL OF BILLBUGS, AND WHITE GRUBS OF: Japanese beetle, european chafer and southern chafer:
 Apply 2.5 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

FOR CONTROL OF HYPERODES WEEVIL (New York only):
 Apply 2 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

SPOT TREATMENT

Ants: Apply 1 1/2 teaspoons of Prentox Diazinon 5G in and around each individual ant hill.

Digger wasps: Apply 1 1/2 teaspoons of Prentox Diazinon 5G in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundation of houses as well as the lawn. Certain of these insects may even enter the house.

ANTS, BROWN DOG TICKS, CLOVER MITES, CRICKETS, EARWIGS, FLEAS, MILLIPEDES, SOWBUGS, SPRINGTAILS (COLLEMBOLA).

For control of the above, apply a five-foot band of Prentox Diazinon 5G next to the foundation of the house. Distribute the granules uniformly using a rate of 3.2 oz. for each 6' by 20' area to be treated.

NOTE: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting if necessary in order to assure correct rate of application.

SPREADER Make and Model	SPREADER SETTING NO. (per 1000 sq. ft.)		
	FOR INSECTS LISTED BELOW* Grass and St. Augustine Lawns 1 - 4 lb.	St. Augustine Lawn 3 lbs.	White Grubs and Billbugs only 2 1/2 lbs.
Central Imperial Mark No. 1662 G	6 1/2	7 1/4	7
Central Imperial No. 1622-0	5	7 1/2	7
Cyclone Model B1 Rotary	4 1/2	5	4 1/4
Greenfield Model 105	6 1/2	7 1/4	6 1/4
J.C. Penney Model 6005	7 1/4	8	7 1/2
J.C. Penney Model 35	5	7	6 1/2
Scott Model 35	4	6	4 1/2
Sears Craftsman Model 671-19190	6	7	6 1/2

* Ants, Armyworms, Bermuda Mites, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Earwigs, Fleas, Hyperodes Weevil (New York only), Lawn Moths (Sod Webworms), Leafhoppers, Millipedes, Sowbugs, Springtails (Collembola), White Grubs.

ACCEPTED

OCT 15 1979

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under E.P.A. Reg. No. 655-556

