

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

FIELD AND FORAGE INSECTS

BEANS (PINTO) Lesser Cornstalk Borers

CORN Corn Rootworm Larvae

Lesser Cornstalk Borers

Suggested Rates According to Corn Row Spacing

Table with 3 columns: Row spacing, Linear feet of corn row per acre, Amount of Diazinon 14G per acre. Rows include 18, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90, 96, 102, 108, 114, 120, 126, 132, 138, 144, 150, 156, 162, 168, 174, 180, 186, 192, 198, 204, 210, 216, 222, 228, 234, 240, 246, 252, 258, 264, 270, 276, 282, 288, 294, 300.

European Corn Borers

Fall Armyworms

Southern Corn Borers

Cutworms

Seed Corn Maggots

Garden Symphylans (Centipedes)

Wireworms

PEANUTS Southern Corn Rootworms

Cutworms

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

Lesser Cornstalk Borers

PRENTOX DIAZINON® 14G

ACTIVE INGREDIENT: O,O-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 14.3%
INERT INGREDIENTS 85.7%
TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED—Call a physician immediately
IF INHALED—Get away from the area to fresh air and apply respiration if needed
IF ON SKIN—Wash promptly with soap and water. Rinse thoroughly
IF IN EYES—Rinse eyes for at least 15 minutes with water and call a physician immediately
NOTE TO PHYSICIAN: 2-PAM is antidotal and may be administered in conjunction with atropine sulfate

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (& DOMESTIC ANIMALS) CAUTION

HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FOOD AND FEED. DO NOT STORE NEAR FOOD OR FEED PRODUCTS. AVOID INHALATION OF DUST. AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WASH THOROUGHLY AFTER HANDLING. MAY BE ABSORBED THROUGH SKIN.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL:

CONTAINER DISPOSAL: Do not reuse empty containers. Wrap container and put in trash collection
STORAGE: Keep out of reach of children, pets and other animals. Avoid moisture. Avoid contamination of food and foodstuffs by storage or disposal. Open burning and dumping prohibited. Do not reuse empty containers.

DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or local disposal authorities.

SEE SIDE PANEL FOR OTHER PRECAUTIONS.

EPA EST. NO. 675-NJ-1

EPA REG. NO. 655

NET CONTENTS _____ lbs.

Manufactured by

Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue Newark, N.J. 07105

Office: 363 7th Avenue New York, N.Y. 10001

TOBACCO Cutworms: Broadcast 14-18 lbs per acre before transplanting. Work into soil 2-4 inches
Mole Crickets: Apply 7 lbs per acre when insects are present. Make application to plant beds or rows 7-10 days before seeding or transplanting and work into soil 1-2 inches

Seed Corn Maggots: Broadcast 14-18 lbs per acre 4 weeks before transplanting. Work into soil 2-4 inches

Wireworms: Broadcast 14-18 lbs per acre 4 weeks before transplanting. Work into soil 2-4 inches

FORAGE CROPS (Alfalfa, Clover, Cowpeas, and Leppeds) Cutworms: Broadcast 14-18 lbs per acre before planting. Work into soil 2-3 inches

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, AND CAULIFLOWER Root Maggots: Broadcast 14-21 lbs per acre before planting or transplanting. Work into soil 2-4 inches

ONIONS Onion Maggots: Apply 14-28 lbs per acre as a broadcast soil treatment just prior to planting. Thoroughly mix the chemical into the top 3-4 inches of soil by disk or harrow or other suitable means.

RADISHES Root Maggots: Apply 10-14 lbs per acre (2-24.5 oz per 1,000 linear feet of row) in 1-2 inch furrows at planting time.

SWEET CORN Wireworms: Broadcast 14-18 lbs per acre before planting. Work into soil 2-4 inches

TOMATOES Vinegar Flies (Drosophila spp.): Broadcast 14-21 lbs per acre 7-10 days before planting. Work into soil 2-4 inches

OTHER VEGETABLE INSECTS

CUTWORMS (Surface and Subterranean) Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Endive, Kale, Lettuce, Lima Beans, Mustards, Parsley, Peas, Peppers, Pole Beans, Snap Beans, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash

Endive (escarole), Kale, Lettuce, Lima Beans, Mustards, Parsley, Peas, Peppers, Pole Beans, Snap Beans, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash

GARDEN SYMPHYLANS (Centipedes) Cutworms, Endive, Kale, Lettuce, Lima Beans, Mustards, Parsley, Peas, Peppers, Pole Beans, Snap Beans, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash

MOLE CRICKETS Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Endive, Kale, Lettuce, Lima Beans, Mustards, Parsley, Peas, Peppers, Pole Beans, Snap Beans, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash

WIREWORMS Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Endive, Kale, Lettuce, Lima Beans, Mustards, Parsley, Peas, Peppers, Pole Beans, Snap Beans, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

Lesser Cornstalk Borers: Apply 14-18 lbs per acre before planting. Work into soil 2-4 inches

