

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds and other wildlife. Birds feeding in treated areas may be killed. Do not apply where runoff is likely to occur. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

STORAGE—Keep out of reach of children, pets and other animals. Avoid moisture. Avoid contamination of feed or foodstuffs by storage or disposal. Open burning and dumping prohibited. Do not reuse empty container

DISPOSAL—Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State, or Local disposal authorities.

CONTAINER DISPOSAL—Do not reuse empty container. Wrap container and put in trash collection.

OVERALL LAWN APPLICATION:

Prentox Diazinon 2G is for use on the following insects on bermuda grass and Dichondra lawns:

CHINCH BUGS, EARWIGS, ANTS, LAWN MOTHS (SOD WEBWORMS), BERMUDA MITES, CRICKETS, ARMYWORMS, BROWN DOG TICKS, CHIGGERS, CLOVER MITES, CUTWORMS, FLEAS, LEAFHOPPERS, SOWBUGS, MILLIPEDES AND SPRINGTAILS (COLLEMBOLA).

Apply Prentox Diazinon 2G to your lawn with a fertilizer spreader at the rate shown on the side panel chart. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary.

For application with other types of equipment, apply at the rate of 5 lbs. per 1000 sq. ft. (an area 20' by 50' per 8 oz. per 100 sq. ft. (an area 10' by 10'). For heavy lawns, apply at the rate of 10 lbs. per 1000 sq. ft. (an area 20' by 50').

TO CONTROL BILLBUGS, WHITE GRUBS OF JAPANESE BEETLE, EUROPEAN CHAFER AND SOUTHERN CHAFER:

Apply 8 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application for grub control make application any time between late July and October. For beetle control, apply when adults are first observed or when chewed or damaged areas are first observed. Repeat as necessary.

TO CONTROL HYPERODES WEEVIL (New York only):
Apply 5 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after. Treat problem areas in mid-April and again in mid-May.

**PRENTOX®
DIAZINON® 2G**

For control of: Ants, armyworms, bermuda mites, billbugs, brown dog ticks, chiggers, chinch bugs, clover mites, crickets, cutworms, digger wasps, earwigs, fleas, lawn moths, leaf hoppers, millipedes, springtails, sod webworms, sowbugs, white grubs of: Japanese beetle, european chafer, southern chafer.

ACTIVE INGREDIENT:

(1) O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 2.0%
INERT INGREDIENTS 98.0%
TOTAL 100.0%

PRENTOX® Registered Trademark, Pontiss Drug & Chemical Co. Inc.
DIAZINON® Registered Trademark of CIBA-GEIGY

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans
(and Domestic Animals)

CAUTION

AVOID BREATHING OF DUST; AVOID CONTACT WITH SKIN, EYES AND CLOTHING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. WASH THOROUGHLY AFTER HANDLING. HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. TREAT LAWNS IN LATE AFTERNOON. KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL THE FOLLOWING AFTERNOON.

See side panel for additional precautionary statements.

E.P.A. Reg. No.: 655-559
E.P.A. Est. No.

NET CONTENTS: _____ Lbs.

Manufactured by

Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
Newark, N.J. 07105

Office: 363 7th Avenue
New York, N.Y. 10001

SPOT TREATMENT

Ants: Apply one tablespoon of Prentox Diazinon 2G over and around each individual ant nest.

Digger Wasps: Apply one tablespoon of Prentox Diazinon 2G in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION:

The following insects are commonly found around or near the foundation of houses and well as in the lawn. Certain of these insects may even enter the house.

ANTS, BROWN DOG TICKS, CLOVER MITES, CRICKETS, EARWIGS, FLEAS, MILLIPEDES, SOWBUGS, SPRINGTAILS (COLLEMBOLA).

For control of the above insects, apply a five-foot band of Prentox Diazinon 2G next to the foundation of the house. Distribute the granules uniformly at a rate of 8 oz. for each 5' by 20' area to be treated.

NOTE: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting if necessary in order to assure correct rate of application.

SPREADER Make and Model	SPREADER SETTING NO. (per 1000 sq. ft.)		
	FOR INSECTS LISTED BELOW* Grass and Dichondra Lawn 5 lbs.	St. Augustine Lawn 7 1/2 lbs.	White Grubs and Billbugs only 8 lbs.
Central Imperial Mark No 1662-G	7 1/2	8 1/2	8
Central Suburban Lawn Spreader—No. 1622-0	7 1/2	8 1/2	8
Cyclone Model B1 Rotary	6	7 1/2	6 1/2
Greenfield Model 105	8	8 1/2	8 1/2
J.C. Penney Model 6005	8 1/2	9 1/2	9
J.C. Penney Rotary Spreader	8	10	9
Scott Model 35	6 1/2	7 1/2	7
Scars Craftsman Model 671-19198	7 1/2	9	8

Ants
Armyworms
Bermuda Mites
Brown Dog Ticks
Chiggers
Chinch Bug
Clover Mites
Crickets
Earwigs
Fleas
Hyperodes Weevil (New York only)
Lawn Moths (Sod Webworms)
Leafhoppers
Millipedes
Sowbugs
Springtails (Collembola)

ACCEPTED
OCT 15 1979
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, no pesticide is to be registered unless it meets the requirements of the Act.
655-559



BEST DOCUMENT AVAILABLE