

# prentox<sup>®</sup>

## DIAZINON 2G

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS  
CAUTION**

Avoid breathing of dust. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling. Harmful if swallowed. May be absorbed through skin. Treat lawns in late afternoon. Keep children and pets off treated areas until the following afternoon.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds and other wildlife. Birds feeding in treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**DIRECTIONS FOR USE:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**STORAGE:** If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup, wear protective equipment as required to prevent contact with the product. Sweep on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bag in a sanitary landfill or by incineration if allowed by State and local authorities.

**OVERALL LAWN APPLICATION:**

Controls the following lawn insects on both grass and Dichondra lawns.

**CHINCH BUGS, EARWIGS, ANTS, LAWN MOTHS (SOD WEBWORMS), BERMUDA MITES, CRICKETS, ARMYWORMS, BROWN DOG TICKS, CHIGGERS, CLOVER MITES, CUTWORMS, FLEAS, LEAFHOPPERS, SOWBUGS, MILLIPEDES AND SPRINGTAILS (COLLEMBOLA).**

Apply Prentox Diazinon 2G to your lawn with a fertilizer spreader. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary.

For application with other types of equipment apply at the rate of 5 lbs per 1000 sq ft (an area 20' by 50') or 8 oz per 100 sq ft (an area 10' by 10'). For heavy lawns such as St. Augustine lawns, application should be increased to 7.5 lbs per 1000 sq ft (or 12 oz per 100 sq ft).

**For control of: Ants, armyworms, bermuda mites, billbugs, brown dog ticks, chigger, chinch bugs, clover mites, crickets, cutworms, digger wasps, earwigs, fleas, lawn moths, leafhoppers, millipedes, springtails, sod webworms, sowbugs, white grubs of: Japanese beetle, european chafer, southern chafer.**

**ACTIVE INGREDIENT:**

Diazinon -(0, 0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate) ..... 2.0%  
**INERT INGREDIENTS:** ..... 98.0%  
**TOTAL** ..... 100.0%


**PRENTOX<sup>®</sup>** — Registered Trademark of Prentiss Drug & Chemical Co., Inc.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

E.P.A. REG. NO.: 655-559

E.P.A. EST. NO.

NET CONTENTS 50 LBS.



Manufactured by



**prentiss**  
Drug & Chemical Co., Inc.

Plant: Kaolin Road  
Sandersville, GA. 31082

Office: C.B. 2000  
Ficral Park, N.Y. 11001

**TO CONTROL BILLBUGS, WHITE GRUBS OF JAPANESE BEETLE, EUROPEAN CHAFER AND SOUTHERN CHAFER:**

Apply 6 lbs per 1000 sq ft of lawn. Water grass thoroughly after application. For grub control make application on any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

**TO CONTROL HYPERODES WEEVIL (New York only)**

Apply 5 lbs per 1000 sq ft of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

**SPOT TREATMENT**

**Ants:** Apply one tablespoon of this product over and around each individual ant hill.

**Digger Wasps:** Apply one tablespoon of this product around each individual ground nest opening. Make the application about dusk when the wasps are not active.

**BAND TREATMENT AROUND FOUNDATION.**

The following insects are commonly found around or near the foundation of houses as well as the lawn. Certain of these insects may even enter the house.

**ANTS, BROWNDOG TICKS, CLOVER MITES, CRICKETS, EARWIGS, FLEAS, MILLIPEDES, SOWBUGS, SPRINGTAILS (COLLEMBOLA).**

For control of the above insects apply a five-foot band of this product next to the foundation of the house. Distribute the granules uniformly using a rate of 8 oz for each 5' by 20' area to be treated.

**NOTE:** The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting if necessary in order to assure correct rate of application.

**SPREADER SETTING NO.  
(per 1000 sq. ft.)**

| SPREADER Make and Model                         | FOR INSECTS LISTED BELOW*<br>Grass and Dichondra Lawn<br>2 lbs. | St. Augustine Lawn<br>3 lbs. | White Grubs and Billbugs Only<br>2 1/2 lbs. |
|---|---|------------------------------|---|
| Central Imperial<br>Mark No. 1662-G             | 7   | 8                            | 8   |
| Central Superior<br>Lawn Spreader<br>No. 1622-D | 7   | 8                            | 8   |
| Cyclone<br>Model B1<br>Rotary                   | 6   | 7                            | 6   |
| Greenfield<br>Model 105                         | 8   | 8 1/2                        | 8 1/2                                       |
| J.C. Penney<br>Model 600A                       | 8   | 9                            | 9   |
| J.C. Penney<br>Rotary<br>Spreader               | 8   | 10                           | 9   |
| Scott<br>Model 35                               | 6   | 7                            | 7   |
| Sears Craftsman<br>Model 621-19128              | 7   | 9                            | 8   |
| Ants  |   |                              | Earwigs                                     |
| Armyworms                                       |   |                              | Fleas                                       |
| Bermuda Mites                                   |   |                              | Hyperodes Weevil (New York only)            |
| Brown Dog Ticks                                 |   |                              | Lawn Moths (Sod Webworms)                   |
| Chiggers  |   |                              | Leafhoppers                                 |
| Chinch Bugs                                     |   |                              | Millipedes                                  |
| Clover Mites                                    |   |                              | Sowbugs                                     |
| Crickets  |   |                              | Springtails (Collembola)                    |
| Cutworms  |   |                              |   |