PRECAUTIONARY STATEMENTS HAZAKDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION

Harmful if swallowed. Avoid breathing dust. May be absorbed through skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep out of reach of children,

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.


This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not reuse empty container. Wrap container & place in trash collection.



prentox

DIAZINON[®] **4**D

ACTIVE INGREDIENT:

O,O-diethyl O-(2-isopropyl-6-methyl-4pyrimidinyi) phosphorothicate 4.0% TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED -- Call a physician immediately. IF INHALED - Remove victim to fresh air and apply respiration, if indicated.

IF ON SKIN - Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES -- Rinse eyes for at least 15 minutes with water and call a physician immediately NOTE TO PHYSICIAN: Atropine sulfate is antidotal and may be

administered in conjuction with 2-PAM SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.


PRENTOX8 - Registered Trademark of Prentiss Drug & Chemical Co., Inc.

E.P.A Reg. No. 655-560

NET CONTENTS_

LOT#_





DIRECTIONS FOR USE (Could.)

Soli Application: To control cutworms and wireworms prior to planting of the following vegetables:

Beans Broccol: Cabbage Carrots Caut figuer	Corn Cucumbers Endive Lettuce Metons	Onions Parships Peus Peppers Potatoes	Spinact Squash Tomatoes
--	--	---	-------------------------------

Apply evenly as a broadcast treatment over the surface of the soil. Immediately after spreading, mix thoroughly with the soil to a depth of 6-8 inches. Apply ½ lb. per 225 sq. ft.

Foliage Application: Apply dust at first signs of insects. Repeat as necessary. For best results, dust in early morning or late evening when temperature is lower and air is still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. Number in parentheses, (), indicates days required between last application and harvest.

Planta	Insects
Beans (7)	Aphids Mexican Bean Beetles Leefhoppers
Peas (0)	Aphids
Broccoli (5) Cabbage (7) Cautifower (5)	Aphids Diamondback Moths Imported Cabbageworm
Cucumber (7) Melons Canta- loupe Muskmelo., Watermelon (3), Squash Summer (7) Winter (3)	Aphids Easthoppers Serpentine Lea'miners, Miles Thrips
Peppers (5)	Aphids Serpentine Leafminers
Tomatoes (3)	Aphids Dipterous Leafminers Banded Cucumber Beetles, Fatt Armyworm Southern Armyworm
Sweet Corn (2)	Com Earworm*
Strawbernes (5)	Aphids Twospotted Mites
Azaleas Camellias, Carnations Chrysan themums, Gladiola Junipers Lifacs, Rhododendron Roses	Aphids, Mites, Leathopper, Meelybugs, Whitelines, Scale Crewlers, Lacubuge

Note: Do not use on Ferns, Poinsetties, Hibiscus since injury may occur.

Apply to silks as soon as they appear. Make applications at 3 to 4 day intervals as infestation warrants. Reapply immediately after rain

36

7/84

Office: 363 7th Avenue New York, N.Y. 10001