

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION

ACCESSED

FEB 15 1985

prentox

LINDANE 20% EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS:

Lindane (Gamma Isomer of Benzene Hexachloride)

Xylene Range Aromatic Solvent

INERT INGREDIENTS:

(Contains 1.83 pounds of Lindane per gallon) - TOTAL:

20.0%

~~67.0%~~ 62.2%

~~13.0%~~ 17.8%

100.0%

PRENTOX® - Registered Trademark of Prentiss Drug & Chemical Co. Inc.

E.P.A. REG. NO. 655-578

E.P.A. EST. NO. 655-GA-1

NET CONTENTS

LOT NO

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Call a physician - Poison Control immediately. Contains petroleum distillates. Do not induce vomiting. Get medical attention immediately. If inhaled - Remove victim to fresh air. Apply artificial respiration if necessary. If on skin - Remove contaminated clothing and wash area immediately with soap and warm water. If in eyes - Flush with plenty of water and get prompt medical attention.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be absorbed through skin. Harmful if swallowed. Avoid inhalation of spray mist. Avoid skin contact. In case of contact, wash immediately with soap and water. Wash thoroughly after using and before eating or smoking. Avoid contamination of feed and foodstuffs. Food utensils such as teaspoons should not be used for food purposes after use in measuring pesticide. Do not use on household pets and humans. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift to areas treated. Do not use in areas where there is a high concentration of disposal of wastes.

BENEFICIAL INSECT CAUTION

This pesticide is highly toxic to bees, used to control treatment of colonies of bees in the treated area. Do not apply when bees are actively visiting the crop. Cover crop or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AERIAL APPLICATION OF LINDANE IS PROHIBITED

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops being treated with this product, consult your State Department of Agriculture for further information.

Manufactured by



prentiss

Drug & Chemical Co. Inc.

Plant: Kaolin Road
Sandersville, GA. 31082

Office: C.B. 2000
Floral Park, N.Y. 11001

USE TO APPLY:
to ground. Do not apply to fruit or foliage. Do not apply within 80 days of harvest.

PECANS: Tom Control P&A Phylloxera and Short Curculionids 20 ounces 2 1/2 to 10 gallons of water. Thoroughly wet the foliage at approximately 30 gallons of spray per tree. Do not apply more than 450 gallons of finished spray per acre (1 gal. 6 ozs. gallons of this product). Apply by air blast sprayer when the pecan buds are beginning to burst. This is early to mid April. Make only one application. **DO NOT GRAZE TREATED PEAS.** Do not apply within 28 days of harvest.

TOMATOES: Aphids. Use 1 gallon of 0.03% spray per 150 square feet. Do not apply after fruit starts to form.

CABBAGE CELERY COLLARDS, CUCUMBERS, LETTUCE, OKRA, PEPPERS, SQUASH, MELONS, TOMATOES: White Grubs, Cutworms, Wireworms. For preplant soil application only. Use 2 gallons for each 250 square feet. Work into soil thoroughly before or during planting. Do not treat land to be planted in next crops.

BULBS (DAFFODIL, GLADIOLUS, HYACINTH, IRIS, LILIES, NARCISSUS TULIPS): Aphids, Bulb Fly, European corn borer, etc. Apply when insects appear. Repeat as necessary.

FLOWERS (CARNATIONS, CHRYSANTHEMUMS, COLEUS, DAISIES, GARDENIA, GERANIUMS, HIBISCUS, NASTURTIUMS, PANSIES, PETUNIAS, SWEET PEAS, ZINNIA): Aphids, Hooter beetles, Leafhoppers, Thrips. Apply when insects appear. Repeat as necessary.

ORNAMENTALS (AZALEAS, CAMELLIAS, HEATHER, LILACS, RHODODENDRONS, ROSES): Aphids, Blackvine weevil, Cucumber beetles, Thrips, White Fly. Apply when insects appear. Repeat as necessary.

TICKS AND CHIGGERS ON VEGETATION: A minimum dosage of 5 pounds of actual Lindane per acre for control of ticks and chiggers on vegetation. Lawns, recreation areas, campgrounds, golf courses. Mix 1 quart thoroughly in 25 gallons of water and apply at the rate of 20 to 25 gallons per acre or about 1 gallon to 100 square feet. Cylindrical or knapsack type hand sprayers may be used over small areas, but power equipment will be most satisfactory for large acreages. Treat ground first and the lower parts of plants with this spray.

MEASURE ACCURATELY - STIR THOROUGHLY BEFORE USE

DO NOT SPRAY ANIMALS WITHIN 30 DAYS OF SLAUGHTER

HORNFLIES ON BEEF CATTLE: Spray animals thoroughly with 0.3% Lindane solution prepared by adding 1 tablespoonful Lindane Concentrate to 2 1/2 gallons of water or 1/2 pint to 40 gallons. Repeat as necessary. Do not spray calves under 3 months of age.

LICE AND TICKS ON BEEF CATTLE, HORSES, SHEEP, GOATS: Spray animals with 0.03% Lindane solution prepared by adding 1 tablespoonful Lindane Concentrate to 2 1/2 gallons of water or 1/2 pint to 40 gallons. Wet skin thoroughly. Repeat as necessary. Do not spray calves under 3 months of age.

LICE ON HOGS: Spray animals with 0.06% Lindane Solution prepared by adding 2 tablespoonfuls Lindane Concentrate to 2 1/2 gallons of water or 1/2 pint to 40 gallons. Repeat as necessary. Do not apply to young animals under 3 months of age.

SARCOPTIC MANGE ON BEEF CATTLE AND HOGS AND PSOROPTIC MANGE (SHEEP SCABIES) ON SHEEP AND GOATS: Spray animals with 0.03% spray prepared by adding 1 tablespoonful Lindane Concentrate to 2 1/2 gallons of water. Wet animals thoroughly. Use 2 or more treatments at 7 to 10 day intervals as needed. Pens and shelters should also be sprayed to prevent re-infestation. Do not spray calves under 3 months of age.

FLIES, ROACHES, SPIDERS, ANTS, MOSQUITOES IN LIVESTOCK PENS AND BUILDINGS: Spray with a 0.3% solution prepared by adding 4 tablespoonfuls Lindane Concentrate to 1 gallon of water or 1/2 pint to 4 gallons. Apply 1 gallon of mix for each 500 square feet of surface area or until thoroughly wet. Shut doors off. For crawling insects apply thoroughly to cracks, crevices, baseboards and other hiding places. Remove animals before spraying. Do not contaminate milk or milking equipment feed or drinking water.

INSECTICIDAL PAINT: Use 1.25% Lindane solution prepared by adding 1/2 pint Lindane Concentrate to 1 gallon of water. Apply with brush on any and around hiding places as a spot treatment. This concentration may blister paint. Do not use in the home. Do not contaminate milk, milking equipment, feed or drinking water.

BEST AVAILABLE COPY

