

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. DO NOT BREATHE DUST OR SPRAY MIST. AVOID CONTACT WITH SKIN AND EYES. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. DO NOT USE IN POULTRY HOUSES.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding on treated exposed seed may be killed. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

STORAGE & DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE DISPOSAL: Pesticide, Spray Mixture or Rinseate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or Local Disposal Authorities.

CONTAINER DISPOSAL: Consult Federal, State or Local Disposal Authorities for approved procedures.

29

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

For Seed Treatment

ACTIVE INGREDIENTS:

Lindane (Gamma Isomer of Benzene Hexachloride) --- 25%

INERT INGREDIENTS: ----- 75%
TOTAL: 100%

KEEP OUT OF REACH OF CHILDREN

WARNING

Statement of Practical Treatment

- IF SWALLOWED:** Call a physician immediately.
- IF INHALED:** Remove victim to fresh air and apply respiration if indicated.
- IF ON SKIN:** Wash immediately with soap and warm water.
- IF IN EYES:** Flush eyes for at least 15 minutes with water. Call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DIRECTIONS FOR USE: (to

SEED TREATMENT: For protection of seeds and seedlings against wireworm and seedcorn maggot, except where populations are very high, uniform coverage of seeds listed or follow your Station. May be used dry or and may be combined with treating fungicides, or a which already have fungicide. Excessive amount may reduce. Treat seeds within 3 weeks. Treated seed should not be high temperatures as this duction in germination and. Do not use treated seed for purposes. TREATED SEEDS surface will be hazardous other wildlife. NOTE: must not be used or mixed animal feed, or processed commercially treated with must be labeled in accordance with requirements of Seed Act.

CORN: Use 4 to 8 oz./100 (use lowest dosage in N.Y. and highest dosage in California).
LARGE SEEDED LIMA BEANS: of seed.

OTHER PEAS: Use 2 oz./bu. Use protective fungicide on beans.

PEAS: Use 4 oz./ 100 lbs.

SUGARBEETS: Use 16 oz./100 lbs.

COTTON: Use 6 to 8 oz./100 lbs.

SMALL GRAINS (wheat, oats, barley, rye, etc.): Use 3 to 4 oz. on quantity planted per acre. This is usually from 1 1/2 to 4 oz. per acre.

NET CONTENTS

POUNDS

E.P.A. Reg. No. 655-580

E.P.A. EST. NO. 655-580

Manufactured by

PROLIFE
Drug & Chemical Co., Inc.

Plant: 538 Wilson Avenue
Newark, N.J. 07105

Office: 333 7th Avenue
New York, N.Y. 10001

ACCEPTED
1980

655-580

2

LINDANE 20W

For Seed Treatment

ACTIVE INGREDIENTS:
Lindane (Gamma Isomer of Benzene Hexachloride) --- 25%
INERT INGREDIENTS: ----- 75%
TOTAL: 100%

KEEP OUT OF REACH OF CHILDREN
WARNING
Statement of Practical Treatment
IF SWALLOWED: Call a physician immediately.
IF INHALED: Remove victim to fresh air and apply respiration if indicated.
IF ON SKIN: Wash immediately with soap and warm water.
IF IN EYES: Flush eyes for at least 15 minutes with water. Call a physician immediately.
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT (cont'd)
Apply Lindane 20W to a mixture of seed and seedlings against wireworms and seed corn maggot, except when insect populations are very high, apply to give uniform coverage of seed at cost listed or follow your State recommendations. May be used dry or as slurry, and may be combined with certain seed treating fungicides, or applied to seeds which already have fungicide treatment. Excessive amount may reduce germination. Treat seeds within 3 weeks of planting. Treated seed should not be stored at high temperatures as this will cause reduction in germination and emergence. Do not use treated seed for food or food purposes. TREATED SEEDS exposed on soil surface will be hazardous to birds and other wildlife. NOTE: Treated seed must not be used or mixed with food or animal feed, or processed for oil. Seed commercially treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act.

CORN: Use 4 to 8 oz./100 lbs. of seed (use lowest dosage in N.Y. & N.J., medium dosage in California & Southern States, high doses elsewhere).
LARGE SEED LIMA BEANS: Use 1 oz./bu. of seed.
OTHER BEANS: Use 2 oz./bu. of seed. Use protective fungicide with insecticide on beans.
PEAS: Use 4 oz./ 100 lbs. of seed.
SUNFLOWERS: Use 16 oz./100 lbs. of seed.
GRITTS: Use 6 to 8 oz./100 lbs. of seed.
SMALL GRAINS (wheat, oats, barley): Use 3 to 4 oz. on quantity of seed planted per acre. This will vary usually from 1/2 to 4 oz. per bu. of seed.

STATEMENTS HANS & DOMESTIC ANIMALS

IF SWALLOWED. MAY BE AB-
H SKIN. DO NOT BREATHE
INIST. AVOID CONTACT
EYES. AVOID CONTAMI-
D AND FOODSTUFFS. DO
LTRY HOUSES.

HAZARDS

is toxic to fish, birds,
life. Birds and other
ing on treated exposed
illed. Keep out of lakes,
onds. Do not apply where
ely to occur. Do not ap-
her conditions favor
reated areas. Do not con-
y cleaning of equip-
sal of wastes. Apply
only as specified on

USE

ION OF FEDERAL LAW TO
CT IN A MANNER INCON-
TS LABELING.


DISPOSAL

INATE WATER, FOOD OR
GE OR DISPOSAL.

DISPOSAL: Pesticide,
or Rinsate that can-
or chemically reproces-
disposed of according
approved by Federal,
Disposal Authorities.

DISPOSAL: Consult
or Local Disposal
or approved proce-

NET CONTENTS _____ POUNDS
E.P.A. Reg. No. 655-580
E.P.A. EST. NO. 655-580
Manufactured by
MONROVIA Plant: 330 Wilson Avenue
Newark, N.J. 07105
Drug & Chemical Co., Inc. Office: 333 7th Avenue
New York, N.Y. 10001



29A