

**CAUTIONARY STATEMENTS
HARMFUL TO HUMANS & DOMESTIC ANIMALS**

IT MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. DO NOT BREATHE DUST OR SPRAY MIST. AVOID CONTACT WITH SKIN AND EYES. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. DO NOT USE IN POULTRY HOUSES.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding on treated exposed seed may be killed or put out of action, blinded and paralyzed. Do not spray when other conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of mix. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in treatment area. Applications should be timed to provide maximum possible interval between treatment and the next visit of bees to the crop.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

ORAGE & DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE DISPOSAL: Pesticides, spray containers, and other materials that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or Local Health Authorities.

WASTE DISPOSAL: Consult Federal, State or Local Health Authorities for approved procedures.

PLANTS: Aphids - 1 lb. in 100 gals. water (200 to 400 gals. spray per acre). Concentrate Spray - 3 to 4 lbs. per acre in water to cover. Treat under leaves also. Do not apply within 60 days of harvest.

BEANS, BROCCOLI, BRUSSELS SPROUTS, CABBAGE, KALE, COLLARDS: Aphids, Flea Beetle, Cutworms, etc. Lettuce - 1 lb. in 100 gals. water (200 gals. spray per acre). Do not apply after the edible parts start to form. Cabbage - 4 lbs. per acre in water to cover. Apply to soil around base of plants when plants are 2 to 3 inches tall. Repeat at 7 to 10 day intervals.

PEPPERS: Aphids, Leaf Miners - 1 lb. in 100 gals. water (200 gals. spray per acre). Armyworms - 2 1/2 lbs. per acre in water to cover. Do not apply after the edible parts start to form.

SOIL INSECTICIDE: Wireworms, Cutworms, White Grubs, Starvation Larvae on Celery, Cucumbers, Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Strawberries, Lettuce, Lettuce, Spinach, Tomatoes, etc. (Do not harvest in fall prior to winter use) - Pre-Plant Treatment - 1 to 2 lbs. per acre. Apply to soil or dust directly on soil. Disc as soon as possible to give thorough distribution in upper 4 to 6 inches of soil. Can be mixed with acid or neutral fertilizers and applied with fertilizer only. Do not apply to root crops within 1 year. Do not use in soil to be planted to potatoes within 2 years.

SPINACH: Aphids, Leaf Miners - 1 lb. in 100 gals. water (200 gals. spray per acre). Use only in seedling stage.

STORAGE PRODUCTS INSECTICIDE: Strawberry Weevil, San Jose Scale, Pear Scale, Combed Pear Scale, Corn or Rice Weevil, Angoumois Grain Weevil, Indian Meal Weevil (Grain Bin, Storage Bin, Warehouse) - 16 1/2 lbs. per 100 gals. water. Spray bins and warehouses when empty. Prior to spraying clean walls and floor. Apply as a spot treatment. Spray into breeding places, areas where insects congregate or onto insects directly. Repeat as necessary. Controls only susceptible stages of these pests. Apply only so that feedstuffs or grain will not be contaminated.

prentox LINDANE 25W

ACTIVE INGREDIENTS:
Lindane (Gamma Isomer of Benzene Hexachloride) 25%

INERT INGREDIENTS 75%

TOTAL: 100%

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED - Call a physician immediately.
- IF SWALLOWED - Remove victim to fresh air and apply respiration if indicated.
- IF ON SKIN - Wash immediately with soap and warm water.
- IF IN EYES - Flush eyes for at least 15 minutes with water. Call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL DIRECTIONS FOR USE

E.P.A. Reg. No. 688-688 E.P.A. Sol. No. 688-688-1

GRASSHOPPERS: Fence Rows, Barrier Zones, Waste Lands, and Uncultivated Areas. As a dust apply 1.5-2 lbs. per acre directly to the infested area. Where grasshoppers are migrating into field put a 10 foot barrier strip of dust around field or on side from which they are approaching. As a spray apply 7 1/2 to 10 lbs. per 100 gals. water at the same rate per acre as given for the dust application.

GRUBS: Thrips - 1 lb. per 100 gals. water (200 gals. spray per acre). First application when thrips average 5 to 6 per plant. Repeat in 5 to 6 days. A third application may be necessary. Do not use on green or spring onions. Onion Maggot - 4 lbs. per acre in water to cover. Apply to soil around base of plants when plants are 2 to 3 inches tall. Repeat at 7 to 10 day intervals.

PEPPERS: Aphids, Leaf Miners - 1 lb. in 100 gals. water (200 gals. spray per acre). Armyworms - 2 1/2 lbs. per acre in water to cover. Do not apply after the edible parts start to form.

SOIL INSECTICIDE: Wireworms, Cutworms, White Grubs, Starvation Larvae on Celery, Cucumbers, Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Strawberries, Lettuce, Lettuce, Spinach, Tomatoes, etc. (Do not harvest in fall prior to winter use) - Pre-Plant Treatment - 1 to 2 lbs. per acre. Apply to soil or dust directly on soil. Disc as soon as possible to give thorough distribution in upper 4 to 6 inches of soil. Can be mixed with acid or neutral fertilizers and applied with fertilizer only. Do not apply to root crops within 1 year. Do not use in soil to be planted to potatoes within 2 years.

SPINACH: Aphids, Leaf Miners - 1 lb. in 100 gals. water (200 gals. spray per acre). Use only in seedling stage.

STORAGE PRODUCTS INSECTICIDE: Strawberry Weevil, San Jose Scale, Pear Scale, Combed Pear Scale, Corn or Rice Weevil, Angoumois Grain Weevil, Indian Meal Weevil (Grain Bin, Storage Bin, Warehouse) - 16 1/2 lbs. per 100 gals. water. Spray bins and warehouses when empty. Prior to spraying clean walls and floor. Apply as a spot treatment. Spray into breeding places, areas where insects congregate or onto insects directly. Repeat as necessary. Controls only susceptible stages of these pests. Apply only so that feedstuffs or grain will not be contaminated.

Manufactured by



prentiss
Drug & Chemical Co., Inc.

Plant: Keolin Road
Sandersville, GA. 31082
Office: C.B. 206
Floral Park, N.Y. 11001

BEST DOCUMENT

DIRECTIONS FOR USE (Cont'd.)

STRAWBERRIES: Use heavy spray 200 gals. water (200 gals. spray per acre). Use 1 lb. per acre in water to cover. Apply only after the fruit sets in to form.

TOMATOES: Use heavy spray 200 gals. water (200 gals. spray per acre). Use 1 lb. per acre in water to cover. Apply only after the fruit sets in to form.

USES ON ANIMALS: Do not treat animals under 3 months old. Do not treat sheep, goats, or hogs within 60 days of slaughter. Do not treat dairy cattle or deer within 30 days of milking or within 30 days of slaughter. Do not treat swine within 30 days of slaughter. Do not treat poultry within 30 days of slaughter. Do not treat horses within 30 days of slaughter. Do not treat dogs within 30 days of slaughter. Do not treat cats within 30 days of slaughter. Do not treat birds within 30 days of slaughter. Do not treat fish within 30 days of slaughter. Do not treat shellfish within 30 days of slaughter. Do not treat honey bees within 30 days of harvest. Do not treat bees within 30 days of flight. Do not treat bees within 30 days of nesting. Do not treat bees within 30 days of swarming. Do not treat bees within 30 days of queen introduction. Do not treat bees within 30 days of hive relocation. Do not treat bees within 30 days of hive expansion. Do not treat bees within 30 days of hive contraction. Do not treat bees within 30 days of hive destruction. Do not treat bees within 30 days of hive abandonment. Do not treat bees within 30 days of hive relocation. Do not treat bees within 30 days of hive expansion. Do not treat bees within 30 days of hive contraction. Do not treat bees within 30 days of hive destruction. Do not treat bees within 30 days of hive abandonment.

BEET CATTLE, SHEEP, GOATS, HOGS: Use heavy spray 200 gals. water (200 gals. spray per acre). Use 1 lb. per acre in water to cover. Apply only after the animal has been shorn or sheared. Do not apply to the animal's face, eyes, or ears. Do not apply to the animal's mouth or nose. Do not apply to the animal's feet. Do not apply to the animal's tail. Do not apply to the animal's udder. Do not apply to the animal's teats. Do not apply to the animal's vulva. Do not apply to the animal's penis. Do not apply to the animal's scrotum. Do not apply to the animal's testicles. Do not apply to the animal's ovaries. Do not apply to the animal's uterus. Do not apply to the animal's vagina. Do not apply to the animal's cervix. Do not apply to the animal's rectum. Do not apply to the animal's anus. Do not apply to the animal's perineum. Do not apply to the animal's vulva. Do not apply to the animal's penis. Do not apply to the animal's scrotum. Do not apply to the animal's testicles. Do not apply to the animal's ovaries. Do not apply to the animal's uterus. Do not apply to the animal's vagina. Do not apply to the animal's cervix. Do not apply to the animal's rectum. Do not apply to the animal's anus. Do not apply to the animal's perineum.

SEED TREATMENT

For protection of seed and seedling from soil-borne fungi, insects, and nematodes, apply to give uniform coverage of the seed. Follow your State recommendations and may be combined with certain fungicides, to seeds which already have been treated with this product. Excess amount may reduce germination and vigor of seedling. Treated seed should be stored at 50 to 60 degrees Fahrenheit. Do not use on treated seed. **TREATED SEEDS:** Do not use on soil, birds, and other wildlife. **NOTE:** Treat mixed with food or animal feed. Do not use on seed which has been previously treated with this product. Use an appropriate rate according to the following table: **CORN:** Use 4 to 6 oz. / 100 lbs. of seed. **SOYBEANS:** Use 4 to 6 oz. / 100 lbs. of seed. **PEAS:** Use 4 to 6 oz. / 100 lbs. of seed. **SUGARBEETS:** Use 1 to 2 oz. / 100 lbs. of seed. **COTTON:** Use 6 to 8 oz. / 100 lbs. of seed. **SMALL GRAINS:** (wheat, oats, barley, rye, etc.) Use 4 to 6 oz. / 100 lbs. of seed. **NOTE:** This will vary by seed.

STORAGE AND

Do not open dump

STORAGE -

container with pest dust or into a sa

PESTICIDE

pesticide these was contact y once.

CONTAINER

Then disp if allowe out of sm

AVISO

producto

Do not

drift exp

vented b

Do not en

dried

Because c

various c

of Agricu

IT CONTENTS

SEP 15 1983

Stew
9/15/83

Mr. Nick Sanchez
Prentiss Drug & Chemical Co., Inc.
C.B. 2000
Floral Park, NY 11001

300/104413
17/3

Dear Mr. Sanchez:

Subject: Amendment - Revise Label According to
FR Notices 83-2 and 83-3
Prentox Lindane 25W
EPA Registration No. 655-580
Your Application Dated August 23, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

At the next label printing, please make the following revisions to your label.

- a. A Statement of Practical Treatment must be added to the labeling. We suggest the following statement; however, if your toxicological data suggest otherwise, please submit a draft Statement of Practical Treatment for prior approval:

If swallowed - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with a finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin - Remove contaminated clothing and wash affected areas with soap and water.

If in eyes - Flush eyes with plenty of water. Call a physician immediately.

CONCURRENCES

SYMBOL								
SURNAME								
DATE								

- b. In the directions for seed treatment add the following paragraph.

Labels attached to treated seed must read:

TREATED SEED SHOULD NOT BE USED FOR OR MIXED WITH FOOD
OR ANIMAL FEED, OR PROCESSED FOR OIL. EXPOSED TREATED
SEEDS MAY BE HAZARDOUS TO BIRDS AND OTHER WILDLIFE.
DISPOSE OF ALL EXCESS TREATED SEED BY BURIAL AWAY FROM
BODIES OF WATER.

- c. In the precautionary block - environmental hazard section add the following sentences "Do not apply to feed or forage to be fed to or grazed by livestock or poultry. Do not contaminate feed or food crops."

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

RD/IRB:LARocca:DCR-28682:WANG-0301C:pjb:Raven:479-2023:9/12/83.