

# prentox<sup>®</sup>

## PYRONYL POULTRY HOUSE & BARN FLY SPRAY "A"

(For Application by Trained Personnel)

### CONTROLS HOUSE FLIES, STABLE FLIES, HORN FLIES.

#### ACTIVE INGREDIENTS:

Pyrethrins .....	0.62%
Piperonyl Butoxide Technical* .....	5.03%
Petroleum Hydrocarbons .....	94.35%
<b>TOTAL</b>	<b>100.00%</b>

\*Equivalent to 4.02% (Butylcarbityl)(6-Propylpiperonyl) Ether and 1.01% related compounds.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

E.P.A. REG. NO. 655-582

E.P.A. EST. NO. 655-NJ-1

**ACCEPTED**

FEB 27 1980

NET CONTENTS

Under 1 GAL. LOT #  
Fungicide, and Insecticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 655-582

Manufactured by

# prentiss

Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue  
Newark, N.J. 07105

Office: 363 7th Avenue  
New York, N.Y. 10001

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND (DOMESTIC ANIMALS) CAUTION

**HARMFUL IF SWALLOWED.** Avoid inhalation. Remove pets, birds and cover fish aquaria before spraying. In case of contact, wash immediately with soap and water. Wash arms, hands and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Wear natural rubber gloves, water-proof protective clothing and goggles. Keep all unprotected persons out of operational area while applying the insecticide, or working in treated areas during posted period, wear a mask or respirator of a type passed by the U.S. Bureau of Mines. Avoid contamination of feed and foodstuffs.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Remove birds and cover fish aquariums before spraying.

#### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinsate that cannot be used or chemically processed, should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**CONTAINER DISPOSAL: 30 gallons and larger:** a) Reuse container and offer for reconditioning, or b) Triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in an approved landfill, or bury in a safe place.

**Smaller Containers:** As in b) above

**GENERAL:** Consult Federal, State or Local Disposal Authorities for approved alternative procedures.

#### DIRECTIONS FOR USE:

**CONTACT SPRAY — Diluted for Use with Sprayers that give large, wetting spray particles —** To kill Cockroaches and accessible stages of Granary Insects, Rice & Granary Weevil, Red Flour Beetle, Indian Meal Moth Larvae, Caddis, Grain Mites and Saw-Toothed Grain Beetles.

#### DIRECTIONS FOR USE (Cont'd)

Cover or remove exposed food a Concentrate with Oil at the rate of 19 parts Oil, and mix well. Spray out flooring, about base of milking baseboards, areas behind cabinet other insect harbourages using 1 sq. ft. of surface. Force spray into Repeat applications as necessary products should be fumigated or methods using an approved procedure

**SPACE SPRAY — Undiluted in Aer kill Flies, Mosquitoes, Small Flying or remove exposed food and Close room and shut off all air conc equipment. Apply at the rate of 2 Concentrate per 1,000 cu. ft. of spa aerosol generator upward towards and ceiling. Leave room closed for remain in treated areas. Ventilate th is completed. Sweep up and destroy as necessary.**

**SPACE SPRAY — Dilute 1 plus 4 w Barns & Poultry Houses, Warehouse Storage Rooms —** To control House Horn Flies, apply as a fine mist in stanchions or holding pens or over t in pens. Use a fogger or mist ge directed and applied above the anim center of the area where animals ar per 1,000 cu. ft., so that the spra uniformly throughout the entire a ventilate the area for at least 10 m been applied. If used in Dairy Ba thoroughly before milking. Repeat afternoon, or as required. Keep spr nozzle clean, to ensure proper deli fog. Product to be used in mechan give particle size aerosol range or s type.

**SPACE SPRAY — For Use in Equipment —** Kills Flies, Mosquitoes & Gnats. Cover or remove expos handling surfaces. Close room, mak air conditioning or ventilation. Concentrate with Deodorized Kero part Concentrate with 9 part Deodc mix well. Apply at the rate of 50 fl. oz 25,000 cu. ft. of space. Make sure upward towards upper corners of r room closed for one half hour after remain in treated area. Ventilate completed.

BEST DOCUMENT AVAILABLE

30



# prentox<sup>®</sup>

## PYRONYL POULTRY HOUSE & BARN FLY SPRAY "A"

(For Application by Trained Personnel)

### CONTROLS HOUSE FLIES, STABLE FLIES, HORN FLIES.

#### ACTIVE INGREDIENTS:

Pyrethrins .....	0.62%
Piperonyl Butoxide Technical* .....	5.03%
Petroleum Hydrocarbons .....	94.35%
<b>TOTAL</b>	<b>100.00%</b>

\*Equivalent to 4.02% (Butylcarbityl)(6-Propylpiperonyl) Ether and 1.01% related compounds

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

E.P.A. REG. NO. 655-582

E.P.A. EST. NO. 655-NJ-1

**ACCEPTED**

FEB 27 1980

NET CONTENTS

Under U.S. GAL. LOT #  
Fungicide, and Insecticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 655-582

Manufactured by

**prentiss**

Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue  
Newark, N.J. 07105

Office: 363 7th Avenue  
New York, N.Y. 10001

#### DIRECTIONS FOR USE (Cont'd)

Cover or remove exposed food and close room. Dilute Concentrate with Oil at the rate of 1 part Concentrate plus 9 parts Oil and mix well. Spray outside walls of grain bins, flooring about base of milling machinery, walls and baseboards, areas behind cabinets and under pipes and other insect harbourages using 1 gal. of Spray per 1,000 sq. ft. of surface. Force spray into all cracks and crevices. Repeat applications as necessary. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

**SPACE SPRAY — Undiluted in Aerosol Generators** — To kill Flies, Mosquitoes, Small Flying Moths & Gnats. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilation equipment. Apply at the rate of 2 to 6 oz. of undiluted Concentrate per 1,000 cu. ft. of space. Direct output from aerosol generator upward towards upper corners of room, and ceiling. Leave room closed for one half hour. Do not remain in treated areas. Ventilate the area after treatment is completed. Sweep up and destroy fallen insects. Repeat as necessary.

**SPACE SPRAY — Dilute 1 plus 4 with Base Oil for Use in Barns & Poultry Houses, Warehouses, Storage Bins & Storage Rooms** — To control House Flies, Stable Flies & Horn Flies, apply as a fine mist over the livestock in stanchions or holding pens or over the poultry in cages or pens. Use a fogger or mist generator, so spray is directed and applied above the animals and direct spray to center of the area where animals are housed. Apply 1 oz. per 1,000 cu. ft. so that the spray mist is distributed uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in Dairy Barns, aerate the barn thoroughly before milking. Repeat daily, morning and afternoon, or as required. Keep sprayer unit reservoir and nozzle clean, to ensure proper delivery of spray mist or fog. Product to be used in mechanical sprayers, which give particle size aerosol range or smaller, such as ULV type.

**SPACE SPRAY — For Use in Mechanical Fogging Equipment** — Kills Flies, Mosquitoes, Small Flying Moths & Gnats. Cover or remove exposed food and food handling surfaces. Close room, making sure to shut off all air conditioning or ventilation equipment. Dilute Concentrate with Deodorized Kerosene at the rate of 1 part Concentrate with 9 part Deodorized Kerosene and mix well. Apply at the rate of 50 fl. oz. of diluted spray per 25,000 cu. ft. of space. Make sure to direct the spray upward towards upper corners of room & ceiling. Keep room closed for one half hour after spraying. Do not remain in treated area. Ventilate after treatment is completed.

12/78

BEST DOCUMENT AVAILABLE

N 30-A

**CAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
(DOMESTIC ANIMALS)**  
**CAUTION**  
ALLOWED. Avoid inhalation. Remove cover fish aquaria before spraying. In case of eye contact, immediately with soap and water. Wash face thoroughly with soap and water. Wash all contaminated clothing before reuse. Wear natural rubber protective clothing and goggles. Exclude persons out of operational area from insecticide, or working in treated areas. In case of fire, wear a mask or respirator of a type approved by the Bureau of Mines. Avoid contamination of water.

**ENVIRONMENTAL HAZARDS**  
Toxic to fish, birds and other wildlife. Keep away from streams or ponds. Do not contaminate water, soil, or disposal of wastes. Remove from aquariums before spraying.

**DIRECTIONS FOR USE**  
CONFORMANCE WITH FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

**STORAGE AND DISPOSAL**  
Store in original container. Do not use for domestic purposes. Do not dump in water, food or feed by storage or disposal. Dumping is prohibited.  
**DISPOSAL:** Pesticide, spray mixture or residue should not be used or chemically processed, but should be disposed of in a landfill approved for pesticides. Keep away from water supplies.  
**DISPOSAL: 30 gallons and larger:** a) Reseal container for reconditioning, or b) Triple rinse container and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place.  
**Labels:** As in b) above.  
Consult Federal, State or Local Disposal Authority for approved alternative procedures.

**FOR USE:**  
**SPRAY — Diluted for Use with Sprayers that Produce Fine Spray Particles** — To kill Cockroaches, Flies, Mosquitoes, Small Flying Moths & Gnats, Indian Meal Moth Larvae, Mites and Saw-Toothed Grain Beetles.

