

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged, stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings in a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

DIRECTIONS FOR USE (Cont'd.)

To control **CARPET BEETLES:** Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as necessary.

To control **BEDBUGS:** Spray mattresses slightly, particularly around tuft and seams. Take bed apart and spray all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control **FLEAS AND BROWN DOG TICKS:** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

AS A SPACE SPRAY: Dilute at a rate of 1 part concentrate to 10 parts water or oil. Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of **FLIES, FRUIT FLIES, GNATS, MOSQUITOES, SKIPPER FLIES, WASPS, HORNETS, ANTS, MOSQUITOES, SKIPPERS, MEDITERRANEAN FLOUR MOTHS, ANGIOPODID GRAMMID MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, SMALL FLYING MOTHS, AND TOBACCO MOTHS:** Spray direct at an upward angle distributing it uniformly throughout the entire area at a rate of 1 ounce per 1,000 cubic feet of space. Keep area closed for at least 10 minutes. Ventilate and sweep up and destroy dead and fallen insects before reoccupying.

U.L.V. APPLICATION: For rapid kill of exposed or accessible stages of the insects named on this label, dilute 1 part concentrate to 1 to 9 parts water or oil. Apply using conventional mechanical or compressed air equipment (non-thermal) following directions for space spraying. Apply the above dilutions at a rate of 1 to 10 ounces per 1,000 cubic feet of space. When used in meat and poultry processing plants, a dilution of no less than 1 to 1 (50%) is acceptable, and it must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns).

FOR USE ON LIVESTOCK: To control **HORN FLIES, HOUSE FLIES, MOSQUITOES, AND GNATS:** Dilute 1 part concentrate and 39 parts water and apply to wet the hair thoroughly with particular attention to the topline, underline, flanks, withers and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.

To control **STABLE FLIES, HORSE FLIES AND DEER FLIES:** Dilute 1 part concentrate and 9 parts water and apply one quart per adult animal to wet hair thoroughly with particular attention to the ears, flanks, barrel, topline and other body areas commonly attacked by these flies. Repeat treatment each week or as needed.

To control **FACE FLIES:** Dilute one part concentrate and 9 parts water. Use a spray which produces large wetting droplets. Apply to the face of the animal in the morning before releasing to pasture. Apply sufficient to wet the face but not more than 1 1/2 ounces per animal.

For effective control of **MOTILE STAGES OF (BITING) YELLOW LICE AND (BUCKING) BLUE LICE** on cattle, horses, sheep, goats and hogs, dilute this concentrate at a rate of 1 part in 100 parts of water. Spray to thoroughly wet hair of the animal including the head and brush of the tail. Repeat treatment in 10 days to kill newly hatched lice.

To control **POULTRY LICE:** Using a dilution of 1 to 100 in water, dip birds by submerging in a barrel, tank or tub of the solution holding the wings away from the body to insure thorough wetting of all infested areas. In order to avoid chilling the fowls, dip them on a warm day or in a warm building.

For control of **BEDBUGS AND MITES** in poultry houses: Dilute 1 part of concentrate and 9 parts of water and spray crevices of roost poles, cracks in walls, cracks in nests where the bedbug and mites hide.

To control **SHEEP "TICK" OR RED:** Dilute at the rate of 1 part of concentrate to 100 parts of water and thoroughly wet all portions of the body by dipping or by spraying with sufficient pressure and with a nozzle adjustment to give penetration of the wool. Treat at a rate sufficient to wet the animal.

To kill **FLEAS ON LIVESTOCK AND PETS**, and to obtain protection against reinestation: Dilute at a rate of 1 part concentrate and 9 parts water and wet the animal by dipping or spraying. For best results, repeat

prentox®

EMULSIFIABLE SPRAY CONCENTRATE #96

Authorized by U.S.D.A. for use in federally inspected meat and poultry plants

ACTIVE INGREDIENTS:

Pyrethrin	8.00%
Piperonyl Butoxide "Technical"	9.00%
Petroleum Distillate	81.00%
INERT INGREDIENTS	2.00%
TOTAL	100.00%

*Equivalent to 7.00% (Butylcarbitol) (6-propylpiperonyl) ether and 1.00% related compounds.

E.P.A. REG. NO. 655-987

E.P.A. EST. NO. 655-GA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

How to use: — Call a physician or Poison Control Center immediately. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia. **If inhaled** — Remove victim to fresh air. **If on skin** — Remove contaminated clothing and wash affected areas with soap and water. **If in eyes** — Flush eyes with plenty of water. Call a physician immediately, if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist. Avoid skin and eye contact. Wash thoroughly after handling and before eating and smoking. Do not allow children or pets to contact treated surfaces until spray has dried. Do not use on plants. Do not allow spray to contact food, feedstuffs or water supplies. Thoroughly wash dishes and food handling utensils contaminated with this product. Remove pets and cover fish bowls before spraying.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS: Do not use, or store near heat or open flame; exposure to temperatures above 130 F. may cause bursting.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

USE PRECAUTIONS: The use of this product as a space spray in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Remove any birds, pets, or fish aquaria before spraying.

When diluting with water, prepare only the amount needed for immediate use. Holding a water dilution more than 24 hours is not recommended. Agitate occasionally. For oil dilutions, a deodorized petroleum oil is recommended.

Facile premises after treatment and ventilate before reoccupying. Sweep up and remove dead insects. Repeat treatment as necessary. For maximum effectiveness, a combination of localized surface application and space treatment is recommended.

Do not apply directly to food. Do not apply space spray while food processing is underway. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions indicated.

AS A SURFACE SPRAY: For initial clean up of severe insect infestations, dilute at a rate of 1 part concentrate and 4 parts water or oil. For use in edible product areas of federally inspected meat and poultry plants and for normal infestations, dilute 1 part concentrate and 9 parts water or oil. To control **ROACHES, ANTS, SILVERFISH, SPIDERS, CRICKETS, CLOVERMITES, CHEESEMITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUSTY FLOUR BEETLES, SAW-TOOTHED GRILLA BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAMMID MOTHS, SCORFICHS, BOX ELDER BUGS AND CADELLES:** Use a good sprayer adjusted to deliver coarse wet spray. Direct the spray in cracks and crevices, under pallets, around containers of stored foods, around the bases of machinery, behind shelves and drawers. For **SILVERFISH** spray; bookcase. For **ANTS** spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

Manufactured by



prentiss
Drug & Chemical Co., Inc.

Plant: Kaolin Road
Sandersville, GA. 31082
Office: C.B. 2000
Floral Park, N.Y. 11001

flies and ticks on dog should also be treated. To control **SPINOSER** and 24 parts of water pouring 1 to 2 ounces ears of dogs.

In barns, milking parlors, **MOSQUITOES AND** and nine parts of water spray toward the ceiling misting about 20 minutes. Repeat application as needed. **MOSQUITO CONTROL:** using green parts water or oil. Apply particular attention to rest also low or marsh. Repeat as necessary. With mobile or truck, result if a 400 foot as dispersing spray at a 1 Application using five in 4 gallons of water p. To control **MOSQUITO** concentrate to 150 per 25 gallons per acre. Ag lands, swamps, marsh not exceed this dosage **ON GROWING CROPS** (greenhouses) up to or as a clean-up or pre-t used due to residue re use cautions and direct with other insecticide, "exciter" to flush its residues.

On the following crops acre in sufficient va gallons) Coverage of control:

ASPARAGUS: ASPAR BEANS: APHIDS, LE BEETLE, 12-SPOTTED BROCCOLI, CABBAGE APHIDS, CABBAGE DIAMONDBACK MOT, IMPORTED CABBAGE CELERY: GREEN PEA CRABAPPLES: FIRE EGGPLANTS: BLIST FLEA BEETLE, GREE LETTUCE: CABBAG MONDBACK MOT, L

MUSTARD GREENS, CABBAGEWORM, DI BAGE LOOPER, PEPPERS: GREEN PE POTATOES: APHIDS, LEAF HOPPER, RADISH: APHIDS, FLI SPINACH: APHIDS, C TOMATOES: GREEN, STINK BUG, FLEA BE ORNAMENTALS: (AF GONIAS, CAMELLI, DAHLIAS, DOGWOC ROSES, RUBBER PLA BEETLES, LEAFHOP

DROSOPHILA OR FR concentrate with 1 200 with 4 gallons of water. Make treatments as fo

1. Apply liberally to plants. Use sprayers a parts of the diluted spr. for a maximum covera spray between and ber flies, but the emulsio washed from the fruit.

2. Spray the raw stoc 3. Dip baskets in the adhering larvae and p 4. After washing and c just prior to bringing should be sprayed at a parts with 7 gallons 3p the rate of 1 gallon to 7 floors, paying special crevices.

NET CONTENTS

LOT #

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

300/115232
873

REC
FEB

Mr. M. Sanchez
Prentiss Drug & Chemical Co., Inc.
C.D. 200
21 Vernon Street
Floral Park, NY 1101

Dear Mr. Sanchez:

Subject: Prentox Emulsifiable Spray Concentrate 996
EPA Registration No. 655-587
Your Letter Dated January 6, 1984

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure
DCR-05649:A. Hayward:RD-20:1/27/84:del.2/20/84:LM

CONCURRENCES							
SYMBOL	TS-767						
SURNAME	Allegretti						
DATE	2.1.84						

EPA Form 1320-1 (4-81)

OFFICIAL FILE COPY