



prentox

0.5%

CHLORPYRIFOS

(Contains Dursban®)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS
(AND DOMESTIC ANIMALS)

CAUTION

HARMFUL IF SWALLOWED IF SWALLOWED DO NOT INDUCE VOMITING CALL A PHYSICIAN IMMEDIATELY AVOID BREATHING VAPOR OR SPRAY MIST AVOID CONTACT WITH SKIN EYES OR CLOTHING IN CASE OF EYE CONTACT FLUSH WITH PLENTY OF WATER GET MEDICAL ATTENTION IF IRRITATION PERSISTS KEEP AWAY FROM FOOD FEEDSTUFFS AND DOMESTIC WATER SUPPLIES WASH THOROUGHLY AFTER HANDLING DO NOT SMOKE WHILE USING REMOVE ALL PETS AND AQUARIUMS FROM AREA BEFORE SPRAYING

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

NOTE TO PHYSICIAN: Contains petroleum distillate. If gastric lavage is deemed necessary, undertake only after proper placement of a cuffed endotracheal tube. Active ingredients contain a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

- Prohibitions:** DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE, DISPOSAL OR CLEANING OF EQUIPMENT. OPEN DUMPING IS PROHIBITED.
- Pesticide Disposal:** Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.
- Container Disposal:** Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill or by other approved State and local procedures.
- Consumer Packing — 1 gallon or less:** Do not reuse empty container. Wrap container and put in trash collection.

For control of cockroaches, ants, silverfish, spiders, crickets, earwigs and brown dog ticks. Use in and around homes, apartments, garages, similar residential structures, institutions, restaurants and food processing plants. Use outdoors as an aid in reducing annoyance from flies and mosquitoes entering homes.

ACTIVE INGREDIENTS:

Chlorpyrifos [O,O-Diethyl O-(3,5,6-Trichloro-2-Pyridyl) Phosphorothioate]	0.500%
Xylene Range Aromatic Solvent	0.283%
Petroleum Distillates	99.180%
INERT INGREDIENTS:	0.037%
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Do not induce vomiting. Call physician immediately.

IF INHALED - Take person into fresh air immediately.

IF ON SKIN - Wash with plenty of soap and water.

IF IN EYES - Flush with plenty of water. Get medical attention if irritation persists.

SEE ADDITIONAL SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRENTOX® Registered Trademark of Prentiss Drug & Chemical Co. Inc.

DURSBAN® Registered Trademark of The Dow Chemical Co.

E.P.A. Reg. No. 655-581

E.P.A. EST. NO. 655-NJ-1

NET CONTENTS _____

LOT # _____

Manufactured by



prentiss

Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
Newark, N.J. 07105

Office: 363 1 Avenue
New York, N.Y. 10001

DIRECTIONS FOR FOOD HANDLING ESTABLISHMENTS
(Places other than private residential establishments where food is held, processed, prepared or served.)

FOOD AREAS - APPLICATION LIMITED TO CRACK & CREVICE TREATMENT ONLY - Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) and serving areas (when food is exposed and facility is in operation).

To control roaches, ants, earwigs, waterbugs, silverfish, spiders and crickets, apply in small amounts directly into cracks and crevices using equipment capable of delivering a pin stream of insecticide in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases, in conduits, motor housings, junction or switch boxes. Prior to application, de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switches in ON position. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contaminating of food or food processing surfaces.

APPLICATION OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A CRACK AND CREVICE TREATMENT IS NOT PERMITTED.

SERVING AREAS - Facilities where foods are served, such as dining rooms, but excluding areas where foods may be prepared or held, apply as a spot treatment to selective surfaces, such as baseboards, under elements of construction and into cracks and crevices. Avoid treating surfaces likely to be contacted by food. Do not apply when facility is in operation or foods are exposed.

DIRECTIONS FOR HOUSEHOLD AND OUTDOOR USE IN NON-FOOD AREAS - Including garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mwp, closets and storage (after canning or bottling). Apply to baseboard areas around water pipes, surfaces behind and beneath sinks, lockers, tables, pallets and similar areas where insects hide or through which they may enter. Repeat treatment as required.

BROWN DOG TICKS - Apply thoroughly as a spot treatment to infested areas, such as pet beds and resting quarters, nearby cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor covering, where these pests may be present. Fresh bedding should be placed in animal quarters following treatment. Do not spray pets with this product.

FLIES AND MOSQUITOES - Use only on outdoor surfaces. Spray outside surfaces of windows and door frames and other areas where these pests may enter the home. Also spray dark corners and localized resting areas, such as under eaves, porches, patios, and in garages where these insects may congregate. Flies and mosquitoes coming to rest on treated surfaces will be killed. Repeat treatment when effectiveness diminishes.