

# prentox

BEST DOCUMENT AVAILABLE

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Avoid breathing of dust. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling, harmful if swallowed. May be absorbed through skin. Treat lawns in late afternoon. Keep children and pets off treated areas until the following afternoon.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds and other wildlife. Do not apply where runoff is likely to occur. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping prohibited.

**STORAGE:** If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spills, wear protective equipment as required to prevent contact with the product or its vapors. Sweep on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary land fill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## INSECTICIDE DUST ROTENONE 1%

Controls Worms and Insects.  
 See specific names listed on directions panel on Vegetables

**ACTIVE INGREDIENTS:**  
 Rotenone ..... 1.0%  
 Other Cube Resins ..... 2.0%  
**INERT INGREDIENTS** ..... 97.0%

PRENTOX®—Registered Trademark of Prentiss Drug & Chemical Co., Inc.

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**If swallowed** — Call a physician immediately

**If inhaled** — Remove victim to fresh air

**If on skin** — Wash promptly with soap and water. Rinse thoroughly.

**If in eyes** — Flush eyes for at least 15 minutes with water and call a physician immediately.

**SEE SIDE PANEL  
 FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

E.P.A. REG. NO. 655-600      E.P.A. EST. NO. 655-QA-1

NET WEIGHT \_\_\_\_\_ LBS.

LOT # \_\_\_\_\_

**INSECTICIDE DUST** may be applied at any time when dusting conditions are favorable (on calm days). The DUST may be applied during the daytime to either moist or dry foliage. Apply DUST regularly at 7 to 10 day intervals, beginning soon after plants are above ground.

Dosage may vary depending on cultural practices, amounts of foliage, dust equipment, etc. For best results keep foliage evenly coated with the DUST. This is especially important on new growth. Do not apply 1% Rotenone dust to crops listed below less than one day of Harvest.

#### CUCUMBERS, MELONS, SQUASH, PUMPKINS

To control Striped and Twelve Spotted Cucumber Beetles, apply when insects first appear and continue applications weekly until danger of insect injury is past. Redust after rains and continue with regular applications of insecticide dust.

#### TOMATOES

To control Potato Flea Beetles, Colorado Potato Beetles and Small Tomato Hornworms, begin application 2 or 3 days after plants are set in the field. To control Tomato Worms apply when worms are still small. Repeat as necessary.

#### POTATOES and EGG PLANTS

To control Colorado Beetles and Potato Flea Beetles begin application when insects make appearance, repeat at 5 to 7 day intervals as necessary when danger of insect injury is past. Continue regular applications of insecticide dust.

#### BEANS

Mexican Bean Beetles and Bean Thrips, Bean Leaf Beetles. Start when insect and disease pests first appear. Apply to the underside of the leaves as well as top side. Repeat treatment as often as necessary.


#### CABBAGE

Cabbage Lopper Worms, Southern Cabbage Worms, Cross-Striped Cabbage Worms, Imported Cabbageworm. Apply DUST when worms first appear. Repeat as often as necessary.

#### ASPARAGUS

Asparagus Beetle and Spotted Asparagus Beetle. The spears should be dusted with this product and when possible apply when the insects are predominately in the larval state. Repeat as often as necessary.

ACCEPTED  
 JUL 7 1984  
 655-600



Manufactured by  
**prentiss**  
 Drug & Chemical Co., Inc.

Plant: Kaolin Road  
 Sandersville, GA. 31082  
 Office: C.B. 2000  
 Floral Park, N.Y. 11001