

# prentox

## BERRY BUSH SPRAY CONCENTRATE

MIXES EASILY IN WATER

FOR

- Raspberries
  - Loganberries
  - Dewberries
- Blackberries
  - Boysenberries
  - Strawberries

**KILLS:** Aphids, Thrips, Mites, Leafhoppers, Japanese Beetles, Spittlebugs, Rose Scale, Lygus Bug, Field Crickets, Whiteflies, Sap Beetle and Strawberry Leafroller (see chart on Direction panel)

**ACTIVE INGREDIENTS:**

Mineral Oil ..... 91.50%  
Malathion\* ..... 5.50%

**INERT INGREDIENTS:** ..... 3.00%

\*O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.

### KEEP OUT OF REACH OF CHILDREN CAUTION

**STATEMENT OF PRACTICAL TREATMENT**

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. AVOID BREATHING SPRAY MIST  
**IF SWALLOWED:** GIVE EMETIC. CALL A PHYSICIAN  
**IF INHALED:** REMOVE PERSON TO FRESH AIR AT ONCE  
**IF ON SKIN:** WASH THOROUGHLY WITH SOAP AND HOT WATER  
**IF IN EYES:** WASH WITH PLENTY OF WATER FOR 15 MINUTES  
CALL PHYSICIAN  
**NOTE TO PHYSICIAN:** MALATHION IS A CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

E.P.A. Reg. No. 655-614

E.P.A. Est. No. 655-NJ-1

NET CONTENTS \_\_\_\_\_

LOT # \_\_\_\_\_

Manufactured by

**prentiss**  
Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue  
Newark, N.J. 07105

Office: 363 7th Avenue  
New York, N.Y. 10001

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS:**  
**CAUTION:** Harmful if swallowed. Avoid breathing of spray mist and contact with skin. Wash thoroughly after using. Do not use stronger than recommended.

**ENVIRONMENTAL HAZARDS:**  
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**PHYSICAL OR CHEMICAL HAZARDS:**  
Do not use or store near heat or open flame.

**Directions for Use:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL:**

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide, Spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**CONTAINER DISPOSAL:**

**30 & 55 gallon drums:** (a) Reseal container and offer for reconditioning, or (b) triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place.

**SMALLER CONTAINERS: Containers 1 gallon or less—** Do not reuse empty container. Wrap and dispose in trash collection.

**GENERAL:** Consult federal, state or local disposal authorities for approved alternative procedures.

DIRECTIONS FOR USE: Cont d

**AN EFFECTIVE  
BERRYBUSH INSECTICIDE**

One quart Berry-Bush spray will make 10 gallons of finished spray ready to use.

**May Be Used in Hand or Knapsack Sprayer**

**SPRAYING:** Full-coverage spray applications should be made, paying particular attention to underside of leaves. Always observe the printed CAUTIONS and follow the recommendations of State Agricultural Extension Service of your local vicinity where spraying is to be done. Repeat applications as needed.  
**CAUTION:** Make no spray application later than 7 days before harvest.

**EASY SPRAY CHART**

CROP	PEST	AMT. TO USE PER GALLON OF WATER
Blackberries	Mites	6 tablespoons (Approximately 2 fluid ozs.)
Boysenberries	Thrips	6 tablespoons (Approximately 2 fluid ozs.)
	Leafhoppers	6 tablespoons (Approximately 2 fluid ozs.)
Dewberries	Japanese Beetle	6 tablespoons (Approximately 2 fluid ozs.)
Loganberries	Aphids, Rose Scale	12 tablespoons (Approximately 5 fluid ozs.)
Raspberries		6 tablespoons (Approximately 2 fluid ozs.)
Raspberries	Sap Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Do not apply when foliage is wet.	
Strawberries	Aphids, Spider Mites	6 tablespoons (Approximately 2 fluid ozs.)
	Strawberry Root Weevil	6 tablespoons (Approximately 2 fluid ozs.) Apply to soil surface before planting and work into top 6 to 8 inches.
	Lygus Bug	6 tablespoons (Approximately 2 fluid ozs.)
	Spittlebugs	6 tablespoons (Approximately 2 fluid ozs.)
	Field Crickets	6 tablespoons (Approximately 2 fluid ozs.)
	Thrips	6 tablespoons (Approximately 2 fluid ozs.)
	Potato Leafhopper	6 tablespoons (Approximately 2 fluid ozs.)
	Strawberry Leafroller	6 tablespoons (Approximately 2 fluid ozs.)
	Whiteflies	6 tablespoons (Approximately 2 fluid ozs.)

