

ACCEPTED
1981
655-614

BERRY BUSH SPRAY CONCENTRATE

MIXES EASILY IN WATER
FOR

- Raspberries
- Loganberries
- Dewberries
- Blackberries
- Boysenberries
- Strawberries

KILLS: Aphids, Thrips, Mites, Leafhoppers, Japanese Beetle, Spittlebugs, Rose Scale, Lygus Bug, Field Crickets, Whiteflies, Sap Beetle, and Strawberry Leafminer (See chart on Data Panel)

ACTIVE INGREDIENTS: Malathion* 5.50%
INERT INGREDIENTS: 94.50%
 *O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.

DIRECTIONS FOR USE (Cont'd)

AN EFFECTIVE BERRYBUSH INSECTICIDE

One quart Berry-Bush spray will make 10 gallons of finished spray ready to use

May Be Used in Hand or Knapsack Sprayer

SPRAYING: Full-coverage spray applications should be made, paying particular attention to underside of leaves. Always observe the printed **CAUTIONS** and follow the recommendations of State Agricultural Extension Service of your local vicinity where spraying is to be done. Repeat applications as needed.
CAUTION: Make no spray application later than 7 days before harvest.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS:
CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist and prolonged contact with skin. Wash thoroughly after using. Do not use stronger than recommended.
ENVIRONMENTAL HAZARDS:
 This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label. This product is highly toxic to bees exposed to direct treatment or residues on crop. Protective information may be obtained from your Cooperative Agricultural Extension Service.
PHYSICAL OR CHEMICAL HAZARDS:
 Do not use or store near heat or open flame.

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL:
 Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
PESTICIDE DISPOSAL: Pesticide spray mixture or rinse that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
CONTAINER DISPOSAL:
30 & 55 gallon drums: (a) Reseal container and offer for reconditioning, or (b) triple rinse (or equivalent) and offer for recycling, reconditioning or disposal at approved landfill or bury in a safe place.
SMALLER CONTAINERS: Containers 1 gallon or less - Do not reuse empty containers. Do not burn or incinerate.
GENERAL: Consult Federal, State, or local authorities at the time of any fire, spill, or other emergency.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT
 HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. AVOID BREATHING SPRAY MIST!
IF SWALLOWED: GIVE EMETIC. CALL A PHYSICIAN.
IF INHALED: REMOVE PERSON TO FRESH AIR AT ONCE.
IF ON SKIN: WASH THOROUGHLY WITH SOAP AND HOT WATER.
IF IN EYES: WASH WITH PLENTY OF WATER FOR 15 MINUTES. CALL PHYSICIAN.
NOTE TO PHYSICIAN: MALATHION IS A CHOLINESTERASE INHIBITOR. ATROPINE IS AN ANTIDOTE.
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

EPA Reg. No. 655-614 EPA Est. No. 655-NJ-1

NET CONTENTS

EASY SPRAY CHART

CROP	PEST	AMT. TO USE PER GALLON OF WATER
RASPBERRIES	Aphids	6 tablespoons (Approximately 2 fluid ozs.)
	Thrips	6 tablespoons (Approximately 2 fluid ozs.)
	Leafhoppers	6 tablespoons (Approximately 2 fluid ozs.)
LOGANBERRIES	Sap Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Whiteflies	6 tablespoons (Approximately 2 fluid ozs.)
DEWBERRIES	Japanese Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Field Crickets	6 tablespoons (Approximately 2 fluid ozs.)
BLACKBERRIES	Japanese Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Field Crickets	6 tablespoons (Approximately 2 fluid ozs.)
BOYSENBERRIES	Japanese Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Field Crickets	6 tablespoons (Approximately 2 fluid ozs.)
STRAWBERRIES	Strawberry Leafminer	6 tablespoons (Approximately 2 fluid ozs.)
	Whiteflies	6 tablespoons (Approximately 2 fluid ozs.)
RASPBERRIES	Japanese Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Field Crickets	6 tablespoons (Approximately 2 fluid ozs.)
LOGANBERRIES	Sap Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Whiteflies	6 tablespoons (Approximately 2 fluid ozs.)
DEWBERRIES	Japanese Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Field Crickets	6 tablespoons (Approximately 2 fluid ozs.)
BLACKBERRIES	Japanese Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Field Crickets	6 tablespoons (Approximately 2 fluid ozs.)
BOYSENBERRIES	Japanese Beetle	6 tablespoons (Approximately 2 fluid ozs.)
	Field Crickets	6 tablespoons (Approximately 2 fluid ozs.)
STRAWBERRIES	Strawberry Leafminer	6 tablespoons (Approximately 2 fluid ozs.)
	Whiteflies	6 tablespoons (Approximately 2 fluid ozs.)



Manufactured by
Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
 Newark, N.J. 07105
 Office: 363 7th Avenue
 New York, N.Y. 10001