

Prentox[®] 3 WAY DUST GARDEN INSECTICIDE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or inhaled. Do not get into eyes, on tender parts of body or clothing. In case of contact wash thoroughly with soap and water. If it gets into eyes wash out eyes at once with plenty of water. Avoid breathing dust.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning equipment, or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Keep this package away from excessive heat. Avoid storage near steam pipes and radiators.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping prohibited.

STORAGE: If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spills wear protective equipment as required to prevent contact with the product or its vapors. Sweep onto a shovel and put the sweepings into a salvage drum. Dispose of wastes as below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ACTIVE INGREDIENTS:

Sulfur	10.00%
Rotenone	1.00%
Other Cube Resins	2.00%
INERT INGREDIENTS:	87.00%
TOTAL:	100.00%

PRENTOX[®] — Registered Trademark of Prentiss Drug & Chemical Co., Inc.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If on skin — Wash thoroughly with soap & water.

If in eyes — Wash out with plenty of water.

SEE SIDE PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS.

E.P.A. REG. NO. 655-617

E.P.A. EST. NO.

NET CONTENTS

LOT #

DIRECTIONS FOR USE (Cont'd.)

GENERAL INSTRUCTIONS

May be applied at anytime when dusting conditions are favorable (on a calm day). The dust may be applied during the daytime to either moist or dry foliage. Apply dust regularly at 7 to 10 day intervals, beginning soon after plants are above ground. For best results keep foliage evenly coated with the dust. This is especially important on new growth. Thoroughly dust upper and lower leaf surfaces.

CUCUMBERS, MELONS, SQUASH, PUMPKINS

To control striped and twelve spotted cucumber beetles: Apply when insects appear and continue applications weekly until danger of insect injury is past. Re-dust after rains and continue with regular application of 3-Way Dust as often as necessary.

TOMATOES

To control potato flea beetles and Colorado potato beetles and small tomato Hornworms: Begin application 2 or 3 days after plants are set in the field. To control tomato worms apply when worms are still small. Continue dusting with 3-Way Dust as often as necessary.

POTATOES and EGG PLANTS

To control Colorado beetles and potato flea beetles: Begin application when insects make appearance. Repeat at 5 to 7 day intervals as necessary when danger of insect injury is past. Continue regular applications of 3-Way Dust as often as necessary.

CABBAGE

Important Cabbage Worms. Apply when worms appear. Repeat dusting as often as necessary.

ASPARAGUS

Asparagus Beetle and Spotted Asparagus Beetle: The spears should be dusted with this product and when possible apply when the insects are predominately in the larval state. Repeat dusting as often as necessary.

ROSE BUSHES

Slugs on Rose Bushes. Dust bushes thoroughly while wet with dew. Repeat dusting in a few days if necessary.

BEST DOCUMENT AVAILABLE



Manufactured by

prentiss
Drug & Chemical Co., Inc.

Plant: Kaolin Road
Sandersville, GA. 31082

Office: C.B. 2000
Floral Park, N.Y. 11001

