

DO NOT APPLY DIRECTLY TO WATER

# prentox<sup>®</sup>

## POTATO

### TOMATO POTATO DUST or SPRAY

200-305

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION**— Harmful if swallowed inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Avoid breathing dust or spray mist. Wash thoroughly after dusting or spraying. Avoid contamination of feed and foodstuff.

#### ENVIRONMENTAL HAZARDS.

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. (X)

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its label.

#### STORAGE AND DISPOSAL

**PROHIBITIONS** — Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL** — Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**CONTAINER DISPOSAL** — Dispose of in an incinerator or landfill approved for pesticide containers or buried in a safe place.

*See attached sheet*

#### ACTIVE INGREDIENTS:

Methoxychlor-technical<sup>\*</sup> 10.00%

<sup>\*</sup>Equivalent to 8.8% 2,2 bis (p-methoxyphenyl)

-1,1,1-trichloroethane and 1.2% of related compounds.

Captan<sup>\*</sup> 6.00%

<sup>\*</sup>N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

**INERT INGREDIENTS** 84.00%

#### KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**If Swallowed** — Get medical attention immediately.

**If Inhaled** — Remove victim to fresh air. Apply respiration if indicated.

**If on Skin** — Wash area immediately with soap and warm water.

**If in Eyes** — Flush with plenty of water and get prompt medical attention.

#### THIS PRODUCT CONTAINS

prentox contains both fungicide and insecticide ingredients. Controls Early and Late Blight, Colorado Potato Beetles, Striped Cucumber Beetle, Leafhoppers, Flea Beetles, Fruit Worms, Hornworm and Spotted Cucumber Beetles.

#### DUSTING

For tomatoes, prentox may be applied to either moist or dry foliage at any time when dusting conditions are favorable. Apply at the rate of 35 pounds per acre at 7 to 10 day intervals, beginning soon after plants are above ground. Do not apply to tomatoes less than 7 days before harvest.

#### SPRAYING

For potatoes mix 25 pounds to 100 gals. of clean water. Apply 100 gals. of the diluted spray per acre. For making smaller quantities mix 2 1/2 pounds to 10 gals. of clean water. Be sure mixing tank and sprayer are thoroughly clean and free from residue before using. Apply at 7 to 10 day intervals, beginning soon after plants are above the ground.

**CAUTION** — Harmful if swallowed. Avoid breathing spray. Avoid contact with skin, eyes, and clothing. Wash thoroughly if it gets into the eyes. Flush with plenty of water at once. Avoid contamination of food and foodstuffs.

This product is toxic to fish. This product is toxic to bees and should not be applied when bees are actively visiting the area. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water. Do not contaminate water by cleaning of equipment or disposal of wastes. (X)

**DO NOT FEED CULLED POTATOES TO DAIRY ANIMALS**

Do not feed culled potatoes to dairy animals.

E.P.A. Reg. No. 655-625

E.P.A. EST. NO. 655-NJ-1

NET CONTENTS \_\_\_\_\_

LOT # \_\_\_\_\_

BEST DOCUMENT AVAILABLE

Manufactured by

# prentiss

Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue  
Newark, N.J. 07106

Office: 363 7th Avenue  
New York, N.Y. 10001



*See attached sheet*

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label.

\* Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.  
Open dumping prohibited.

STORAGE: If container is damaged: STOP any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear proper protective gear. On cleanup, wear protective equipment as required to prevent contact with the product. Sweep on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 19 1983

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No. 655-625



SEP 19 1

Mr. Nick Sanchez  
Frontier Drug & Chemical Co., Inc.  
363 Seventh Ave.  
New York, NY 10001

Dear Mr. Sanchez:

Subject: Frontier Fumate Tomato ~~Label~~ Dust or Spray  
EPA Registration No. ~~011-51~~  
Your Application Dated August 22, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger  
Product Manager (12)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767)

Enclosure

TS-767:Ellenberger;DCR-28415;WANG-0342C;KIM;Raven;479-2013;9/14/83

CONCURRENCES							
SYMBOL ▶							
SIGNATURE ▶							
DATE ▶							