

prentox

FOGGING INSECTICIDE PF

PRECAUTIONARY STATEMENTS

Hazards to Humans &
Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water supplies by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARD

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL — Pesticide spray mixture or rinse that cannot be used should be disposed of according to procedures approved by Federal, State or Local disposal authorities.

CONTAINER DISPOSAL — Under 30 gallon sizes: Triple rinse (or equivalent) and dispose of in an approved landfill or bury in a safe place. 30 gallons sizes and over: a) reseal containers and offer for reconditioning or b) Triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place.

GENERAL — Consult Federal, State or local disposal authorities for approved alternative procedures.

In mushroom production and processing: To control mushroom flies or fungus gnats in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 5 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and grading plants, use at a rate of 1 ounce to 2000 cubic feet of space.

Designed for Use in Edible Product Areas
of Official Establishments Operating
Under the Meat, Poultry, Shell Egg
Grading and Egg Products Inspection
Programs and in Other Food Processing
Plants.

Apply with Equipment Designed and Adjusted to
Deliver a True Aerosol Mist

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
*Piperonyl Butoxide, Technical	5.0%
Petroleum Distillate	94.5%
TOTAL	100.0%

*Equivalent to 4.0% butylcarbitol isopropylpiperonyl ether and 1.0% related compounds.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

EPA REG NO 655-643

EPA EST NO 655-NJ-1

NET CONTENTS

LOT #

Manufactured by



prentox

Drug & Chemicals

Plant: 538 Wilson Avenue
Newark, N.J. 07105

Office: 363 7th Avenue
New York, N.Y. 10001

DIRECTIONS FOR USE (Cont'd.)

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Stored food products which are infested should be fumigated or treated by other effective methods using an approved product intended for that purpose.

Space Spray: For control of accessible, exposed stages of flies including fruit flies, mosquitoes, small flying moths, gnats, wasps, hornets, cheese skippers, and these common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, Tobacco moths, meal worms and grain mites. Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (in droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. For best control of wasps and hornets, apply directly to nests evening when the insects have returned to the nests. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

Space and Contact Spray: For control of accessible, exposed stages of silverfish, crickets, roaches, spiders, scorpions, clover mites, cheese mites, earwigs, and ants, first using equipment which can deliver a pin spray, direct spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects are known to seek shelter. Contact as many insects as possible, spray directly at ant trails. Do not allow spray to contact stored food. Then, shutting all doors, windows and shutting off all ventilating systems proceed with space spraying directions as given above, use a dosage rate of 1 ounce per 1000 cubic feet of space. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.