

prentox[®]

FOGGING INSECTICIDE PF

Designed for Use in Edible Product Areas
of Official Establishments Operating
Under the Meat, Poultry, Shell Egg
Grading and Egg Products Inspection
Programs and in Other Food Processing
Plants.

Apply with Equipment Designed and Adjusted to
Deliver a True Aerosol Mist.

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
Piperonyl Butoxide, Technical*	5.0%
Petroleum Distillates	94.5%
TOTAL:	100.0%

*Equivalent to 4.0% (butylcarbitol) (6-propylpiperonyl)
ether and to 1.0% related compounds

PRENTOX[®] — Registered Trademark of Prentiss Drug &
Chemical Co., Inc.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If **swallowed** — Call a physician or Poison Control Center
immediately. **DO NOT INDUCE VOMITING**. Vomiting may
cause aspiration pneumonia.

If **inhaled** — Remove victim to fresh air.

If **on skin** — Remove contaminated clothing and wash affected
areas with soap and water.

If **in eyes** — Flush eyes with plenty of water. Call a physician
immediately, if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY
STATEMENTS

E.P.A. REG. NO. 655-643

E.P.A. EST. NO. 655-QA-1

NET CONTENTS _____

LOT# BEST DOCUMENT AVAILABLE

Manufactured by



prentiss

Drug & Chemical Co., Inc.

Plant: Kaolin Road
Sandy Ille, GA. 31082

Office: C.B. 2nd J
Floral Park, N.Y. 11001

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin.
Avoid breathing vapors or spray mist. Avoid
contact with skin and eyes. In case of contact,
immediately flush eyes or skin with plenty of
water. Obtain medical attention if irritation
persists. If swallowed, do not induce vomiting.
Call a physician immediately.

Do not apply directly to food. In commercial food
handling facilities, cover or remove any food and
food processing equipment during application.
Do not apply while food processing is under way.
After space spraying in meat packing plants,
bakeries and other food processing plants, wash
with potable water all equipment, benches,
shelving, etc., where exposed food will be
handled. In the home, all food processing surfaces
and utensils should be covered during treatment
or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums
before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other
wildlife. Do not apply directly to water. Do not
contaminate water by disposal of wastes. Apply
this product only as specified on this label.

PHYSICAL HAZARDS

Do not use, or store near heat or open flame.
Exposure to temperatures above 130°F. may
cause bursting.

Directions for Use:

It is a violation of Federal law to use this product in
a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage
or disposal.

Open dumping is prohibited.

STORAGE: If container is damaged Stop any
leaks by repositioning the container or by
patching or otherwise repairing the leaks. Take
care to avoid contact with pesticide and wear
protective gear. On cleanup of spilled liquids,
wear protective equipment as required to prevent
contact with the product or its vapors. Cover the
spilled areas with generous amounts of absorbent
material, such as clay, diatomaceous earth, sand
or sawdust. Sweep the contaminated absorbent
on to a shovel and put the sweepings into a salvage
drum. Dispose of wastes as below. Place any
leaking container into a similar drum or glass
container. Do not store, use, pour or spill near heat
or open flame.

STORAGE AND DISPOSAL (cont'd)

PESTICIDE DISPOSAL: Wastes resulting from
the use of this product may be disposed of on site
or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equiva-
lent). Then offer for recycling or reconditioning,
or puncture and dispose of in a sanitary landfill, or
by other procedures approved by state and local
authorities.

Directions for Use:


It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

In mushroom production and processing: To control mushroom
flies or fungus gnats in growing houses, apply at a rate of 1 ounce
per 1000 cubic feet of space. Fogger should be operated 8 to 10
feet away from beds and directed slightly upward over them.
Circulating fans should be left in operation, but for best results,
refrigeration should be turned off. In packing and canning
plants, use at a rate of 1 ounce to 2000 cubic feet of space.

The use of this product in food processing establishments
should be confined to time periods when the plant is not in
operation. Foods should be removed or covered during
treatment. All food processing surfaces should be covered
during treatment or thoroughly cleaned before use. Stored
food products which are infested should be fumigated or treated
by other effective methods using an approved product intended
for that purpose.

Space Spray: For control of accessible, exposed stages of flies,
including fruit flies, mosquitoes, small flying moths, gnats,
wasps, hornets, cheese skippers, and these common stored
product pests: granary weevils, rice weevils, confused flour
beetles, saw-toothed grain beetles, spider beetles, cigarette
beetles, Drugstore beetles, Angoumois grain moths,
Mediterranean flour moths, Indian meal moths, Tobacco moths,
meal worms, and grain mites. Close windows and doors and shut
off ventilating systems. Use a fogger or vaporizer adjusted to
deliver an aerosol spray (no droplets over 50 microns in diameter
and 80% less than 30 microns). Apply at the rate of 1 ounce per
1000 cubic feet of space. Direct the spray toward the ceiling and
upper corners of the area and behind obstructions. For best
control of wasps and hornets, apply directly to nests in the
evening when the insects have returned to the nests. Keep the
area closed for at least 15 minutes. Vacate the treated area and
ventilate before reoccupying. Repeat treatment as necessary.

Space and Contact Spray: For control of accessible, exposed
stages of silverfish, crickets, roaches, spiders, scorpions, clover
mites, cheese mites, earwigs, and ants, first using equipment
which can deliver a pin spray direct spray into all cracks and
crevices in woodwork, walls, floors, underneath sinks, behind
pipes and in all places where these insects are known to seek
shelter. Contact as many insects as possible, spray directly all
ant trails. Do not allow spray to contact stored food. Then,
shutting all doors, windows and shutting off all ventilating
systems proceed with space spraying directions as given above.
Use a dosage rate of 1 ounce per 1000 cubic feet of space. Keep
the area closed for at least 15 minutes. Vacate the treated area
and ventilate before reoccupying. Repeat treatment as
necessary.



10/83

BEST DOCUMENT AVAILABLE

prentox[®]

FOGGING INSECTICIDE PF

Designed for Use in Edible Product Areas
of Official Establishments Operating
Under the Meat, Poultry, Shell Egg
Grading and Egg Products Inspection
Programs and in Other Food Processing
Plants.

Apply with Equipment Designed and Adjusted to
Deliver a True Aerosol Mist

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
Piperonyl Butoxide, Technical*	5.0%
Petroleum Distillates	94.5%
TOTAL:	100.0%

*Equivalent to 4.0% (butylcarbitol - propylpiperonyl) ether and to 1.0% related compounds

PRENTOX[®] Registered Trademark of Prentiss Drug & Chemical Co. Inc.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.

If inhaled - Remove victim to fresh air.

If on skin - Remove contaminated clothing and wash affected areas with soap and water.

If in eye - Flush eyes with plenty of water. Call a physician immediately if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG NO 655-643

EPA EST. NO 655-QA-1

NET CONTENTS _____

LOT # _____

Manufactured by



prentiss
Drug & Chemical Co. Inc.

Plant: Kaolin Road
Sandusky, N.Y. 14882
Office: C.B. 2nd
Floral Park, N.Y. 11001

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. If swallowed, do not induce vomiting. Call a physician immediately. Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash with potable water all equipment, benches, shelving, etc., where exposed food will be handled. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets and birds, and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use, or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Open dumping is prohibited.

STORAGE: If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour or spill near heat or open flame.

STORAGE AND DISPOSAL (cont'd)

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer to recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

In mushroom production and processing: To control mushroom flies or fungus gnats in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2000 cubic feet of space.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Stored food products which are infested should be fumigated or treated by other effective methods using an approved product intended for that purpose.

Space Spray: For control of accessible, exposed stages of flies, including fruit flies, mosquitoes, small flying moths, gnats, wasps, hornets, cheese skippers, and these common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths, mealworms, and grain mites. Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. For best control of wasps and hornets, apply directly to nests in the evening when the insects have returned to the nests. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

Space and Contact Spray: For control of accessible, exposed stages of silverfish, crickets, roaches, spiders, scorpions, clover mites, cheese mites, earwigs, and ants. First using equipment which can deliver a pin spray, direct spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects are known to seek shelter. Contact as many insects as possible, spray directly at ant trails. Do not allow spray to contact stored food. Then, shutting all doors, windows and shutting off all ventilating systems proceed with space spraying, directions as given above. Use a dosage rate of 1 ounce per 1000 cubic feet of space. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

655-643