PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful 1 swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. If swallowed, do not induce vomiting, Call a physician immediately.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying in meat packing plants, bakeries and other food processing plants, facilities and equipment must be thoroughly washed with an acceptable detergent solution and then rinsed with potable water to remove all traces of contamination.

Remove pets and birds, and cover lish aquariums before spraying

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use, or store near heat or open flame. Exposure to temperatures above 130°F, may cause bursting.

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Open dumping .s prohibited.

STORAGE: If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent. contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour or spill near heat or open flame.

prentox BEST AVAILABLE CBPY

FOGGING **INSECTICIDE PF**

Designed for Use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants.

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

ACTIVE INGREDIENTS:

Pyrethrina		0.5%
Piperonyl Butoxide, Technical*	· · · · · · · · · · · · · · · · · · ·	5.0%
Petroleum Distillates		<u>. 94.5%</u>
-	TOTAL	100.0%

*Equivalent to 4.0% (butylcarbityl) (6-propylpiperonyl) ether and to 1.0% related compounds.

PRENTOX[®] -- Registered Trademark of Prentiss Drug & Chemical Co., Inc.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed -- Call a physician or Poison Control Center immediately DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia

If inhaled - Remove person to fresh air.

If on skin - Remove contaminated clothing and wash affected areas with soap and water.

If in eyes - Flush eyes with plenty of water. Call a physician immediately, if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. REG. NO. 655-643

E.P.A. EST. NO. 655-QA-1

NET CONTENTS _

the use of this product may be disposed of on site. or at an approved waste disposal facility.

CONTAINER DISPOSAL: Tripletinse (or equivalent). Then offer for recycling or reconditioning. or puncture and dispose of in-a sanktary lanofill, or by other procedures approved by state and local authorities.

PESTICIPE DISPOSAL: Wastes resulting from

STORAGE AND DISPOSAL (cont'd)

Directions for Use:


It is a violation of Federal law to use this product in a manner Inconsistent with its labeling.

In mushroom production and processing: To control mushroom flies or fungue gnate in growing houses, apply at a rate of 1 ounce. per 1000 cubic feet of space. rogger should be operated 8 to 30 feet sway from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2000 cubic feet of space.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Stored food products which are infested should be furnigated or treated. by other affective inethode using an approved product intended. for that purpose.

Space Spray: For control of accessible, exposed stages of flies, including fruit flies, mosquitoes, small living moths, gnats, wasps, hornets, cheese skippers, and these common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, Angoumois grain moths, Mediterannean flour moths, Indian meal moths, Tobaccomoths, meal worms, and grain mites: Close windows and doors and shut off ventilating systems. Use a logger or vaporizor adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. For best control of wasps and hornets, apply directly to nests in the evening when the insects have returned to the nests. Keep the area closed for at least 15 minutes. Vacate the treated area and ventiliate before reoccupying. Repeat treatment as necessary

Space and Contact Spray: For control of accessible, exposed stages of ailverfish, crickets, roaches, spiders, scorpions, clover mites, cheese mites, earwigs, and ants, first using equipment which can deliver a pin spray, direct spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects are known to seek shelter. Contact as many insects as possible, spray directly all ant trails. Do not allow spray to contact stored food. Then, shutting all doors, windows and shutting off all ventilating systems proceed with space spraying directions as given above. Use a dosage rate of 1 ounce per 1000 cubic feet of space. Keep the area closed for at least 15 minutes. Vacate the treated area end ventilate before reoccupying. Repeat treatment as necessary.



2/85



