

prentox[®]

DIAZINON[™]

DDVP RESIDUAL CONCENTRATE

AN INSECTICIDE FOR FORMULATING HOUSEHOLD RESIDUAL INSECTICIDE

ACTIVE INGREDIENTS:

0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	25.00%
2,2-Dichlorovinyl dimethyl phosphite	23.25%
Related Compounds	1.75%
Aromatic Petroleum Derivative Solvent	46.26%
INERT INGREDIENTS:	3.74%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN



**POISON
DANGER**



STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled — Move victim to fresh air. Apply respiration if indicated. Call a physician immediately.

If on skin — Remove contaminated clothing and wash immediately with soap and water. Call a physician immediately if symptoms of poisoning occur.

If in eyes — Flush eyes for at least 15 minutes with water. Call a physician immediately.

DIAZINON[™] is the registered trademark of Ciba-Geigy Corp.
E.P.A. Reg. No. 655-881 E.P.A. EST. NO. 655-MJ-1

NET CONTENTS _____


LOT# _____

Manufactured by



prentiss
Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
Newark, N.J. 07105
Office: 363 7th Avenue
New York, N.Y. 10001



Note to Physician

Symptoms: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

Treatment: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine should be injected at ten-minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. 2-PAM is also antidotal and may be administered in conjunction with atropine. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by blood test.

STORAGE AND DISPOSAL

PROHIBITIONS

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. OPEN DUMPING IS PROHIBITED.

PESTICIDE DISPOSAL

PESTICIDE, SPRAY MIXTURE, OR RINSATE THAT CANNOT BE USED OR CHEMICALLY PROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

(a) RESEAL CONTAINER AND OFFER FOR RECONDITIONING, OR (b) TRIPLE RINSE (OR EQUIVALENT) AND OFFER FOR RECYCLING, RECONDITIONING, OR DISPOSAL IN APPROVED LANDFILL, OR BURY IN A SAFE PLACE. IF LESS THAN 30 GALLONS: TRIPLE RINSE (OR EQUIVALENT) AND OFFER FOR RECYCLING, RECONDITIONING, OR DISPOSAL IN APPROVED LANDFILL OR BURY IN A SAFE PLACE.

GENERAL

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES.

ACCEPTED

MAY 6 1981

Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended, for the pesticide registered under EPA Reg. No. 655-651

1/81

23