


PRENTOX(R) DIAZINON DDVP RESIDUAL CONCENTRATE

An insecticide concentrate for formulating use only

ACTIVE INGREDIENTS:

0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate.....	25.00%
2,2 Dichlorovinyl dimethyl phosphate	23.25%
Related Compounds	1.75%
Xylene Range Aromatic Solvent.....	46.26%
INERT INGREDIENTS:.....	3.74%
TOTAL	100.00%

PRENTOX(R) - Registered Trademark of PRENTISS DRUG & CHEMICAL CO., INC.




STATEMENT OF PRACTICAL TREATMENT
If swallowed - Call a physician immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
If inhaled - Move victim to fresh air. Apply respiration if indicated. Call a physician immediately.
If on skin - Remove contaminated clothing and wash immediately with soap and water. Call a physician immediately if symptoms of poisoning occur.
If in eyes - Flush eyes for at least 15 minutes with water. Call a physician immediately.

DIAZINON® is the registered trademark of Ciba Geigy Corp.
E.P.A. Reg. No. 655-651 E.P.A. EST. NO. 656GA-1

NET CONTENTS _____

LOT# _____



Manufactured by
prentiss
Drug & Chemical Co., Inc.

Plant: Keolin Road
Sandersville, GA. 31082
Office: C.B. 2000
Floral Park, N.Y. 11001

LEFT PANEL

665. 657

NOTE TO PHYSICIAN

SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

Treatment: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine should be injected at ten-minute intervals in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. 2-PAM is also antidotal and may be administered in conjunction with atropine. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by blood test.

STORAGE

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE - If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled area with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent or to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance."

CONTAINER DISPOSAL Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

BEST DOCUMENT AVAILABLE

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....