

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS & DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Avoid breathing vapor. Avoid contact with skin, eyes or clothing. Wear mask or respirator of type recommended by NIOSH (National Institute for Occupational Safety and Health) to give adequate protection against this material. Wear protective clothing. Wash exposed skin with soap and water after handling and before eating. Wash all contaminated clothing with soap and water separately from other clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Use for use, or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Directions for Use (Continued)

INDOOR USE AS A SPACE SPRAY UNDOUBTED: The use of this product in food processing or food handling establishments should be confined to time periods when the plants or facilities are not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 10 microns). Apply at a rate of 1 ounce per 1,000 cubic feet of space. Direct the spray toward ceiling and upper corners of the area behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. Sweep up and destroy fallen insects.

The method of application will control exposed, accessible stages of FLIES (including FRUIT FLIES), MOSQUITOES, SMALL FLYING MOTHS, GNATS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS, BOXELDER BUGS, EARWIGS, ANTS and the following common stored product pests: GRAIN ARMYWORMS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANGIOUMS, GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, YELLOW MEALWORMS, DARK MEALWORMS, GRAIN MITES, CADELLS and RED FLOUR BEETLES. DRIED FRUIT PRODUCTS in storage or being processed for control of CONFUSED FLOUR BEETLES, DRIED FRUIT BEETLES, INDIAN MEAL MOTHS AND SAW TOOTHED GRAIN BEETLES. Remove and destroy infested products. As noted above, disperse this product as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use one-half ounce per 1,000 cubic feet of space. Repeat application at 14 days in any given month. Leave the room closed for one hour. Do not remain in treated areas and ventilate the area before re-entry. Sweep up and destroy fallen insects.

STORED FOOD: (Stored in bulk, wall bags or cloth bags in warehouses, storage bins, flour mills, grain elevators, granaries for control of ANGIOUMS, GRAIN MOTHS, CADELLS, CONFUSED FLOUR BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RICE WEEVILS, SAW TOOTHED GRAIN BEETLES, YELLOW MEALWORMS, CHEESE MITES, CHEESE SKIPPERS, CIGARETTE BEETLES, GRAIN MITES, GRAIN ARMYWORMS, DARK MEALWORMS, RED FLOUR BEETLES. Remove and destroy infested products. As noted above, disperse this product first by directing spray into corners, under pallets, and around stacks of packaged foods lying as many insects as possible and then treat as a space spray.

Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose. Use a dosage of one-half ounce per 1,000 cubic feet of space. Do not remain in treated areas and ventilate area before re-entry. This application will kill accessible, exposed stages of those stored product pests listed above. Sweep up and destroy fallen insects.

CONTACT AND SURFACE SPRAY: For the control of roaches dilute 12 1/2 ounces to make one gallon of finished solution in oil. Use a good sprayer adjusted to

prentox[®]

Pyronyl Cil

Concentrate #12294-A

For Use by Trained Personnel Only

Authorized by U.S.D.A. for use in
 Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide Technical*	2.00%
N-Octyl Bicycloheptene Dicarboximide	2.94%
Petroleum Distillates	94.06%
TOTAL	100.00%

*Consists of 1.60% (butylcarbityl) (6-propylpiperonyl) Ether and 0.40% related compounds

PRENTOX[®] - Registered Trademark of Prentiss Drug & Chemical Co., Inc.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.

If inhaled - Remove victim to fresh air.

If on skin - Remove contaminated clothing and wash affected areas with soap and water.

If in eyes - Flush eyes with plenty of water. Call a physician immediately if irritation persists.

SEE SIDE PANEL FOR
 ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. Reg. No. 655-665

NET CONTENTS

LOT #

Manufactured by

prentiss

Drug & Chemical Co., Inc.

Plant: Kaolin Road
 Sandersville, GA. 31082

Office: C.B. 2000
 Floral Park, N.Y. 11001

deliver a coarse, wet spray. Direct the spray into hiding places, cracks and crevices under pallets, around containers of stored foods, around bases of machinery, behind shelves and drawers. Spray roaches directly whenever possible. Repeat application as necessary.

INDOOR USE AS A SPACE SPRAY: For use in conventional mechanical logging equipment to kill FLIES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

SURFACE SPRAY: To achieve maximum control of crawling insects (ANTS, BOXELDER BUGS, CLOVER MITES, DRUGSTORE BEETLES, EARWIGS, LICE, SPIDER BEETLES) dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well and apply as a coarse wetting spray using approximately 1/2 gallon of spray mixture to 750 square feet of surface. Spray walls, floor and other surfaces of bins and storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Treat unloading, handling and processing areas and inside conveying processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

For in transit protection, clean and spray truck sides, boxcars and ship's hold before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked.

To control INDIAN MEAL MOTHS and ANGIOUMS GRAIN MOTHS in stored grain, apply to the surface of binned grain at a rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of moths and continuing until early fall. In case of severe infestation break up webbing with rake before spraying and make a second application in two weeks. As an aid in control of MOSQUITOES, GNATS, WASPS AND SKIPPERS, treat walls, ceilings, moldings, screens, door and window frames, beams, light cords and similar resting places.

ANIMAL QUARTER USE: (Cattle barns, horse barns, poultry houses, swine houses, etc.) As a space spray diluted for use in conventional mechanical logging equipment to kill FLIES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS. Dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well. Apply at the rate of 2 fluid ounces per 1,000 cubic feet of space above the animals. Direct spray towards the upper portions of the enclosure filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

MUSHROOM PRODUCTION AND PROCESSING: To control MUSHROOM FLIES or FUNGUS GNATS in growing houses, dilute 1 part of concentrate plus 29 parts Methylene Chloride and mix well and apply at a rate of 1 ounce per 1,000 cubic feet of space. Fogger should be operated 10 to 15 feet away from beds and should be directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off in packaging and curing plants. Use at a rate of one-half ounce per 1,000 cubic feet of space.

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS: Dilute one part concentrate with 3 parts base oil (refined kerosene) and use with a ULV spraying device at the rate of one ounce per 1,000 cubic feet of room space. Apply the spray into all cracks and crevices in woodwork, walls, floors, under drains, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Repeat as necessary for control of exposed or accessible stages of ROACHES, SILVERFISH, WATERBUGS, EARWIGS, ANTS, SPIDERS, FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS. ~~Remove exposed food products and packaging materials. Must be covered or removed. After use, equipment and vessels must be thoroughly rinsed with potable water before operation and allowed to dry. Do not spray over open flames or heat sources.~~

OUTDOOR GROUND APPLICATION UNDOUBTED: To control ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS. For best results, apply when there is a light breeze of approximately 1 mph and temperature is cool (50° or less). With a ULV spraying device spray at a dosage rate of 5 ounces per minute up wind of the area to be controlled. If the equipment is truck-mounted, keep the speed to approximately 5 mph with a swath width of no more than 300 feet.

To control MOSQUITO LARVAE, dilute at the rate of 15 ounces per 50 gallons of water and apply at the rate of 20 to 25 gallons per acre. Apply as a coarse wetting spray to the border or on shallow water of wet lands, swamps and marshes where mosquito larvae are known to breed.

TEMPORARY REDUCTION OF ANNOYANCE FROM FLIES, MOSQUITOES AND SMALL FLYING MOTHS OUTDOORS: The above mentioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in public gathering places (drive-in restaurants, drive-in theaters, golf courses, municipalities, parks, playgrounds and recreational areas). Direct application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. Application should be made prior to attendance. Repeat as necessary.

In additional outdoor areas (corals, feed lots, swine lots and pens, horse water, drinking fountains and animal feed before use. Fill area with mist directing application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage in roos, avoid exposure of reptiles to the product. Repeat as necessary.

FOR USE ON ANIMALS: To protect beef and dairy cattle, horse and dogs from FLIES, HOUSE FLIES, MOSQUITOES AND GNATS, dilute 1 part of concentrate plus 29 parts oil (refined kerosene) and mix well. Apply a light mist sufficient to wet the tips of the hair. To control STABLE FLIES, HORSE FLIES and DEER FLIES on beef and dairy cattle and horses, apply 2 ounces per adult animal sufficient to wet the hair but not to soil the hide. Repeat treatment once or twice daily or at intervals to give continued protection.

To control the mite stages of BLUDD LICE on beef and dairy cattle, apply the spray to thoroughly wet the hair. Use 2 ounces or less per animal and repeat 7 to 10 days as required.

BEST AVAILABLE



A 4x4 grid of 16 small, stylized, abstract figures. Each figure is composed of black dots and lines arranged in a unique pattern, representing different combinations of the four basic elements (Earth, Water, Fire, Air) and their interactions. The figures are arranged in a grid that is 4 units wide and 4 units high.