

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. Storage and disposal should be in accordance with local, State, and Federal regulations.

**STORAGE:** If container is damaged, stop any leaks by tightening the container or by patching or otherwise repairing the leak. Take care to avoid contact with pesticide and wear protective gear. Do not use if spilled liquid wears protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material such as clay, cat litter, or sawdust. Sweep up and place in a sealed container. Dispose of wastes as below. Do not store use prior to expiration date.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved State and local authorities.

**DIRECTIONS FOR USE (Cont'd.)**

**INDOOR USE AS A SPACE SPRAY UNDILUTED:** Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 90% less than 5 microns). Apply at a rate of 1 ounce per 1,000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. Sweep up and destroy fallen insects.

This method of application will control... and accessible stages of FLIES (including FRUIT FLIES), MOSQUITOES, SMALL FLYING MOTHS, GNATS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS, BOXELDER BUGS, EARWIGS, ANTS, and the following common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDER BEETLES, GARRULE BEETLES, DRUGSTORE BEETLES, ANGIOUMS, GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, YELLOW MEALWORMS, DARK MEALWORMS, GRAIN MITES, CADELLES, and RED FLOUR BEETLES.

**DRIED FRUIT PRODUCTS:** In storage or being processed, for control of CONFUSED FLOUR BEETLES, DRIED FRUIT BEETLES, INDIAN MEAL MOTHS, AND SAW-TOOTHED GRAIN BEETLES. Remove and destroy infested products. As noted above, disperse this product at a rate of 1 ounce per 1,000 cubic feet of space. Repeat application at this dosage in any quarter. Leave the room closed for one hour. Do not remain in treated area. Ventilate the area before reentry. Sweep up and destroy fallen insects.

**STORED FOOD:** Stored in multi-wall bags or unit bags in warehouses, storage rooms, flour mills, grain elevators, granaries, for control of ANGIOUMS, GRAIN MOTHS, CADELLES, CONFUSED FLOUR BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RICE WEEVILS, SAW-TOOTHED GRAIN BEETLES, YELLOW MEALWORMS, CHEESE MITES, CHEESE SKIPPERS, GARRULE BEETLES, GRAIN MITES, GRANARY WEEVILS, DARK MEALWORMS, RED FLOUR BEETLES. Remove and destroy infested products. As noted above, disperse this product at a rate of 1 ounce per 1,000 cubic feet of space. Repeat application at this dosage in any quarter. Leave the room closed for one hour. Do not remain in treated area. Ventilate the area before reentry. This application will kill accessible stages of those stored product pests listed above. Sweep up and destroy fallen insects.

**INDOOR USE AS A SPACE SPRAY DILUTED:** For use in conventional fogging equipment to control FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, AND GNATS. Cover or remove exposed food and handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute this concentrate plus 25 parts of water (see label mix). Apply at a rate of 1/2 fluid ounce per 1,000 cubic feet of space. Repeat application at this dosage in any quarter. Vacate treated area and ventilate before reentry. Do not remain in treated area when re-ventilation begins.

**ALLETHRIN OIL CONCENTRATE #525**

FOR USE BY TRAINED PERSONNEL ONLY.

Authorized by U.S.D.A. for use in Federally inspected meat and poultry plants.

**ACTIVE INGREDIENTS:**

ALLETHRIN	5.00%
RELATED COMPOUNDS	0.56%
PIPERONYL BUTOXIDE TECHNICAL*	25.00%
MINERAL OIL	69.44%
<b>TOTAL</b>	<b>100.00%</b>

\*Consists of 20% (Butylcarbityl) (6-propylpiperonyl) Ether and 5% related compounds

PRENTOX\* - Registered Trademark of Prentiss Drug & Chemical Co., Inc.

E.P.A. REG. NO. 655-053

E.P.A. EST. NO. 655-QA-1

**KEEP OUT OF REACH OF CHILDREN CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

If swallowed: Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. If inhaled: Remove victim to fresh air. If on skin: Remove contaminated clothing and wash affected areas with soap and water. If in eyes: Flush eyes with plenty of water. Call a physician immediately if irritation persists. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Effective in the control of the below-listed insects if the applicator follows the directions for use as set forth in this label:

- ANGIOUMS, GRAIN MOTHS - ANTS, BOXELDER BUGS - CADELLES - CHEESE MITES - CHEESE SKIPPERS - CIGARETTE BEETLES - CLOVER MITES - CONFUSED FLOUR BEETLES - DARK MEALWORMS - DEER FLIES - DRIED FRUIT BEETLES - DRUGSTORE BEETLES - EARWIGS - FRUIT FLIES (DROSOPHILA) - FUNGUS GNATS - GNATS - GRAIN MITES - GRANARY WEEVILS - HORN FLIES - HORNETS - HORSE FLIES - HOUSE FLIES - INDIAN MEAL MOTHS - LICE - MEDITERRANEAN FLOUR MOTHS - MOSQUITOES - MUSHROOM FLIES - RED FLOUR BEETLES - RICE WEEVILS - SAW-TOOTHED GRAIN BEETLES - SMALL FLYING MOTHS - SPIDER BEETLES - STABLE FLIES - TOBACCO MOTHS - WASPS - YELLOW MEALWORMS

**PRECAUTIONARY STATEMENTS**

**HAZARD TO HUMANS AND DOMESTIC ANIMALS CAUTION**

Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist. Avoid skin and eye contact. Wash thoroughly after handling and before eating or smoking. Do not allow children or pets to contact treated surfaces until spray has dried. Do not use on plants. Do not allow spray to contact food, feedstuffs, or water supplies. Thoroughly wash dishes and food handling utensils contaminated with this product. Remove pets and cover fish bowls before spraying.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

**PHYSICAL HAZARDS**

Do not use or store near heat or open flame. Exposure to temperatures above 130 F may cause bursting.



Manufactured by  
**prentiss**  
Drug & Chemical Co., Inc.

Plant: Kaolin Road  
Sandersville, GA. 31082  
Office: C.B. 2000  
Floral Park, N.Y. 11001

821

