

prentox[®]

ACCEPTED

AUG 18 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 655-663

ALLETHRIN OIL CONCENTRATE #525

FOR USE BY TRAINED PERSONNEL ONLY.

Authorized by U.S.D.A. for use in Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

ALLETHRIN	5.00%
RELATED COMPOUNDS	0.56%
PIPERONYL BUTOXIDE TECHNICAL*	25.00%
INERT INGREDIENT	69.44%
TOTAL	100.00%

*Consists of 20% (Butylcarbityl) (6-propylpiperonyl) ether and 5% related compounds.

PRENTOX[®]—Registered Trademark of Prentiss Drug & Chemical Co., Inc.

E.P.A. REG. NO. 655-663

E.P.A. EST. NO. 655-GA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If **swallowed**: Call a physician or Poison Control Center immediately. **DO NOT INDUCE VOMITING**. Vomiting may cause aspiration pneumonia. If **inhaled**: Remove victim to fresh air. If **on skin**: Remove contaminated clothing and wash affected areas with soap and water. If in **eyes**: Flush eyes with plenty of water. Call a physician immediately if irritation persists. **SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

Effective in the control of the below-listed insects if the applicator follows the directions for use as set forth in this label.

ANGOUMOIS GRAIN MOTHS — ANTS BOXELDER BUGS — CADELLES — CHEESE MITES — CHEESE SKIPPERS — CIGARETTE BEETLES — CLOVER MITES — CONFUSED FLOUR BEETLES — DARK MEALWORMS — DEER FLIES — DRIED FRUIT BEETLES — DRUGSTORE BEETLES — EARWIGS — FRUIT FLIES (DROSOPHILA) — FUNGUS GNATS — GNATS — GRAIN MITES — GRANARY WEEVILS — HORN FLIES — HORNETS — HORSE FLIES — HOUSE FLIES — INDIAN MEAL MOTHS — LICE — MEDITERRANEAN FLOUR MOTHS — MOSQUITOES — MUSHROOM FLIES — RED FLOUR BEETLES — RICE WEEVILS — SAW-TOOTHED GRAIN BEETLES — SMALL FLYING MOTHS — SPIDER BEETLES — STABLE FLIES — TOBACCO MOTHS — WASPS — YELLOW MEALWORMS.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist. Avoid skin and eye contact. Wash thoroughly after handling and before eating or smoking.

Do not allow children or pets to contact treated surfaces until spray has dried. Do not use on plants. Do not allow spray to contact food, feedstuffs or water supplies. Thoroughly wash dishes and food handling utensils contaminated with this product. Remove pets and cover fish bowls before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use, or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE (Cont'd)

SURFACE SPRAY: To insure maximum control of crawling insects (ANTS, BOXELDER BUGS, CLOVER MITES, DRUGSTORE BEETLES, EARWIGS, LICE, SPIDER BEETLES), dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well and apply as a coarse wetting spray using approximately 1 gallon of spray mixture to 750 square feet of surface. Spray walls, floor and other surfaces of bins and storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Treat unloading, handling and processing areas and inside conveying processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

For in transit protection, clean and spray truck beds, boxcars, and ship's hold before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked.

To control INDIAN MEAL MOTHS, and ANGOUMOIS GRAIN MOTHS in stored grain, apply to the surface of binned grain at a rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of moths and continuing until early fall. In case of severe infestation, break up webbing with rake before spraying and make a second application in two weeks.

As an aid in control of MOSQUITOES, GNATS, WASPS AND SKIPPERS: treat walls, ceilings, mouldings, screens, door and window frames, beams, light cords and similar resting places.

IN MUSHROOM PRODUCTION AND PROCESSING: To control MUSHROOM FLIES or FUNGUS GNATS in growing houses, dilute 1 part of concentrate plus 29 parts Methylene Chloride and mix well and apply at a rate of 1 ounce per 1,000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and should be directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packaging and canning plants, use at a rate of one-half ounce per 1,000 cubic feet of space.

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS: Dilute one part concentrate with 5 parts base oil (refined kerosene) and use with a ULV spraying device at the rate of one ounce per 1,000 cubic feet of room space. Apply the spray into cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Repeat as necessary for control of exposed or accessible stages of ROACHES, SILVERFISH, WATERBUGS, EARWIGS, ANTS, SPIDERS, FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS. Before using, exposed food products and packaging materials must be covered or removed. After use, equipment and utensils must be thoroughly rinsed with potable water before operations are resumed. Use as a space spray only when plant is not in operation.

OUTDOOR GROUND APPLICATION, UNDILUTED: To control ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS. For best results, apply when there is a light breeze of approximately 5 mph and temperature is cool (75° or less). With a ULV spraying device spray at a dosage rate of 5 ounces per minute up-wind of the area to be controlled. If the equipment is truck mounted, keep the speed to approximately 5 mph with a swath width of no more than 300 feet.

To control MOSQUITO LARVAE, dilute at the rate of 9 ounces per 50 gallons of oil and apply at the rate of 20 to 25 gallons per acre. Apply as a coarse wetting spray to the border or on shallow water of wet lands, swamps and marshes where mosquito larvae are known to breed.

TEMPORARY REDUCTION OF ANNOYANCE from FLIES, MOSQUITOES and SMALL FLYING MOTHS OUTDOORS. The above-mentioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in public gathering places (drive-in restaurants, drive-in theaters, golf courses, municipalities, parks, playgrounds and recreational areas). Direct application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. Application should be made prior to attendance. Repeat as necessary.

In additional outdoor areas (corrals, feed lots, swine lots and zoos), cover water, drinking fountains and animal feed before use. Fill area with mist, directing application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage in zoos, avoid exposure of reptiles to the product. Repeat as necessary.

FOR USE ON HORSES OR DOGS: To protect horses and dogs from HORN FLIES, HOUSE FLIES, MOSQUITOES, AND GNATS, dilute 1 part of concentrate plus 49 parts oil (refined kerosene) and mix well. Apply a light mist, sufficient to wet the tips of the hair. To control STABLE FLIES, HORSE FLIES, and DEER FLIES on horses, apply 2 ounces per adult animal, sufficient to wet the hair but not to soak the hide. Repeat treatment once or twice daily or at intervals to give continued protection.

To be applied on horses not intended for consumption.



Manufactured by

prentiss

Drug & Chemical Co., Inc.

Plant: Kaolin Road
Sandersville, GA. 31082

Office: C.B. 2000
Floral Park, N.Y. 11001