

LEFT 10-11

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Open dumping prohibited.

STORAGE: If container is damaged: STOP any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. ~~Dispose of on to a shovel and put the sweepings into a salvage drum.~~ Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PLASTIC CONTAINERS ONLY: DO NOT STORE IN CAR TRUNK WHERE TEMPERATURES MAY EXCEED 120°F.

BEST DOCUMENT AVAILABLE

prentox[®]

Pyronyl Oil Concentrate #12294

For Use by Trained Personnel Only

Authorized by U.S.D.A. for use in Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide Technical*	2.00%
N-Octyl bicycloheptane dicarboximide	2.94%
Mineral Oil	90.06%
Petroleum Distillates	4.00%
TOTAL	100.00%

*Consists of 1.60% (butylcarbityl) (6-propylpiperonyl) Ether and 0.40% related compounds

Pyronyl Oil Concentrate #12294 is effective in the control of the below-listed insects if the applicator follows the directions for use as set forth in this label.

ANGOUMOIS GRAIN MOTHS
ANTS
BOXELDER BUGS
CADELLES
CHEESE MITES
CHEESE SKIPPERS
CIGARETTE BEETLES
CLOVER MITES
CONFUSED FLOUR BEETLES
DARK MEALWORMS
DEER FLIES
DRIED FRUIT BEETLES
DRUGSTORE BEETLES

EARWIGS
FRUIT FLIES
(DROSOPHILA)
FUNGUS GNATS
GNATS
GRAIN MITES
GRANARY WEEVILS
HORN FLIES
HORNETS
HORSE FLIES
HOUSE FLIES
INDIAN MEAL MOTHS
LICE

MEDITERRANEAN FLOUR MOTHS
MOSQUITOES
MUSHROOM FLIES
RED FLOUR BEETLES
RICE WEEVILS
SAW-TOOTHED GRAIN BEETLES
SMALL FLYING MOTHS
SPIDER BEETLES
STABLE FLIES
TOBACCO MOTHS
WASPS
YELLOW MEALWORMS

KEEP OUT OF REACH OF CHILDREN
CAUTION

E.P.A. Reg. No. 855-884

E.P.A. EST. NO. 855-88-1

NET CO



Manufactured by

prentiss

One Chemical Building

LOT #

Plant: Kroll Road
Sandersville, GA. 31082

Office: 333 7th Avenue

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION: HARMFUL IF SWALLOWED. IF INGESTED DO NOT INDUCE VOMITING. CALL A PHYSICIAN AT ONCE. DO NOT APPLY DIRECTLY TO FOOD.
ENVIRONMENTAL HAZARDS
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage or disposal. Pesticide disposal: Do not dump or pour. Pesticide disposal: Pesticide spray mixture residues that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
CONTAINER DISPOSAL Containers: (a) Rinse or more: Rinse at container and offer for reconditioning or (b) Recycle: Rinse (or equivalent) and offer for recycling or conditioning for disposal in approved landfill or bury in a safe place. Containers less than 20 gallons—As in (b) above.
REMARKS—Consult Federal, State or local disposal authorities for approved alternative procedures.

INDOOR USE AS A SPACE SPRAY UNLIMITED The use of this product in food processing or food handling establishments should be confined to time periods when the plants or facilities are not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.
Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray in droplets over 50 microns in diameter and 80% less than 30 microns. Apply at a rate of 1 ounce per 1,000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area behind constructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. Sweep up and destroy fallen insects.
This method of application is for control of exposed, accessible stages of FLIES (including FRUIT FLIES), MOSQUITOES, SMALL FLYING MOTHS, GNATS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS, BOXELDER BUGS, EARWIGS, ANTS and the following common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANGOUMOIS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, YELLOW MEALWORMS, DARK MEALWORMS, GRAIN MITES, CADELLES and RED FLOUR BEETLES.
DRIED FRUIT PRODUCTS in storage or being processed for control of CONFUSED FLOUR BEETLES, DRIED FRUIT BEETLES, INDIAN MEAL MOTHS and SAW TOOTHED GRAIN BEETLES. Remove and destroy infested products. As noted above, use this product as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use one-half ounce per 1,000 cubic feet of space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not remain in treated areas and ventilate the area before re-entry. Sweep up and destroy fallen insects.
STORED FOOD (Stored in multi-wall bags or cloth bags) in warehouses, storage rooms, flour mills, grain elevators or granaries for control of ANGOUMOIS GRAIN MOTHS, CADELLES, CONFUSED FLOUR BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RICE WEEVILS, SAW TOOTHED GRAIN BEETLES, YELLOW MEALWORMS, CHEESE MITES, CHEESE SKIPPERS, CIGARETTE BEETLES, GRAIN MITES, GRANARY WEEVILS, DARK MEALWORMS, RED FLOUR BEETLES. Remove and destroy infested products. As noted above, disperse this product first by directing spray into corners, under pallets and around stacks of packaged foods hitting as many insects as possible and then treat as a space spray.
Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose. Use a dosage of one-half ounce per 1,000 cubic feet of space. Do not remain in treated areas and ventilate area before re-entry. This application is for accessible exposed stages of these stored product pests listed above. Sweep up and destroy fallen insects.

INDOOR USE AS A SPACE SPRAY LIMITED For use in conventional mechanical fogging equipment to control FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS and GNATS. Cover or remove exposed food and food handling equipment before application. Do not use in ventilated systems.

BEST DOCUMENT AVAILABLE

prentox[®]

Pyronyl Oil Concentrate #12294

For Use by Trained Personnel Only

Authorized by U.S.D.A. for use in Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide Technical	2.00%
N-Octyl bicycloheptane dicarboximide	2.84%
Mineral Oil	90.00%
Petroleum Distillates	4.00%
TOTAL	100.00%

*Consists of 1.60% (butylcarbityl) (6-propylpiperonyl) Ether and 0.40% related compounds

Prentox[®] is effective in the control of the below-listed insects if the applicator follows the directions for use as set forth in this label

- | | | |
|------------------------|-------------------|---------------------------|
| ANGUINOUS GRAIN MOTHS | EARWIGS | MEDITERRANEAN FLOUR MOTHS |
| ANTS | FRUIT FLIES | MOSQUITOES |
| BOXELDER BUGS | (DIPTERA) | MUSHROOM FLIES |
| CADELLES | FUNGUS GNATS | RED FLOUR BEETLES |
| CHEESE MITES | GNATS | RICE WEEVILS |
| CHEESE SKIPPERS | GRAIN MITES | SAW-TOOTHED GRAIN BEETLES |
| CIGARETTE BEETLES | GRANARY WEEVILS | SMALL FLYING MOTHS |
| CLOVER MITES | HORN FLIES | SPIDER BEETLES |
| CONFUSED FLOUR BEETLES | HORNETS | STABLE FLIES |
| DARK MEALWORMS | HORSE FLIES | TOBACCO MOTHS |
| DEER FLIES | HOUSE FLIES | WASPS |
| DRIED FRUIT BEETLES | INDIAN MEAL MOTHS | YELLOW MEALWORMS |
| DRUGSTORE BEETLES | LICE | |

KEEP OUT OF REACH OF CHILDREN
CAUTION

E.P.A. Reg. No. 655-864

E.P.A. EST. NO. 856-8A-1

NET CO



Manufactured by

prentiss

LOT #

Plant: Krollin Road
Sandersville, GA. 31082

Office: 363 7th Avenue

Directions for Use (continued)

SPRAYS: To insure maximum control of crawling insects (ANTS, BEETLES, COCKROACHES, CRICKETS, FLIES, GRASSHOPPERS, LICE, MOSQUITOES, TERMITES) dilute 1 part concentrate plus 99 parts petroleum solvent and mix well. Apply as a contact spray using approximately 1 gallon of spray mixture to 750 square feet of surface. Spray walls, floor and other exposed surfaces and storage and handling areas thoroughly being especially careful to treat cracks, crevices and similar hiding places. Treat unloading equipment, spraying areas and inside conveying processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

FOR INDOOR USE: Clean and spray such beds, boxes and shop's used for live loading. Thoroughly treat walls and floors and spray each layer of boxes or containers lightly as stacked.

TO CONTROL: MEAL MOTHS, and ANGUINOUS GRAIN MOTHS in stored grain apply to the surface of binned grain at a rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of moths and continuing until early fall. In case of severe infestation break up mounding with rake before spraying and make a second application in two weeks.

As an aid in control of MOSQUITOES, GNATS, WASPS AND SKIPPERS treat walls, ceilings, moldings, screens, door and window frames, beams, light cords and similar hiding places.

INDIAN MEALWORMS: Cattle barns, horse barns, poultry houses, swine houses, etc. As a space spray diluted for use in conventional mechanical fogging equipment to kill FLEES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well.

Apply at the rate of 2 fluid ounces per 1,000 cubic feet of space above the animals. Direct spray toward the upper portions of the enclosure filling the room with mist. Keep enclosed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

TO CONTROL MOSQUITOES AND MOSQUITOES: To control MOSQUITOES AND FUNGUS GNATS in growing houses dilute 1 part of concentrate plus 29 parts kerosene Chloride and mix well and apply at a rate of 1 ounce per 1,000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and should be directed slightly upward over them. Circulating fans should be left in operation but for best results refrigeration should be turned off in packaging and canning plants use at a rate of one-half ounce per 1,000 cubic feet of space.

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS: Use undiluted and use with a ULV spraying device at the rate of one ounce per 1,000 cubic feet of room space. Apply the spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Repeat as necessary for control of excess or accessible stages of ROACHES, SILVERFISH, WATERBUGS, EARWIGS, ANTS, SPIDERS, FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS. Before using exposed food products and packaging materials must be covered or removed. After use equipment, and utensils must be thoroughly rinsed with potable water before operations are resumed. Use as a space spray only when plant is not in operation.

OUTDOOR BARNY APPLICATION: UNDILUTED To control A RUIT MOSQUITOES, GNATS AND SMALL FLYING MOTHS. For best results apply when there is a light breeze of approximately 5 mph and temperature is cool (70° or less). With ULV spraying device spray at a dosage rate of 5 ounces per minute up wind of the area to be controlled. If the equipment is truck mounted, keep the speed to approximately 5 mph with a swath width of no more than 300 feet.

To control MOSQUITO LARVAE dilute at the rate of 15 ounces per 50 gallons of oil and apply at the rate of 20 to 25 gallons per acre. Apply as a coarse wetting spray to the border or on shallow water of wet lands, swamps and marshes where mosquito larvae are known to breed.

TEMPORARY REDUCTION OF ANNOYANCE: FROM FLIES, MOSQUITOES AND SMALL FLYING MOTHS OUTDOORS. The above-mentioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in public gathering places (drive-in restaurants, drive-in theaters, golf courses, municipalities, parks, playgrounds and recreational areas). Direct application into tall grass, shrubbery and around lawns where these pests may cover or rest apply while air is still. Avoid wetting foliage in park areas prior to attendance. Repeat as necessary.

In additional outdoor areas (corridors, food lots, some lots and zoos) cover water drinking fountains and animal feed before use. Fill area with mist directing application into tall grass, shrubbery and around lawns where these pests may hover or rest apply while air is still. Avoid wetting foliage in park areas and exposure of reptiles to the product. Repeat as necessary.

FOR USE ON ANIMALS: To protect beef and dairy cattle, horse and dogs from MOSQUITOES, HOUSE FLIES, MOSQUITOES AND GNATS dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. To control STABLE FLIES, HORSE FLIES AND DEER FLIES on beef and dairy cattle and horses apply 2 ounces per adult animal sufficient to wet the hair but not to soil the skin. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

To control the more stages of the COCKROACHES on beef and dairy cattle dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tips of the hair. Repeat treatment once or twice daily or at the first sign of infestation.

BEST DOCUMENT AVAILABLE

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

- If swallowed - Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.
- If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.
- If on skin - Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes - Flush eyes with plenty of water. Call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DES. 21

LEFT PANEL

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist. Avoid skin and eye contact. Wash thoroughly after handling and before eating or smoking.

Do not allow children or pets to contact treated surfaces until spray has dried. Do not use on plants. Do not allow spray to contact food, feedstuffs or water supplies. Thoroughly wash dishes and food handling utensils contaminated with this product. Remove pets and cover fish bowls before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams and ponds. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use, or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

BEST COPY AVAILABLE

LEFT PAGE

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Open dumping prohibited.

STORAGE: If container is damaged: STOP any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PLASTIC CONTAINERS ONLY: DO NOT STORE IN CAR TRUNK WHERE TEMPERATURES MAY EXCEED 120°F.

BEST DOCUMENT AVAILABLE