

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist. Avoid skin and eye contact. Wash thoroughly after handling and before eating or smoking.

Do not allow children or pets to contact treated surfaces until spray has dried. Do not use on plants. Do not allow spray to contact food, feedstuffs or water supplies. Thoroughly wash dishes and food handling utensils contaminated with this product. Remove pets and cover fish bowls before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use or store near heat-exposed items. Exposure to temperatures above 137°F may cause burning.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticides and other protective gear. On clean up of liquid leaks, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as follows: Place any leaking container into a salvage drum or glass container. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

DIRECTIONS FOR USE (Cont'd.)

INDOOR USE AS A SPACE SPRAY UNDILUTED: The use of this product in kind processing or food handling establishments should be confined to time periods when the plants or facilities are not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Close windows and doors and shut off ventilating systems. Use a fogger or sprayer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at a rate of 1 ounce per 1,000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area behind obstructions. Keep the area closed for at least 15 minutes. Ventilate the treated area and ventilate before reoccupying. Repeat treatment as necessary. Sweep up and destroy fallen insects.

For a method of application and control of exposed accessible stages of FLIES (including FRUIT FLIES), MOSQUITOES, SMALL FLYING MOTHS, GNATS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS, BOXELDER BUGS, EARTHWIGS, ANTS and the following common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANGIOUMOUS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, YELLOW MEAL WORMS, DARK MEAL WORMS, GRAIN MITES, CADELLES and RED FLOUR BEETLES.

DRIED FRUIT PRODUCTS: In storage or being processed for control of CONFUSED FLOUR BEETLES, DRIED FRUIT BEETLES, INDIAN MEAL MOTHS, AND SAW-TOOTHED GRAIN BEETLES. Remove and destroy infested products. As noted above, apply this product as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use one-half ounce per 1,000 cubic feet of space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not remain in treated areas. Ventilate the area before re-entry. Sweep up and destroy fallen insects.

STORED FOOD: (Stored in multi-wall bags or cloth bags) in warehouses, storage rooms, flour mills, grain elevators, granaries for control of ANGIOUMOUS GRAIN MOTHS, CADELLES, CONFUSED FLOUR BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RICE WEEVILS, SAW-TOOTHED GRAIN BEETLES, YELLOW MEAL WORMS, CHEESE MITES, CHEESE SKIPPERS, CIGARETTE BEETLES, GRANARY WEEVILS, DARK MEAL WORMS, RED FLOUR BEETLES. Remove and destroy infested products. As noted above, apply this product by directing spray into corners, under pallets and around stacks of packaged foods hitting as many insects as possible and then treat as a secure facility.

Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose. Use a dosage of one-half fluid ounce per 1,000 cubic feet of space. Do not remain in treated areas and ventilate area before re-entry. This application will kill accessible exposed stages of those stored product pests listed above. Sweep up and destroy fallen insects.

prentox[®]

Pyronyl Oil Concentrate #12294

For Use by Trained Personnel Only

Authorized by U.S.D.A. for use in Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide Technical*	2.00%
N-Octyl bicycloheptene dicarboximide	2.94%
Mineral Oil	90.04%
Petroleum Distillates	4.00%
TOTAL	100.00%

*Consists of 1.60% (butylcarbityl) (6-propylpiperonyl) Ether and 0.40% related compounds

PRENTOX[®]—Registered Trademark of Prentiss Drug & Chemical Co., Inc.

Effective in the control of the below-listed insects if the applicator follows the directions for use as set forth in this label:

ANGIOUMOUS GRAIN MOTHS — ANTS — BOXELDER BUGS — CADELLES — CHEESE MITES — CHEESE SKIPPERS — CIGARETTE BEETLES — CLOVER MITES — CONFUSED FLOUR BEETLES — DARK MEAL WORMS — DEER FLIES — DRIED FRUIT BEETLES — DRUGSTORE BEETLES — EARTHWIGS — FRUIT FLIES — (DROSOPHILA) — FUNGUS GNATS — GNATS — GRANARY WEEVILS — HORN FLIES — HORNETS — HORROR FLIES — HOUSE FLIES — INDIAN MEAL MOTHS — LICE — MEDITERRANEAN FLOUR MOTHS — MOSQUITOES — MUSHROOM FLIES — RED FLOUR BEETLES — RICE WEEVILS — SAW-TOOTHED GRAIN BEETLES — SMALL FLYING MOTHS — SPIDER BEETLES — STABLE FLIES — TOBACCO MOTHS — WASPS — YELLOW MEAL WORMS

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

Harmful: Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. If inhaled, Remove victim to fresh air. If on skin, Remove contaminated clothing and wash affected areas with soap and water. If in eyes, Flush eyes with plenty of water. Call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. REG. NO. 855-664

BEST DOCUMENT AVAILABLE

E.P.A. EST. NO. 855-GA-1

NET CONTENTS

LOT #



Manufactured by
prentiss
Drug & Chemical Co., Inc.

Plant: Kaolin Road
Sandersville, GA. 31082
Office: C.B. 2000
Floral Park, N.Y. 11001

INDOOR USE AS A SPACE SPRAY: For control of the following pests: SMALL FLYING MOTHS AND GNATS, food handling surfaces. Close room ventilating system equipment. Dilute (diluted kerosene) and mix well. Apply 1000 cubic feet filling the room with minutes. Vacate treated area and re-treatment when reinfestation occurs.

SURFACE SPRAY: To insure maximum BOXELDER BUGS, CLOVER MITES, LICE, SPIDER BEETLES, dilute 1 part kerosene) and mix well and apply approximately 1 gallon of spray mixture walls, floor and other surfaces of building thoroughly, being especially careful hiding places. Treat unloading, handling, conveying processing and handling of the base of heavy machinery and equipment possible.

For in-transit protection, clean and spray before loading. Thoroughly treat wall, cartons or containers lightly as stars.

To control INDIAN MEAL MOTHS in stored grain, apply to the surface of bins square feet. Apply at monthly intervals continuing until early fall. In case of infestation, rake before spraying and make a seal. As an aid in control of MOSQUITOES, walls, ceilings, moldings, screens, cords and smaller nesting places.

ANIMAL QUARTER USE: (Cattle barn, houses, zoos) As a space spray dilute fogging equipment to kill FLIES, MOSQUITOES, GNATS, dilute 1 part concentrate and mix well. Apply at the rate of 2 m above the animals. Direct spray toward filling the room with mist. Keep area treated area and ventilate before reinfestation occurs.

IN MUSHROOM PRODUCTION AS ROOM FLIES OR FUNGUS GNATS: concentrate plus 29 parts Methylene of 1 ounce per 1,000 cubic feet of space away from beds and should be circulating fans should be left in operation should be turned off in packaging and once per 1,000 cubic feet of space. FOR USE IN FEDERALLY INSPECTED undiluted and use with a ULV spray 1,000 cubic feet of room space. Apply woodwork, walls, floors, underneath where these insects shelter. Contact necessary for control of exposed SILVERFISH WATERBUGS, EARTHWIGS, MOSQUITOES, SMALL FLY, exposed food products and feed removed. After use equipment and potable water before operations and plant is not in operation.

OUTDOOR GRASSHOPPER APPLICATION: MOSQUITOES, GNATS AND SMALL FLYING MOTHS. Apply when there is a light breeze of cool 175° or less. With a ULV spray 1 ounce per minute upwind of the truck mounted. Keep the speed to 40 or more than 300 feet.

To control MOSQUITO LARVAE, dilute of oil and apply at the rate of 20 to waiting spray to the border or on marshes where mosquito larvae are.

TEMPORARY REDUCTION OF SMALL FLYING MOTHS OUT: for outdoor ground application with from these pests in public gathering places, golf courses, municipal areas. Direct application into the these pests may hover or rest and Application should be made prior in additional outdoor areas (corn water, drinking fountains and an directing application into tall grass) pests may hover or rest, apply with avoid exposure of reptiles to the.

FOR USE ON ANIMALS: To protect from HORSE FLIES, HOUSE FLIES, part of concentrate plus 29 parts light mist sufficient to wet the HORSE FLIES and DEER FLIES 1 ounce per adult animal, apply. Repeat treatment once or twice protection.

To control the mobile stages of cattle, apply the spray to thorax, animal and repeat 2 to 3 weeks.



prentox[®]

Pyronyl Oil Concentrate #12294

For Use by Trained Personnel Only

Authorized by U.S.D.A. for use in Federally Inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide Technical*	2.00%
N-Octyl bicycloheptene dicarboximide	2.94%
Mineral Oil	90.06%
Petroleum Distillates	4.00%
TOTAL	100.00%

*Consists of 1.60% (butylcarbityl) (6-propylpiperonyl) Ether and 0.40% related compounds

PRENTOX[®]—Registered Trademark of Prentiss Drug & Chemical Co., Inc.

Effective in the control of the below-listed insects if the applicator follows the directions for use as set forth in this label.

ANGOUMOIS GRAIN MOTHS — ANTS — BOXELDER BUGS — CADELLES — CHEESE MITES — CHEESE SKIPPERS — CIGARETTE BEETLES — CLOVER MITES — CONFUSED FLOUR BEETLES — DARK MEAL WORMS — DEER FLIES — DRIED FRUIT BEETLES — DRUGSTORE BEETLES — EARWIGS — FRUIT FLIES — (DROSOPHILA) — FUNGUS GNATS — GNATS — GRAIN MITES — GRANARY WEEVILS — HORN FLIES — HORNETS — HORSEFLIES — HOUSE FLIES — INDIAN MEAL MOTHS — LICE — MEDITERRANEAN FLOUR MOTHS — MOSQUITOES — MUSHROOM FLIES — RED FLOUR BEETLES — RICE WEEVILS — SAW-TOOTHED GRAIN BEETLES — SMALL FLYING MOTHS — SPIDER BEETLES — STABLE FLIES — TOBACCO MOTHS — WASPS — YELLOW MEAL WORMS

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

Not Allowed: Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. If inhaled, Remove victim to fresh air. If on skin, Remove contaminated clothing and wash affected areas with soap and water. If in eyes, Flush eyes with plenty of water. Call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. REG. NO. 855-664

E.P.A. EST. NO. 855-GA-1

BEST DOCUMENT AVAILABLE

NET CONTENTS

LOT #



Manufactured by
prentiss
Drug & Chemical Co., Inc.

Plant: Kaolin Road
Sandersville, GA. 31082
Office: C.B. 2000
Floral Park, N.Y. 11001

625-664

INDOOR USE AS A SPACE SPRAY DILUTED: To control SMALL FLYING MOTHS AND GNATS: Close room and shut off all ventilation system equipment. Dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

SURFACE SPRAY: To insure maximum control of crawling insects (ANTS, BOXELDER BUGS, CLOVER MITES, DRUGSTORE BEETLES, EARWIGS, LICE, SPIDER BEETLES) dilute 1 part concentrate plus 29 parts of refined kerosene and mix well and apply as a coarse wetting spray using approximately 1 gallon of spray mixture to 750 square feet of surface. Spray walls, floor and other surfaces of bins and storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Treat unloading, handling and processing areas and inside conveying processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

For transit protection, clean and spray truck beds, boxcars and ship holds before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly, as stacked.

To control INDIAN MEAL MOTHS and ANGOUMOIS GRAIN MOTHS in stored grain, apply to the surface of binned grain at the rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of moths and continuing until they fall in case of severe infestation. Break up webbing with rake before spraying and make a second application in two weeks.

As an aid in control of MOSQUITOES, GNATS, WASPS and SKIPPERS, treat walls, ceilings, moldings, screens, door and window frames, beams, light cords and similar resting places.

ANIMAL QUARTER USE: (Cattle barns, horse barns, poultry houses, swine houses, zoos). As a space spray diluted for use in conventional mechanical fogging equipment to kill FLIES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS, dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 2 fluid ounces per 1,000 cubic feet of space above the animals. Direct spray towards the upper portions of the enclosure filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

IN MUSHROOM PRODUCTION AND PROCESSING: To control MUSHROOM FLIES or FUNGUS GNATS in growing houses, dilute 1 part of concentrate plus 29 parts Methylene Chloride and mix well and apply at rate of 1 ounce per 1,000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and should be directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packaging and canning plants, use at rate of one-half ounce per 1,000 cubic feet of space.

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS: Use undiluted and use with a ULV spraying device at the rate of one ounce per 1,000 cubic feet of room space. Apply the spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Repeat as necessary for control of exposed or accessible stages of ROACHES, SILVERFISH, WATERBUGS, EARWIGS, ANTS AND GNATS, FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS. Before using exposed food products and packaging materials must be covered or removed. After use, equipment and utensils must be thoroughly washed with potable water before operations are resumed. Use as a space spray only when plant is not in operation.

OUTDOOR GROUND APPLICATION, UNDILUTED: To control ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS: For best results apply when there is a light breeze of approximately 5 mph and temperature is cool (75° or less). With a ULV spraying device spray at a dosage rate of 5 ounces per minute upwind of the area to be controlled. If the equipment is truck mounted, keep the speed to approximately 5 mph with a swath width of no more than 300 feet.

To control MOSQUITO LARVAE, dilute at the rate of 15 ounces per 50 gallons of oil and apply at the rate of 20 to 25 gallons per acre. Apply as a coarse wetting spray to the border or on shallow water of wetlands, swamps and marshes where mosquito larvae are known to breed.

TEMPORARY REDUCTION OF ANNOYANCE FROM FLIES, MOSQUITOES AND SMALL FLYING MOTHS OUTDOORS: The aforementioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in public gathering places (drive-in restaurants, drive-in theaters, golf courses, municipalities, parks, playgrounds and recreational areas). Direct application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. Application should be made prior to attendance. Repeat as necessary. In additional outdoor areas (corridors, feed lots, swine lots and zoos), cover water, drinking fountains and animal feed before use. Fill area with mist direct application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage in zoos and exposure of reptiles to the product. Repeat as necessary.

FOR USE ON ANIMALS: To protect beef and dairy cattle, horses and dogs from HORSE FLIES, HOUSE FLIES, MOSQUITOES AND GNATS, dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply a light mist sufficient to wet the tip of the hair. To control STABLE FLIES, HORSE FLIES and DEER FLIES on beef and dairy cattle and horses, apply 2 ounces per adult animal, sufficient to wet the hair, but not to soak the hide. Repeat treatment once or twice daily or at intervals to give continued protection.

To control the mobile stages of BLOOD SUCKING LICE on beef and dairy cattle, apply the spray to thoroughly wet the hair. Use 2 ounces or less per animal and repeat 2 to 3 weeks as required.