

ACCEPTED

AUG 12 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 655-664

prentox®

Pyronyl™ Oil Concentrate #12294

For Use by Trained Personnel Only

Authorized by U.S.D.A. for use in Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide Technical*	2.00%
N-Octyl bicycloheptene dicarboximide	2.94%
Mineral Oil	90.06%
Petroleum Distillates	4.00%
TOTAL	100.00%

*Consists of 1.60% (butylcarbityl) (8-propylpiperonyl) ether and 0.40% related compounds.

PRENTOX® — Registered Trademark of Prentiss Drug & Chemical Co., Inc.

PYRONYL™ — Trademark of Prentiss Drug & Chemical Co., Inc.

Effective in the control of the below-listed insects if the applicator follows the directions for use as set forth in this label.

ANGOUMOIS GRAIN MOTHS — ANTS — BOXELDER BUGS — CADELLES — CHEESE MITES — CHEESE SKIPPERS — CIGARETTE BEETLES — CLOVER MITES — CONFUSED FLOUR BEETLES — DARK MEAL WORMS — DEER FLIES — DRIED FRUIT BEETLES — DRUGSTORE BEETLES — EARWIGS — FRUIT FLIES — (DROSOPHILA) — FUNGUS GNATS — GNATS — GRAIN MITES — GRANARY WEEVILS — HORN FLIES — HORNETS — HORSE FLIES — HOUSE FLIES — INDIAN MEAL MOTHS — LICE — MEDITERRANEAN FLOUR MOTHS — MOSQUITOES — MUSHROOM FLIES — RED FLOUR BEETLES — RICE WEEVILS — SAW-TOOTHED GRAIN BEETLES — SMALL FLYING MOTHS — SPIDER BEETLES — STABLE FLIES — TOBACCO MOTHS — WASPS — YELLOW MEAL WORMS

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. If inhaled: Remove person to fresh air. If on skin: Remove contaminated clothing and wash affected areas with soap and water. If in eyes: Flush eyes with plenty of water. Call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. REG. NO. 655-664

E.P.A. EST. NO. 655-GA-1

NET CONTENTS

LOT #



Manufactured by

prentiss

Drug & Chemical Co., Inc.

Plant: Kaelin Road
Sandersville, GA. 31082
Office: C.B. 2088
Floral Park, N.Y. 11001

PHYSICAL HAZARD STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. If swallowed, do not induce vomiting. Call a physician immediately.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying in meat packing plants, bakeries, and other food processing plants, wash with potable water all equipment, benches, shelving, etc., where exposed food will be handled. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause burning.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticides and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a salvage drum or other container. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

DIRECTIONS FOR USE (Cont'd.)

INDOOR USE AS A SPACE SPRAY UNDILUTED: The use of this product in food processing or food handling establishments should be confined to time periods when the plants or facilities are not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Close windows and doors and shut off ventilating systems. Use a fogger or venturi sprayer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 90% less than 30 microns). Apply at a rate of 1/2 ounce per 1,000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. Sweep up and destroy fallen insects.

This method of application will control exposed, accessible stages of FLIES (including FRUIT FLIES), MOSQUITOES, SMALL FLYING MOTHS, GNATS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS, BOXELDER BUGS, EARWIGS, ANTS and the following common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANGOUMOIS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, YELLOW MEAL WORMS, DARK MEAL WORMS, GRAIN MITES, CADELLES and RED FLOUR BEETLES.

DRIED FRUIT PRODUCTS in storage or being processed, for control of CONFUSED FLOUR BEETLES, DRIED FRUIT BEETLES, INDIAN MEAL MOTHS, AND SAW-TOOTHED GRAIN BEETLES: Remove and destroy infested products. As noted above, disperse this product as a fine mist in the air above the bags and bins. Do not apply to food directly. Use one-half ounce per 1,000 cubic feet of space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not reenter treated areas and ventilate the area before re-entry. Sweep up and destroy fallen insects.

STORED FOOD: (Stored in multi-wall bags or cloth bags) in warehouses, storage rooms, flour mills, grain elevators, granaries for control of ANGOUMOIS GRAIN MOTHS, CADELLES, CONFUSED FLOUR BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RICE WEEVILS, SAW-TOOTHED GRAIN BEETLES, YELLOW MEAL WORMS, CHEESE MITES, CHEESE SKIPPERS, CIGARETTE BEETLES, GRAIN MITES, GRANARY WEEVILS, DARK MEAL WORMS, RED FLOUR BEETLES: Remove and destroy infested products. As noted above, disperse this product first by directing spray into corners, under pallets, and around stacks of packaged goods killing as many insects as possible and then treat or, a space spray.

Infested stored products should be fumigated or treated by other effective methods using unapproved product intended for this purpose. Use dosage of one-half fluid ounce per 1,000 cubic feet of space. Do not reenter treated areas and ventilate area before re-entry. This application will kill accessible, exposed stages of these stored product pests listed above. Sweep up and destroy fallen insects.

DIRECTIONS FOR USE (Cont'd.)
INDOOR USE AS A SPACE SPRAY: In residential logging equipment to kill FLYING MOTHS, and GNATS: C food handling surfaces. Close room or ventilating system equipment. Dilute 1:1 (refined kerosene) and mix well. Apply at 1,000 cubic feet filling the room with mist minutes. Vacate treated area and ventilate before re-occupation.

SURFACE SPRAY: To insure maximum or BOXELDER BUGS, CLOVER MITES, DEER LICE, SPIDER BEETLES, dilute 1 part (or kerosene) and mix well and apply as approximately 1 gallon of spray mixture to walls, floor and other surfaces of bins (thoroughly, being especially careful to hiding places. Treat unloading, handling conveying processing and handling squab the base of heavy machinery and equipment possible.

For in transit protection, clean and spray in before loading. Thoroughly treat walls or cartons or containers lightly as stacked.

To control INDIAN MEAL MOTHS, and A stored grain, apply to the surface of binned square feet. Apply at monthly intervals with continuing until early fall. In case of severe infestation before spraying and make a second infestation. As an aid in control of MOSQUITOES, GNA walls, ceilings, moldings, screens, door cords and similar resting places.

ANIMAL QUARTER USE (Cattle barns, horse houses, zoos): As a space spray diluted for fogging equipment to kill FLIES, MOSQUITOES AND GNATS, dilute 1 part concentrate plus and mix well. Apply at the rate of 2 fluid oz above the animals. Direct spray towards the filling the room with mist. Keep areas close treated area and ventilate before re-occupation.

IN MUSHROOM PRODUCTION AND FRUIT ROOM FLIES OR FUNGUS GNATS: In or concentrate plus 28 parts Methylene Chloride of 1 ounce per 1,000 cubic feet of space. Fog away from beds and should be direct. Circulating fans should be left in operation, should be turned off. In packaging and control ounce per 1,000 cubic feet of space.

FOR USE IN FEDERALLY INSPECTED MEAT UNLUTED and use with a ULV spraying device 1,000 cubic feet of room space. Apply 1/2 oz per woodwork, walls, floors, underneath sinks, where these insects shelter. Contact as may necessary for control of exposed or so SILVERFISH, WATERBUGS, EARWIGS, A FLIES, MOSQUITOES, SMALL FLYING MOTHS exposed food products and packaging removed. Equipment and utensils in Feder plants must be washed with an effective cleanable water rinse after spraying. Use as a in operation.

OUTDOOR BOUND APPLICATION, USE MOSQUITOES, GNATS AND SMALL FLYING MOTHS: Apply when there is a light breeze of approx 1000 (70° or less). With a ULV spraying device ounces per minute up-wind of the area to be treated or - instead, keep the speed to approximately no more than 200 feet.

To control MOSQUITO LARVAE, dilute at the of oil and apply at the rate of 20 to 25 gal/watering spray to the border or on shallow marshes where mosquito larvae are known. TEMPORARY REDUCTION OF ANNOYANCE AND SMALL FLYING MOTHS OUTDOORS: for outdoor ground application will afford less from these pests in public gathering places theaters, golf courses, municipalities, parks, areas). Direct application into tall grass, shrub these pests may hover or rest, apply while a Application should be made prior to attend in additional outdoor areas (corns, feed to water, drinking fountains and animal feed) directing application into tall grass, shrubber pests may hover or rest, apply while air is still avoid exposure of reptiles to the product.

FOR USE ON ANIMALS: To protect beef and from HORN FLIES, HOUSE FLIES, MOSQUITO part of concentrate plus 28 parts oil (refined light mist, sufficient to wet the tips of the HORSE FLIES, and DEER FLIES on head and ounces per adult animal, sufficient to wet the Repeat treatment once or twice daily or 1 protection.

To control the mottle stages of BLOOD SUCK cattle, apply the spray to thoroughly wet the animal and repeat 2 to 3 weeks as required.

BEST AVAILABLE COPY