

prentox CHLORPYRIFOS D.D.V.P. MC



An Insecticide Concentrate for Formulating Use Only

ACTIVE INGREDIENTS:	
Chlorpyrifos [0,0-diethyl 0-(3,5,6-Trichloro-2-Pyridyl) Phosphorothioate]	15.00
2,2-Dichlorovinyl Dimethyl Phosphate	13.95
Related Compounds	1.05
Aromatic Petroleum Derivative Solvent	69.03
INERT INGREDIENTS	0.97
TOTAL	100.00

Directions For Use

Always read the label. Use this product only as directed on the label. Do not use this product if the label directions do not cover the pest you are treating. For more information on this product refer to our literature or contact your distributor.

KEEP OUT OF REACH OF CHILDREN



**POISON
DANGER**



STORAGE AND DISPOSAL


- 1 PROHIBITIONS**—Do not contaminate water bodies or dispose of containers or empty containers in or near bodies of water.
- 2 PESTICIDE DISPOSAL**—Containers must be disposed of in accordance with the label instructions. Do not use for other purposes. Federal Statute, 49 USC 3005, and the Resource Conservation and Recovery Act.
- 3 CONTAINER DISPOSAL**—a. Reread container and label directions. b. Triple rinse (or equivalent) and then recycle where permitted by local and/or state authorities.

Statement of Practical Treatment

If swallowed—If swallowed, induce vomiting immediately by drinking 1 cup of water. Do not induce vomiting if the person has had a seizure or is unconscious. Do not give anything by mouth to a person who is unconscious. If the person is unconscious, place them on their side. If the person is conscious, give them 1 cup of water every 15 minutes. If the person is unconscious, do not give anything by mouth. If the person is conscious, give them 1 cup of water every 15 minutes. If the person is unconscious, do not give anything by mouth.

If inhaled—Remove the person from the area. Apply artificial respiration if necessary. **If on skin**—Wash immediately with soap and water. **If in eyes**—Flush eyes with plenty of water. Call a physician immediately.

NOTE TO PHYSICIAN:
WARNING SYMPTOMS: Symptoms include weakness, headache, dizziness, blurred vision, increased salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.
TREATMENT: Atropine is the specific therapeutic antagonist of cholinergic parasympathetic nervous system stimulation. If there are signs of parasympathetic stimulation, Atropine 5-10 mg should be given at 10 minute intervals in doses of 1 to 2 mg until complete Atropinization has occurred. PAM has an antidotal and may be administered in conjunction with Atropine. Morphine is contraindicated. Clear chest by postural drainage. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to future small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase reactivity has been adequately determined by blood test.



E.P.A. REG. NO. 655-671 E.P.A. EST NO 655-NJ-1

NET CONTENTS _____ LOT NO _____



Manufactured by
prentiss
Drug & Chemical Co., Inc.

Plant: 338 Wilson Avenue
Newark, N.J. 07105
Office: 363 7th Avenue
New York, N.Y. 10001