

LEFT PAGE

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Open dumping prohibited.

STORAGE: If container is damaged: STOP any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required as prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. ~~Dispose of on a shovel and put the sweepings into a salvage drum.~~ Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use near or spill near heat or open flames.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PLASTIC CONTAINERS ONLY: DO NOT STORE IN CAR TRUNK WHERE TEMPERATURES MAY EXCEED 120°F.

BEST PRACTICES MANUAL 2

prentox[®] Pyrenyl Oil Concentrate #15

For Use by Trained Personnel Only

Authorized by U.S.D.A. for use in
Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide Technical [*]	5.00%
Mineral Oil	90.00%
Petroleum Distillates	4.00%
TOTAL	100.00%

^{*}Consists of 4.0% butylcarbitol-6-propylpiperonyl Ether and 1.0% related compounds

is effective in the control of the below listed insects if the applicator follows the directions for use as set forth in this label

- | | | | |
|------------------------|--------------------------|---------------------------|--------------------|
| ANGOUMOIS GRAIN MOTHS | DARK MEALWORMS | HORN FLIES | RICE WEEVILS |
| ANTS | DEER FLIES | HORNETS | SAW-TOOTHED |
| BOXELDER BUGS | DRUG STORE BEETLES | HORSE FLIES | CRAIN BEETLES |
| CADELLES | EARWIGS | HOUSE FLIES | SMALL FLYING MOTHS |
| CHEESE MITES | FRUIT FLIES (DROSOPHILA) | INDIAN MEAL MOTHS | SPIDER BEETLES |
| CHEESE SKIPPERS | FUNGUS GNATS | LICE | STABLE FLIES |
| CIGARETTE BEETLES | GNATS | MEDITERRANEAN FLOUR MOTHS | TOBACCO MOTHS |
| CLOVER MITES | GRANARY WEEVILS | MOSQUITOES | WASPS |
| CONFUSED FLOUR BEETLES | | MUSHROOM FLIES | YELLOW MEALWORMS |
| | | RED FLOUR BEETLES | |

E.P.A. Reg. No. 655-684

E.P.A. Est. No. 655-GA-1

NET CONTENTS

DIRECTIONS FOR USE

Apply to all surfaces of equipment, containers, and other objects in the plant, including floors, walls, ceilings, and equipment. Apply to all surfaces of equipment, containers, and other objects in the plant, including floors, walls, ceilings, and equipment.

LOT

DO NOT USE AS A SPACE SPRAY UNLESS FOR USE IN CONFINED AREAS. Apply to all surfaces of equipment, containers, and other objects in the plant, including floors, walls, ceilings, and equipment.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION
Harmful if swallowed. Avoid breathing vapor. Avoid contact with skin, eyes, or clothing. Wear mask or respirator of type recommended by NIOSH, National Institute for Occupational Safety and Health to give adequate protection against this material. Wear protective clothing. Wash exposed skin with soap after handling and before eating. Wash all contaminated clothing with soap and water separately from other clothing before reuse.
The use of this product in food processing or food handling establishments should be confined to time periods when the plants or facilities are not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.
ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
Store in original container. Do not use if container is damaged. Do not use if contents are empty. Do not use if contents are water. Do not use if contents are water. Do not use if contents are water.

DO NOT USE AS A SPACE SPRAY UNLESS FOR USE IN CONFINED AREAS. Apply to all surfaces of equipment, containers, and other objects in the plant, including floors, walls, ceilings, and equipment. Apply to all surfaces of equipment, containers, and other objects in the plant, including floors, walls, ceilings, and equipment.

BEST DOCUMENT AVAILABLE



Manufactured by

prentiss

Drug & Chemical Co., Inc.

Plant: Kaelin Road
Sandersville, GA. 31082

Office: 363 7th Avenue

Atlanta, GA. 30308

SURFACE SPRAY To control BOXELDER BUGS, COLEOPTER BEETLES, CRICKETS, and other insects, mix well and apply of spray mixture to all surfaces of bins and equipment. Careful to treat cracks, crevices, and other hiding places. Spray the equipment. Spray inside for in transit protective before loading. Spray cartons or containers. To control INDIAN MEAL MOTHS, apply to the surface. Apply at monthly intervals until early fall. As an aid in control of bed bugs, moldings, or similar resting places, apply to the surface. As a space spray, apply at the rate of 2 fluid ounces per 1000 cubic feet. Repeat the application as needed. Repeat the application as needed. Repeat the application as needed.

BEST DOCUMENT AVAILABLE

prentox[®] Pyrenyl Oil Concentrate #15

For Use by Trained Personnel Only

Authorized by U.S.D.A. for use in
Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide Technical*	5.00%
Mineral Oil	90.00%
Petroleum Distillates	4.00%
TOTAL	100.00%

*Consists of 4-(n-butylcarbityl)-(6-propylpiperonyl) Ether and 10% related compounds

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

is effective in the control of the below listed insects if the applicator follows the directions for use as set forth in this label

- | | | | |
|------------------------|---------------------|---------------------------|--------------------|
| ANGOUMOIS GRAIN MOTHS | DARK MEALWORMS | HORN FLIES | RICE WEEVILS |
| ANTS | DEER FLIES | HORNETS | SAW TOOTHED |
| BOXELDER BUGS | DRIED FRUIT BEETLES | HORSE FLIES | GRAIN BEETLES |
| CADELLES | DRUGSTORE BEETLES | HOUSE FLIES | SMALL FLYING MOTHS |
| CHEESE MITES | EARWIGS | INDIAN MEAL MOTHS | SPIDER BEETLES |
| CHEESE SKIPPERS | FRUIT FLIES | LICE | STABLE FLIES |
| CIGARETTE BEETLES | (DROSOPHILA) | MEDITERRANEAN FLOUR MOTHS | TOBACCO MOTHS |
| CLOVER MITES | FUNGUS GNATS | MOSQUITOES | WASPS |
| CONFUSED FLOUR BEETLES | GNATS | MUSHROOM FLIES | YELLOW MEALWORMS |
| | GRANARY WEEVILS | RED FLOUR BEETLES | |

E.P.A. Reg. No. 655-684

E.P.A. Est. No. 655-GA-1

NET CONTENTS

DIRECTIONS FOR USE (Cont'd)

... (text partially obscured) ...

LOT #

... (text partially obscured) ...

SURFACE SPRAY: To insure maximum control of crawling insects (ANT, BOXELDER BUGS, CLOVER MITES, DRUGSTORE BEETLES, EARWIGS, LICE, SPIDER BEETLES) dilute 1 part concentrate plus 29 parts of refined kerosene and mix well and apply as a coarse wetting spray using approximately 1 gallon of spray mixture to 750 square feet of surface. Spray walls, floor and other surfaces in bins and storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Treat unloading and processing areas and inside conveying processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible. For in transit protection, clean and spray truck beds, boxcars and ship's hold before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers slightly as stacked. To control INDIAN MEAL MOTHS and ANGOUMOIS GRAIN MOTHS in stored grain, apply to the surface of binned grain at a rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of moths and continuing until early April. In case of severe infestation, break up a bedding with rake before spraying and make a second application in two weeks. As an aid in control of MOSQUITOES, GNATS, WASPS AND SKIPPERS, treat walls, ceilings, moldings, screens, door and window frames, beams, light cords and similar resting places. **ANIMAL QUARTER USE:** (Cattle barns, horse barns, poultry houses, swine houses, zoos): As a space spray diluted for use in conventional mechanical fogging equipment to kill FLEES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS dilute 1 part concentrate plus 29 parts of refined kerosene and mix well. Apply at the rate of 2 fluid ounces per 1,000 cubic feet of space above the animals. Direct spray into the upper portions of the enclosure hitting the room with mist fog. Repeat treatment when re-infestation occurs. **INDOOR PRODUCTION AND PROCESSING:** To control MUSHROOM FLIES or FUNGUS GNATS in growing houses, dilute 1 part of concentrate plus 29 parts Methylene Chloride and mix well and apply at a rate of 1 ounce per 1,000 cubic feet of space. Sprayer should be operated 8 to 10 feet away from beds and should be directed slightly upward over them. Circulating fans should be left in operation but for best results, air circulation should be turned off in packaging and canning plants. Use at a rate of one half ounce per 1,000 cubic feet of space. **FOOD USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS:** Use and follow with label spraying device at the rate of 1 ounce per 1,000 cubic feet of room space. Apply the spray into all cracks and crevices in woodwork, walls, floors, underneath pipes, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Repeat as necessary for control of exposed or inaccessible stages of ROACHES, SILVERFLY, WATERBUGS, EARWIGS, ANTS, SPIDERS, FLEES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS AND GNATS. Before using, exposed food products and packaging material must be covered or removed. After use, equipment and utensils must be thoroughly rinsed with potable water before operation are resumed. Use as a space spray only when plant is in operation. **OUTDOOR GROUND APPLICATION (UNDILUTED):** To control ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS. For best results, apply when there is a light breeze of approximately 5 mph and temperature is 60° to 75° F. With a fogging device spray at a dosage rate of 5 ounces per minute, upwind of the area to be treated. If the equipment is truck mounted, keep the speed to approximately 5 mph with a swath width of no more than 300 feet. To control MOSQUITO LARVAE dilute at the rate of 15 ounces per 50 gallons of water and apply at the rate of 20 to 25 gallons per acre. Apply as a coarse wetting spray to the border or on shallow water of wet lands, swamps and marshes where mosquito larvae are known to breed. **TEMPORARY REDUCTION OF ANNOYANCE FROM FLEES, MOSQUITOES AND SMALL FLYING MOTHS OUTDOORS:** The air mentioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in public gathering places, drive in restaurants, drive in theaters, golf courses, municipal parks, playgrounds and recreational areas. Direct application into tall grass, shrubbery and around trees where these pests may travel or rest, apply while birds sing. Avoid wetting foliage. Application should be made prior to attendance. Repeat as necessary. In additional outdoor areas, sprays feed lots, stables and zoos, cover water, drinking fountains and animal feed before use. Fill area with mist directing application into tall grass, shrubbery and around trees where these pests may hover or rest, apply while birds sing. Avoid wetting foliage. In zoos, avoid exposure of reptiles to the product. Repeat as necessary. **FOOD USE ON ANIMALS:** Top dress beef and dairy cattle, horse and dogs from ticks, flees, HOUSE FLIES, MOSQUITOES AND GNATS dilute 1 part of concentrate plus 29 parts of refined kerosene and mix well. Apply at night, mist sufficient to wet the top of the hair. For control of STABLE FLIES, HORSE FLIES and DEER FLIES in beef and dairy cattle and horses, apply 2 ounces per adult animal, sufficient to wet the hair but not the face. Repeat treatment once or twice daily or at intervals of several times per week. To control the immature stages of BLACK FLIES, NG LICE on beef and dairy cattle, apply the spray thoroughly with the hair use 2 ounces of less per animal and repeat 2 to 3 weeks as required.

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE



Manufactured by
prentiss
Drug Chemical Co., Inc.

Plant: Kroll Road
Sandersville, GA. 31082
Office: 363 7th Avenue
Atlanta, GA. 30308

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

- if swallowed** - Call a physician or Poison Control Center immediately. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia.
- if inhaled** - Remove victim to fresh air. Apply artificial respiration if indicated.
- if on skin** - Remove contaminated clothing and wash affected areas with soap and water.
- if in eyes** - Flush eyes with plenty of water. Call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PEST DOCUMENT AVAILABLE

LEFT PANEL

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist. Avoid skin and eye contact. Wash thoroughly after handling and before eating or smoking.

Do not allow children or pets to contact treated surfaces until spray has dried. Do not use on plants. Do not allow spray to contact food, foodstuffs, or water supplies. Thoroughly wash dishes and food handling utensils contaminated with this product. Remove pets and cover fish bowls before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams and ponds. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use, or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

.....
.....
.....
.....
.....
.....