Prentox Pyronyl Livestock & Dairy Spray EPA Registration No. 655-694



PYRONYL LIVESTOCK & DAIRY SPRAY

For the control of Houseflies, Stable Flies, Horse Flies, Horn Flies, Mosquitoes, Gnats and Lice.

Can be used on Dairy & Meat Animals and in Dairy Barns, Milk Rooms and Dairies.

ACTIVE INGREDIENTS:	
Pyrethrins	. 0.10%
Piperonyl Butoxide Technical*	1.00%
INERT OTHER INGREDIENTS**:	
TOTAL	100.00%

PRENTOX® — Registered Trademark of Prentiss Incorporated PYRONYL™ — Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN			
CAUTION FIRST AID			
	• Immediately call a Poison Control Center, physician, or the National Pesticide		
if swallowed	Telecommunications Network at 1-800-858-7378.		
	Do not induce vomiting unless told to do so by the Poison Control Center or physician.		
	Do not give any liquid to the person.		
	Do not give anything by mouth to an unconscious or convulsing person.		
	Take off contaminated clothing.		
If on skin or • Rinse skin immediately with plenty of water for 15-20 minutes.			
clothing	Call a Poison Control Center, physician, or the National Pesticide Telecommunications		
	Network at 1-800-858-7378 for treatment advice		
	Hold eye open and rinse slowly and gently with water for 15-20 minutes.		
If in eyes	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.		
	Call a Poison Control Center, physician, or the National Pesticide Telecommunications		
	Network at 1-800-858-7378 for treatment advice		
	Move person to fresh air.		
lf inhaled	If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably		
	mouth-to-mouth, if possible.		
	Call a Poison Control Center, physician, or the National Pesticide Telecommunications		
	Network at 1-800-858-7378 for treatment advice.		
NOTE TO PI	IYSICIAN: Contains petroleum distillate – vomiting may cause aspiration pneumonia		
For informatio	n on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call		

E.P.A. REG. NO. 655-694	E.P.A. EST. NO. 655-GA-1
NET CONTENTS	LOT NO
Manufactured by:	

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

PRENTISS INCORPORATED

ACCEPTED

MAR 2 6 2002

the National Pesticide Telecommunications Network at 1-800-858-7378.

Under the Perieral Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under RPA Beg. No. 655-694 Plant: Kaolin Road, Sandersville, GA 31082 Office: C.B. 2000, Floral Park, NY 11002-2000

Page 1 of 2

^{*}Equivalent to 0.8% (Butylcarbityl) (6-propylpiperonyl) Ether and 0.2% related compounds.

^{**}Contains Petroleum Distillates

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist. Avoid skin and eye contact. Wash thoroughly after handling and before eating or smoking.

Do not allow children or pets to contact treated surfaces until spray has dried. Do not use on plants. Do not allow spray to contact food, feedstuffs or water supplies. Thoroughly was dishes and food handling utensils contaminated with this product. Remove pets and cover fish bowls before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use, or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ON ANIMALS: To protect animals from horn flies, house flies, mosquitoes and gnats, apply as a light mist sufficient to wet the tips of the hair. To control stable flies, horse flies, deer flies, apply 2 ounces per adult animal, sufficient to wet the hair but not to soak the hide. Repeat treatment once or twice daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply the spray lightly to infested areas, using a stiff brush to get the spray to the base of the hair. Use 2 ounces or less per animal. Repeat every two or three weeks as required.

BARNS, MILK ROOM, DAIRY PLANTS: To control flies, mosquitoes, and gnats, apply as a fog or fine mist, directing spray toward the ceiling and upper corners, until the area is filled with mist, using about 2 ounces per 1000 cubic feet of space. For best results, close windows and doors before spraying and keep closed for ten to fifteen minutes. Vacate the area and ventilate before reoccupying. Repeat application as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaccous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Do not store, use, pour or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

2/94