

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get on face, neck, or clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets. Do not allow children to handle areas until contents are dry. DO NOT breathe vapors. Do not use this product on animals simultaneously or within a few days before or after treatment with any agent which interferes with the drug, pesticides or chemicals.

Do not use on fish or aquatic animals, or they show hypersensitivity to organo-phosphates.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on soaked seeds may be killed. Do not apply directly to fish, streams and ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

PERSONAL HAZARDS

Do not use or spray near food or open flame.

USE PRECAUTIONS

IF SWALLOWED: If breathing difficulty occurs, apply artificial respiration. Atropine sulfate and 2-PAM are antidotes.

IF INHALED: In case of acute symptoms (irritative reaction, respiratory difficulty, muscle twitching or convulsions, excitation, headache, blurred vision, contracted pupils) treat with the antidote listed as above.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticides and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or straw. Sweep the contaminated absorbent on to a shovel and put the sweepings into a storage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour, or spill near food or open flame.

POSTUSE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

prentox[®]

FOGGER OIL AND DAIKY SPRAY

For Agricultural/Commercial Use Only.
For Killing Houseflies, Horn Flies, Stable Flies, Face Flies, Mosquitoes, Gnats, Spiders and certain Other Insects in Dairy Barns and Other Farm Buildings. Also for direct application to Dairy and Beef Cattle.

ACTIVE INGREDIENTS:

2, 2-Dichloroethyl Dimethyl Phosphate	8.50%
Related Compounds	8.67%
Petroleum Distillate	82.83%
TOTAL:		100.00%

REGISTERED TRADEMARK OF PRENTISS DRUG & CHEMICAL CO.

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. **DO NOT INDUCE VOMITING** unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply respiration if indicated.

If on skin: Wash contaminated skin with soap and water.

If in eyes: Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

SEE SIDE PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. REG. NO. 655-782 E.P.A. EST. NO. 655-GA-1

NET CONTENTS _____

LOT # _____

MODE OF ACTION

Prentox is a contact insecticide which acts by contact with the body of the insect. It is absorbed into the body and acts systemically. It is effective against houseflies, horn flies, stable flies, face flies, mosquitoes, gnats, spiders and certain other insects.

Prentox is a contact insecticide which acts by contact with the body of the insect. It is absorbed into the body and acts systemically. It is effective against houseflies, horn flies, stable flies, face flies, mosquitoes, gnats, spiders and certain other insects.

Prentox is a contact insecticide which acts by contact with the body of the insect. It is absorbed into the body and acts systemically. It is effective against houseflies, horn flies, stable flies, face flies, mosquitoes, gnats, spiders and certain other insects.

Fogging or Space Spraying: In all flies, mosquitoes, gnats and spiders in dairy and beef cattle barns, apply daily as a fog or mist (about 1 pint per 1,000 cubic feet). One application is sufficient. For mosquitoes, gnats and spiders, apply twice weekly. For flies, apply twice weekly. Do not apply in areas where animals are present. Do not apply in areas where children are present.

HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS: Prentox is a contact insecticide which acts by contact with the body of the insect. It is absorbed into the body and acts systemically. It is effective against houseflies, horn flies, stable flies, face flies, mosquitoes, gnats, spiders and certain other insects.

Dairy Barns and Other Buildings: Dilute one part Fogger Oil and Dalky Spray with one part of fuel oil, diesel or kerosene. Apply this solution as a mist or spray at the rate of a quart per 1,000 square feet to control flies, gnats, mosquitoes, roaches, spiders, fleas, ants, sow bugs, brown dog ticks.

For milk sheds, stables, livestock barns, feeding sheds, pig pens and outdoor areas, treat around doorways, feed storage rooms, alleyways and window sills.

In poultry houses, apply to manure, window sills, outside of penned enclosures and on floor of feed storage rooms.

In treating dog kennels apply to outside dog runways, window sills and ledges. For control of fleas and brown dog ticks increase dosage to 2 quarts per 1,000 square feet.

For control of fly maggots in maggot breeding areas such as garbage dumps and manure piles apply at the rate of 2 quarts per 100 square feet.

ACCEPTED
MAR 27 1984
U.S. DEPARTMENT OF AGRICULTURE
EPA REG. NO. 655-782



Manufactured by
prentiss
Drug & Chemical Co., Inc.

Plant: Kaolin Road
Sandersville, GA. 31082
Office: C.B. 2000
Floral Park, N.Y. 11001