

pm

03

655-702

10/22/97

page 185

OCT 22 1997

Mr. Richard Miller
Prentiss Incorporated
C.B. 2000
Floral Park, NY 11002-2000

Dear Mr. Miller:

Subject: Label Amendment in Response to Basic Suppliers Deletion of Certain DDVP Uses
Prentox Fogger Oil and Dairy Spray
EPA Registration No. 655-702
Your Submission Dated May 8, 1995

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes indicated below prior to release for shipment of product bearing the amended labeling:

1. Under the heading Direct Application add the following: "Do not make direct application to livestock more frequently than once per day."
2. Directly following the statement "For control of fleas and brown dog....." add: "Do not make direct applications to dogs."
3. Revise the Active Ingredient statement as follows: "Dichlorovos [2,2-dichlorovinyl dimethyl phosphate....."
4. Under precautionary statements add a paragraph stating appropriate personal protective equipment, i.e.: "Required Protective Clothing: Persons who handle this product must wear the following protective clothing: Coveralls worn over short sleeved shirt and pants, socks, chemical resistant footwear, chemical resistant (nitrile or butyl) gloves and protective eyewear. A respirator with either or organic vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G) when applying indoors as a space spray of fog."
5. Replace Note to Physician on second page with the Note to Physician on the third page.

- 6. Under Fogging or Space Spraying replace Keep cattle in barn when spraying with:
"Close doors and windows before treatment. Do not remain in treated areas and ventilate the area after treatment is complete. Remove animals prior to treatment. Wear appropriate personal protective equipment for space spraying as indicated under precautionary statements."
- 7. Revise the statement to read: "Do not contaminate water by disposal of equipment washwaters."

Submit two copies of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide Branch
Registration Division (7505C)

3 of 5

Prentox Fogger Oil and Dairy Spray
EPA Reg. No. 655-702

PRENTOX® FOGGER OIL AND DAIRY SPRAY

~~FOR AGRICULTURAL/COMMERCIAL USE ONLY.~~

FOR AGRICULTURAL/COMMERCIAL USE ONLY. NOT FOR SALE FOR ANY OTHER USES.

For Killing Houseflies, Horn Flies, Stable Flies, Face Flies, Mosquitoes, Gnats, Spiders and certain Other Insects in Dairy Barns and Other Farm Buildings. Also for direct application to Dairy and Beef Cattle.

ACTIVE INGREDIENTS:

2,2-Dichlorovinyl Dimethyl Phosphate	0.93%
Related Compounds	0.07%

INERT INGREDIENTS*:	99.00%
TOTAL:	100.00%

*Contains Petroleum Distillates
PRENTOX® - Registered Trademark of Prentiss Incorporated

KEEP OUT OF REACH OF CHILDREN
WARNING
STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. **DO NOT INDUCE VOMITING** unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply respiration if indicated.

If on skin: Wash contaminated skin with soap and water.

If in eyes: Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. REG. NO. 655-702

E.P.A. EST. NO. 655-GA-1

NET CONTENTS _____

LOT # _____

Manufactured by:

PRENTISS INCORPORATED

Plant: Kaolin Road, Sandersville, GA 31082
Office: C.B. 2000, Floral Park, NY 11001

ACCEPTED
with COMMENTS
in EPA Letter Dated

OCT 22 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

Page - 1

655-702

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets. Do not allow children in treated areas until surfaces are dry. DDVP is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticide or chemicals.

Do not use on Braham or Braham cross-cattle, as they show Hypersensitivity to organic phosphates.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to lakes, streams and ponds. Do not apply when weather conditions favor drift from the areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use or store near heat or open flame.

NOTE TO PHYSICIAN

If Swallowed: If breathing difficulty occurs, apply artificial respiration. Atropine Sulfate and 2-PAM are antidotal.

If Inhaled: In case of severe symptoms (excessive salivation, respiratory difficulty, muscle twitches or convulsions, exhaustion, headache, blurred vision, constricted pupils)(acetylcholine effects) treat as above.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour, or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

5 7 5
Prentox Fogger Oil and Dairy Spray
EPA Reg. No. 655-702

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, Atropine Sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams, until complete Atropinization has occurred. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

Morphine is contraindicated. Clear chest by postural drainage. Observe patient continuously for 48 hours.

Repeated exposure to cholinesterase inhibitors may without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

For use only in areas under the general category of farm buildings (farm yards); manure treatments on farm premises; dairy and farm premises; feed lots, stockyards, corrals, holding pens, fences, etc.; animal buildings, including barns, feeding areas, shelters and stables; dairy barns (including milk rooms, equipment and barnyards); livestock feeding areas; pens, poultry droppings, poultry houses (equipment and yards); beef cattle, poultry, dairy cattle, goats, horses (including ponies), sheep, swine and turkeys.

Fogging or Space Spraying: To kill Flies, Mosquitoes, Gnats and Spiders in dairy and beef cattle barns, apply daily as a fog or mist inside building at rate of 1 pint per 8,000 cubic feet. Keep cattle in barn when spraying. Let spray settle before milking. Do not apply in areas where animals have received direct application of this product within 8 hours.

Direct Application: To protect dairy and beef cattle from Face Flies, Houseflies, Stable Flies, Horn Flies, Mosquitoes and Gnats, spray animal daily using 1 to 2 fluid ounces per animal. Apply as a mist spray so as to wet the hair. Do not soak the skin or wet the hide. Pay particular attention to the face, back, flank and legs. Experience indicates that two applications of 1 fluid ounce per animal morning and night give outstanding control of the above pests. Do not apply in excess of 2 fluid ounces per animal per day.

DAIRY BARN AND OTHER BUILDINGS

Dilute one part Fogger Oil and Dairy Spray with one part of fuel oil, diesel or kerosenes. Apply this solution as a coarse wet spray at the rate of a quart per 1,000 square feet to control Flies, Gnats, Mosquitoes, Roaches, Spiders, Fleas, Ants, Sowbugs, Brown Dog Ticks.

For milk sheds, stables, livestock barns, loafing sheds, pig pens and outdoor areas, treat around doorways, feed storage rooms, alleyways and window sills.

In poultry houses, apply to manure, window sills, outside of penned enclosures and on floor of feed storage rooms.

In treating dog kennels apply to outside dog runways, window sills and ledges. For control of fleas and brown dog ticks increase dosage to 2 quarts per 1,000 square feet.

For control of fly maggots in maggot breeding areas such as garbage dumps and manure piles apply at the rate of 2 quarts per 100 square feet.