



prentox METHOXYCHLOR 50W

WETTABLE POWDER FOR DAIRY AND BEEF CATTLE BUILDINGS AND CROPS

Wettable Powder Insecticide for the control of various insects on Livestock, Vegetables, Fruits, Grasses and Legumes, Forest & Shade Trees, Flowers and Ornamentals, Agricultural Premises, Farm Buildings and Grain Storage Bins

ACTIVE INGREDIENT:		
*Methoxychlor Technical	50%
INERT INGREDIENTS:	50%
Total	100%

*50% METHOXYCHLOR TECHNICAL is equivalent to 44% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 6% other isomers and reaction products

KEEP OUT OF REACH OF CHILDREN

CAUTION:

E.P.A. Reg. No. 855-741

E.P.A. Est. No. 855-GA-1

BEST DOCUMENT AVAILABLE

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(& DOMESTIC ANIMALS)

CAUTION: May be harmful if swallowed. Avoid contact with skin. Wash thoroughly after handling. Avoid breathing dust or spray mist. Avoid contamination of feed, seed and foodstuffs.

Environmental Hazards
This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.
This product is toxic to bees and should not be applied when bees are actively visiting the area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

General Information
Methoxychlor 50W insecticide contains 50% technical methoxychlor per pound and may be used as a spray-water suspension for plant and animal protection or may be diluted with suitable diluents (alk. pyrophosphate) to make a finished dusting compound to use on livestock (see directions for specific uses). When applying Methoxychlor 50W use sufficient spray or dust to thoroughly cover all foliage, fruit or other surfaces to be protected.

DIRECTIONS
READ THE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

See additional directions for use on back panel

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

Pesticide Disposal: Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

AGRICULTURAL PREMISES
FARM BUILDINGS: For controlling stable flies, houseflies and fleas use 1 lb. in 2 gallons of water. Apply 1 gallon of spray mixture to approximately 500 sq. ft. of surface treating walls, floors, cracks, crevices, baseboards, pet bedding stalls and pens. Such heavy deposits usually are effective to 3 to 6 weeks unless removed by weathering or other means. Care should be taken to avoid contamination of milk handling equipment, feedstuffs, feed troughs and watering receptacles. Dairy animals should not be present while spraying. The use of any residual fly spray should be supplemented with proper sanitary measures including spraying or removal of breeding sources such as manure in and around buildings. Do not apply in poultry houses.

GRAIN STORAGE BINS: To clean up infestations in empty storage bins use 1 lb. Methoxychlor 50W in 2 gallons of water applied at the rate of 1 gallon of spray per 500

sq. ft. before products are stored. This aids in control of adult rat grain beetle, sawtoothed grain beetle, lesser grain borer, adult flour beetle, long headed flour beetle, rice weevil, confused flour beetle, brown grain beetle, hairy fungus beetle and grain moth. Allow bins to air and dry thoroughly before filling with grain 2 to 4 weeks.

Elevator Tunnels, Gallery Floors, Head Houses: To control weevils, confused flour beetle, flat headed grain beetle, long headed flour beetle, granary weevil, hairy fungus beetle, lesser grain borer, long headed flour beetle, rice weevil, sawtoothed grain beetle and grain moth mix 1 lb. in 2 gallons of water. Apply 1 gallon mixture to 500 sq. ft. clean area thoroughly before spraying. Treat about 3 times per season.

AGRICULTURAL CROPS
Vegetables: For control of the insects listed below use 3 lb. of Methoxychlor 50W per 100 gals. of water and apply as a thorough spray to upper and lower surfaces of foliage and fruit at the rate of 100 gals. per acre. Begin treatment at first sign of insect infestation and repeat at 7-10 day intervals as infestation warrants.

Crop	Insects	Interval From Last Application To Harvest
Potato	Blister beetle, Colorado potato beetle, Flea beetle, Leafhopper	9 days
Cantaloupe	Melon worm	1 day
Cucumber	Cucumber beetle	
Eggplant	Flea beetle, Leafhopper	
Kohlrabi	Flea beetle	
Pepper	Flea beetle, Rose chaffer	
Pumpkin	Cucumber beetle, Squash vine borer	
Squash	Cucumber beetle, Squash vine borer	
Tomato	Blister beetle, Fall army worm, Flea beetle, Japanese beetle, Tomato fruit worm	
Asparagus	Asparagus beetle	If applied later than 3 days before harvest, remove residues by washing or blanching
Beans	Bean leaf beetle, Corn earworm, Fall army worm, Japanese beetle, Leafhopper, Rose chaffer, Mexican bean beetle	3 days. Do not use again for feed or forage
Black-eyed peas	Compebrutid	
Cabbage	Cabbage worm, Flea beetle	3 days
Cauliflower	Cabbage worm, Flea beetle	3 days
Corn	Fall army worm, Flea beetle, Japanese beetle, Rose chaffer	
Melons	Cucumber beetle, Melon worm	
Peas	Rice weevil	
Radish & Rutabaga	Flea beetle	
Beets	Blister beetle, Flea beetle	14 days
Broccoli	Cabbage worm, Flea beetle	
Brussels sprouts	Cabbage worm, Flea beetle	
Carrot	Leafhopper	
Collards & Kale	Flea beetle	
Lettuce	Leafhopper	
Spruce & Turnip	Flea beetle	

BEST DOCUMENT AVAILABLE



Manufactured by

prentiss

Plant: 338 Wilson Avenue
Newark, N.J. 07105
Office: 267 7th Avenue

DOT 6

Prentiss Drug and Chemical Co., Inc.
363 Seventh Avenue
New York, NY 10001

Gentlemen:

Subject: Prentox^(R) Methoxychlor 50 W
EPA Registration No. 655-741
Application Date: August 17, 1982

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

TS-767:Ellenberger:DCR-24758;WANG-0497C:jad:Raven:479-2013:10/04/82