

AUG 8 1985

Connie Yeisley
Prentiss Drug & Chemical Co., Inc.
C.B. 2000
Floral Park, NY 11001

301/153697
17/5

Dear Mrs. Yeisley:

Subject: Amendment-Label Update
Prentox 20 Lindane Emulsifiable Concentrate
EPA Registration No. 655-768
Your Application Dated June 11, 1985

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable. A stamped copy is enclosed for your records.

Add to the next printing of your label: Note to Physician: Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally. If symptoms of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine.

You may delete the reentry restrictions. These are not necessary for a homeowner use.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TR-767)

Enclosure

OPTS/RD:JOE:85843:LaRocca:RD-58:Kendrick & Co:898-1270:8/5/85:Del.8/15/85:eg

CONCURRENCES

SYMBOL ▶								
SURNAME ▶	Nestorik							
DATE ▶	8-8-85							

prentox[®]

20 LINDANE EMULSIFIABLE CONCENTRATE

Applicators must wear the following protective clothing during the application process: Long-sleeved shirt, long pants, waterproof gloves, full foot covering and a head covering.

ACTIVE INGREDIENTS

Lindane (Gamma isomer of Benzene Hexachloride)	20.0%
Xylene Range Aromatic Solvent	62.2%
INERT INGREDIENTS:	17.8%

(Contains 1.63 pounds of Lindane per gallon)

TOTAL: 100.0%

PRENTOX[®]—Registered Trademark of Prentiss Drug & Chemical Co., Inc.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician or Poison Control Center immediately. CONTAINS PETROLEUM DISTILLATES. DO NOT INDUCE VOMITING. Get medical attention immediately. If inhaled — Remove person to fresh air. If on skin — Wash area immediately with soap and warm water. If in eyes — Flush eyes with plenty of water and get prompt medical attention.

**SEE PANEL BELOW
FOR ADDITIONAL PRECAUTIONARY STATEMENTS.**

E.P.A. REG. NO. 665-768

E.P.A. EST. NO. 665-GA-1

NET CONTENTS _____

LOT NO. _____

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be absorbed through skin. Harmful if swallowed. Avoid inhalation of spray mist. Avoid skin contact. In case of contact, wash immediately with soap and water. Wash thoroughly after using and before eating or smoking. Avoid contamination of feed and food-stuffs. Do not use on household pets and humans. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment or disposal of wastes.

BENEFICIAL INSECT CAUTION

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.



Manufactured by

prentiss

Drug & Chemical Co., Inc.

Plant: Keolin Road
Sandersville, GA. 31062
Office: C.B. 2000
Floral Park, N.Y. 11001

BULBS (DAFFODIL, GLADIOLUS, HYACINTH, IREX, LILIES, NARCISSUS, TULIPS): Aphids, Bulb fly, European corn borer, Iris borer, Thrips. Apply when insects appear. Repeat as necessary.

FLOWERS (CARNATIONS, CHRYSANTHEMUMS, COLEUS, DAISSY, GARDENIA, GERANIUMS, Hibiscus, MASTUR-TURN, PANSIES, PETUNIAS, SWEET PEAS, ZIBIBAS): Aphids, blister beetles, Leafhoppers, Thrips. Apply when insects appear. Repeat as necessary.

ORNAMENTALS (AZALEAS, CAMELLIAS, HEATHER, LILACS, RHODODENDRONS, ROSES): Aphids, Blackvine weevil, Cucumber beetles, Thrips, White Fly. Apply when insects appear. Repeat as necessary.

TREES OR SHRUBS	BORER AND OTHER INSECTS	DOSE	TIME TO APPLY
Elm (English and Scotch) Birch, Lacus	Leaf Miners	1 teaspoonful 1 gal water	Apply when small miners appear in early and mid-May. Apply when small miners appear in early July.
Pines	Pine Bark Beetles	1 pint/5 gal water	Spray trunks thoroughly. Wet trunks to run-off.
Pine, Spruce	White Pine Weevil	1 Tablespoonful/gal water	Apply late April when buds begin to swell. Spray upper parts of trees thoroughly.
Taxus, Hemlock	Strawberry Root Weevil Taxus Weevil	1 Tablespoonful/1 gal water	Apply in May or early June to the trunk and around base of trees, soaking the bark and cutting from surface.
Dogwood, Rhododendron, Lilac	Dogwood Borer, Rhododendron Borer, Lilac Borer	1 Tablespoonful/1 gal water	Make 2 to 3 applications 7 to 10 days apart in May and early June.
Apples, Cherries, Peaches	Fruit Tree Bark Beetles, Flat-Headed Borers, Round-Headed Borers	1 Tablespoonful/1 gal water	Apply to bark of tree trunk and large limbs in spring. Apply to trunk, crotch to ground. Soak bark to run-off. Do not apply to fruit or foliage.*
Maple, Aspen, Poplar, Ash, Willows, Elm, Locust	Aphids, Bark Beetles, Wood Boring Beetles	2 Tablespoonful/1 gal water	Apply in spring when buds begin to break. Repeat in 10-14 days. Elms — Apply to foliage when Aphids are present during summer, repeat in fall. Soak trunks to run-off.
Peaches	Peach Tree Borer	1 Tablespoonful/gal water	Spray lower trunk and around base of tree, soaking bark and cutting trunk from crotch to ground. Do not apply to fruit or foliage.*

*DO NOT APPLY WITHIN 60 DAYS OF HARVEST


ACCEPTED
FOR COMMENTS
Re EPA Letter Dated:

AUG 8 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for products registered under No.

6255-768

12/84



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand, or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Do not store, use, pour, or spill near heat or open flame.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

DIRECTIONS FOR USE (Cont'd.)

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops being treated with this product, consult your State Department of Agriculture for further information.

DO NOT ALLOW CHILDREN TO HANDLE OR APPLY THIS PRODUCT. CHILDREN AND PETS SHOULD NOT BE ALLOWED IN TREATED AREAS UNTIL SPRAYED SURFACES ARE DRY.

FRUIT TREES, VEGETABLES, FLOWERS, AND ORNAMENTALS: Apply 0.03% Lindane solution as directed below. Prepare 0.03% solution by adding 1 tablespoonful Lindane Concentrate to 2½ gallons of water or ½ pint to 40 gallons of water.

PEACHES, PLUMS: Aphids, Leafrollers, Thrips: Use 1 gallon of 0.03% spray per 100 square feet. Do not apply within 60 days of harvest.

TOMATOES: Aphids: Use 1 gallon of 0.03% spray per 150 square feet. Do not apply after fruit starts to form.

CABBAGE, CELERY, COLLARDS, CUCUMBERS, LETTUCE, OKRA, PEPPERS, SQUASH, MELONS, TOMATOES: White Grubs, Cutworms, Wireworms: For preplant soil application only. Use 2 gallons for each 250 square feet. Work into soil thoroughly before or during planting. Do not treat land to be planted in root crops.