

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Timothy A. Gardner  
Product Manager (17)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767)

Enclosure

JA

# PRENTOX<sup>(R)</sup> INSECT SPRAY & FOGGING CONCENTRATE

For the control of flies, mosquitoes, small flying moths, gnats, wasps, roaches, waterbugs, silverfish, crickets, carpet beetles, bedbugs, ants, fleas, house flies, stable flies, horn flies, and exposed stages of stored product pests.

### ACTIVE INGREDIENTS:

Pyrethrins.....	0.30%
Piperonyl Butoxide, Technical* .....	0.60%
N-octyl bicycloheptene dicarboximide ...	1.00%
Petroleum Distillates.....	98.10%
TOTAL:	100.00%

\*Equivalent to 0.48% (Butylcarbityl) (6-propylpiperonyl) ether and 0.12% related compounds.

PRENTOX(R) - Registered Trademark of Prentiss Drug & Chemical Co., Inc.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**If swallowed** Call a physician or Poison Control Center immediately. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia.

**If inhaled** Remove person to fresh air.

**If on skin** Remove contaminated clothing and wash affected areas with soap and water.

**If in eyes** Flush eyes with plenty of water. Call a physician immediately if irritation persists.

**SEE SIDE PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS**

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 16 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act  
as amended, the pesticide  
registered under EPA Reg. No.  
655-173

E.P.A. Reg. No. \_\_\_\_\_ E.P.A. EST. NO. 655-GA-1


NET CONTENTS \_\_\_\_\_

LOT # \_\_\_\_\_



Manufactured by  
**prentiss**  
Drug & Chemical Co., Inc.

Plant: Kaolin Road  
Sandersville, GA. 31082  
Office: C.B. 2000  
Floral Park, N.Y. 11001



LEFT PANEL

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. If swallowed, do not induce vomiting. Call a physician immediately.

Do not apply directly to milk. Cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying in barns, poultry houses, warehouses, etc., wash with an approved detergent then rinse where exposed food will be handled to remove all traces of contamination. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

#### ENVIRONMENTAL HAZARDS

The product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water by disposal of waste. Apply this product only as specified on this label.

#### PHYSICAL HAZARDS

Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping prohibited.

**STORAGE:** If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand, or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Do not store, use, pour, or spill near heat or open flame.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

## DIRECTIONS FOR USE (Cont'd)

Clean up and destroy fallen insects. Spray when plant is not in operation. Repeat as necessary.

Cover or remove exposed food and close room. Spray outside walls of grain bins, flooring, about base of milling machinery, walls and baseboards, areas behind cabinets, around pipes and other insect harbourages using 1 gal. of spray per 1,000 sq. ft. of surface. Force spray into all cracks and crevices. Repeat applications as necessary. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

SPACE SPRAY - Undiluted in Aerosol Generators - To kill flies, mosquitoes, small flying moths and gnats, cockroaches, and accessible stages of granary insects: rice & granary weevil, red flour beetle, Indian meal moth larvae, cadelles, grain mites and saw-toothed grain beetles, Angoumois grain moth, chocolate moth, carpet beetle, cigarette beetle, flour grain beetle. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating equipment. Apply at the rate of 4 to 12 oz. of undiluted concentrate per 1,000 cu. ft. of space. Direct output from aerosol generator upward towards upper corners of room and ceiling. Leave room closed for one half hour. Do not remain in treated areas. Ventilate the area after treatment is completed. Sweep up and destroy fallen insects. Repeat as necessary.

SPACE SPRAY - Use in Barns & Poultry Houses, Warehouses, Storage Bins & Storage Rooms - To control house flies, stable flies, and horn flies, apply as a mist over the livestock in stanchions or holding pens or over the poultry in cages or in pens. Use a fogger or mist generator, so spray is directed and applied above the animals and direct spray to center of the area where animals are housed. Apply 2 oz. per 1,000 cu. ft. so that the spray mist is distributed uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in dairy barns, aerate the barn thoroughly before milking. Repeat daily, morning and afternoon, or as required. Keep sprayer unit reservoir and nozzle clean, to ensure proper delivery of spray mist or fog. Product is to be used in mechanical sprayers, which give particle size aerosol range or smaller, such as ULV type.

SPACE SPRAY - For Use in Mechanical Fogging Equipment - Kills flies, mosquitoes, small flying moths & gnats. Cover or remove exposed food and food handling surfaces. Close room, making sure to shut off all air conditioning or ventilation equipment. Dilute with deodorized kerosene at a rate of 1 part concentrate with 4 parts kerosene. Apply at the rate of 50 fl. oz. of spray per 25,000 cu. ft. of space. Make sure to direct the spray upward towards upper corners of room & ceiling. Keep room closed for one half hour after spraying. Do not remain in treated area. Ventilate after treatment is completed.

FOODS SHOULD BE REMOVED OR COVERED DURING TREATMENT. ALL FOOD PROCESSING SURFACES SHOULD BE COVERED DURING TREATMENT OR THOROUGHLY CLEANED BEFORE USING. WHEN USING THIS PRODUCT IN THESE AREAS, APPLY ONLY WHEN THE FACILITY IS NOT IN OPERATION.

#### DIRECTIONS FOR USE (CONTD)

For best results, use a good clean sprayer — free of rust — which produces a fine mist.

**FOR FLYING INSECTS:** Flies, mosquitoes, small flying moths, gnats and wasps: Close doors and windows. Spray freely towards ceiling in all directions until room is fogged with mist. When spraying, keep about 3 feet from interior walls, fabrics and furniture. Doors and windows may be opened after 10 minutes.

**FOR CRAWLING INSECTS:** Roaches, waterbugs, silverfish and ants: Spray thoroughly into cracks and crevices, under and behind sinks, cupboards, drawers, woodwork, refrigerators, pipes and drains, where these insects hide. Spray directly on as many insects as possible. Repeat application as necessary.

**CONTACT SPRAY - FOR CLOTHES MOTHS & CARPET BEETLES:** Clean and spray closets thoroughly, to reach all hiding places. Clean and brush woolen articles. Repeat application every 30 days. When carpets are infested, spray both sides thoroughly. When spraying near fabrics, keep spray in constant motion, at least 36 inches away from fabric. Spraying closer may cause stains. When spraying clothes in storage for adult moths, separate clothing. For the larvae, pay particular attention to seams and folds of garments. Clothing should be sprayed individually. For storing clothing in chests, first treat as per above, and also spray storage container before packing. Repeated monthly applications are advisable.

**FOR BEDBUGS:** Spray baseboards, molding and floor boards. Spray mattresses, particularly around seams and tufts. Take beds apart and spray into all joints. Repeat application as necessary.

**FOR ANTS:** Spray trails, hills, around window frames and openings around pipes and baseboards, where ants crawl hitting as many as possible. Repeat application as necessary.

**FOR FLEAS:** Spray floors, floor coverings and sleeping quarters of pets and other infested areas. Repeat as required. Do not spray pets.

**FOR OUTDOOR USE:** To repel temporarily: Flies, mosquitoes, gnats, wasps, crickets, ants: On porches, terraces, lawns, and at outdoor gatherings, spray frequently around chairs, tables, floors, ground and surrounding air.

**CONTACT SPRAY - Use with Sprayers that give large, wetting spray Particles - To kill cockroaches, and accessible stages of granary insects: rice & granary weevil, red flour beetle, Indian meal moth larvae, cadelles, grain mites and saw-toothed grain beetles, Angoumois grain moth, chocolate moth, carpet beetle, cigarette beetle, flour grain beetle. Use 3-4 oz. per 1000 cu. ft. of space, directing the spray so as to contact the insects. Pay particular attention to cracks, crevices & other hiding places. Repeat treatment as required KILLS INSECTS ONLY WHEN THE SPRAY CONTACTS THEM. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.**

Sanitation is necessary. Clean and remove deposits of food, grain & dust from storage areas. Clean conveyer and handling equipment. For insects affecting stored food, as listed above, flour, grain & cereals - food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Apply by using equipment designed & adjusted to deliver a true aerosol mist. Close doors & windows and spray into air, directing spray up to all 4 corners, filling room with mist. Do not spray on processing equipment. Keep doors & windows closed 15 minutes after spraying.

US ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES PROGRAMS  
REGISTRATION DIVISION (75-767)  
WASHINGTON, DC 20460

EPA REGISTRATION NO.

655-774

DATE OF ISSUANCE

SEP 16 1985

TERM OF ISSUANCE

NOTICE OF PESTICIDE:  REGISTRATION  
 REREGISTRATION  
(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

Prentox Allethrin 0.3% Insect Spray  
and Fogging Insecticide

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Prentiss Drug & Chemical Co., Inc.  
CB 2000  
Floral Park, NY 11001

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
  - a. Add the phrase "EPA Registration No. 655-774."
  - b. Correct the percentage of allethrin declared on the label to read 0.3% rather than 0.03%.
  - c. For the Container Disposal statement, please add the statement for plastic containers and precede the two statements with the container type (e.g., metal, plastic).
  - d. Please provide the Product Chemistry for this product.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

9-16-85