

ACCEPTED

06 APR 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 655-773

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping prohibited.

STORAGE: If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand, or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Do not store, use, pour, or spill near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

PLASTIC CONTAINER DISPOSAL: Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by local authorities, by burning. If burned, stay out of smoke.



INSECT SPRAY AND FOGGING CONCENTRATE #1

For the control of flies, mosquitoes, small flying moths, gnats, wasps, roaches, waterbugs, silverfish, crickets, carpet beetles, bedbugs, ants, fleas, house flies, stable flies, horn flies, and exposed stages of stored product pests.

ACTIVE INGREDIENTS:	
Cyrethrins	0.30%
Piperonyl Butoxide, Technical*	0.60%
N-octyl bicycloheptene dicarboximide	1.00%
INERT INGREDIENTS:	98.10%
TOTAL:	100.0%

*Equivalent to 0.48% (Butylcebitol) (5-propylpiperonyl) ether and 0.12% related compounds.

PRENTOX™ — Registered Trademark of Prentiss Incorporated

** Contains Petroleum Distillates

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician or Poison Control Center immediately. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia. **If inhaled** — Remove person to fresh air. **If on skin** — Remove contaminated clothing and wash affected areas with soap and water. **If in eyes** — Flush eyes with plenty of water. Call a physician immediately, if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

E.P.A. REG. NO. 655-773

E.P.A. EST. NO. 655-GA-1

NET CONTENTS _____

LOT NO. _____

Manufactured by



Plant: Kaolin Rd., Sandersville, GA 31082

Office: C.B. 2000, Floral Park, NY 11001

655-773

04/06/1995

1/2

**PRECAUTIONARY STATEMENTS
HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. If swallowed, do not induce vomiting. Call a physician immediately.

Do not apply directly to milk. Do not apply while food processing is under way. Food should be removed or covered during treatments. All food processing surfaces should be covered during treatment or thoroughly cleaned before use. Remove pets and birds, and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

PHYSICAL HAZARDS

Do not use or store near heat or open flame. Exposure to temperatures above 130°F. may cause bursting.

DIRECTIONS FOR USE (Cont'd)

For best results, use a good clean sprayer — free of rust — which produces a fine mist.

FOR FLYING INSECTS: Flies, mosquitoes, small flying moths, gnats and wasps: Close doors and windows. Spray freely towards ceiling in all directions until room is fogged with mist. When spraying keep about 3 feet from interior walls, fabrics and furniture. Doors and windows may be opened after 10 minutes.

FOR CRAWLING INSECTS: Roaches, waterbugs, silverfish and crickets: Spray thoroughly into cracks and crevices, under and behind sinks, cupboards, drawers, woodwork, refrigerators, pipes and drains, where these insects hide. Spray directly on as many insects as possible. Repeat application as necessary.

CONTACT SPRAY — FOR CLOTHES MOTHS & CARPET BEETLES: Clean and spray closets thoroughly, to reach all hiding places. Clean and brush wooden articles. Repeat application every 30 days. When carpets are infested, spray both sides thoroughly. When spraying near fabrics, keep spray in constant motion, at least 36 inches away from fabric. Spraying closer may cause stains. When spraying clothes in storage for adult moths, separate clothing. For the larvae, pay particular attention to seams and folds of garments. Clothing should be sprayed individually. For storing clothing in chests, first treat as per above, and also spray storage container before packing. Repeat monthly applications are advisable.

FOR BEDBUGS: Spray baseboards, molding and floor boards. Spray mattresses, particularly around seams and tufts. Take beds apart and spray into all joints. Repeat application as necessary.

FOR ANTS: Spray trails, hills, around window frames and openings around pipes and baseboards, where ants crawl hitting as many as possible. Repeat application as necessary.

FOR FLEAS: Spray floors, floor coverings and sleeping quarters of pets and other infested areas. Repeat as required. Do not spray pets.

FOR OUTDOOR USE: To repel temporarily: Flies, mosquitoes, gnats, wasps, crickets, ants. On porches, terraces, lawns, and at outdoor gatherings, spray frequently around chairs, tables, floors, ground and surrounding air.

CONTACT SPRAY — Use with Sprayers that give large, wetting spray Particles — To kill cockroaches, and accessible stages of granary insects: rice & granary weevil, red flour beetle, Indian meal moth larvae, cadelles, grain mites and saw-toothed grain beetles, angoumois grain moth, chocolate moth, carpet beetle, cigarette beetle, flour grain beetle. Use 3-4 oz. per 1000 cu. ft. of space, directing the spray so as to contact the insects. Pay particular attention to cracks, crevices & other hiding places. Repeat treatment as required KILLS INSECTS ONLY WHEN THE SPRAY CONTACTS THEM. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Sanitation is necessary. Clean and remove deposits of food, grain & dust from storage areas. Clean conveyer and handling equipment. For insects affecting stored food, as listed above, flour, grain & cereals — food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Applying using equipment designed & adjusted to deliver a true aerosol mist. Close doors & windows and spray into air, directing spray up to all 4 corners, filling room with

mist. Do not spray on processing equipment. Keep doors & windows closed 15 minutes after spraying. Clean up and destroy fallen insects. Spray when plant is not in operation. Repeat as necessary.

Cover or remove exposed food and close room. Spray outside walls of grain bins, flooring, about base of milling machinery, walls and baseboards, areas behind cabinets, around pipes and other insect harbourages using 1 gal. of spray per 1,000 sq. ft. of surface. Force spray into all cracks and crevices. Repeat applications as necessary. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

SPACE SPRAY — Undiluted in Aerosol Generators — To kill flies, mosquitoes, small flying moths and gnats, cockroaches, and accessible stages of granary insects: rice & granary weevil, red flour beetle, Indian meal moth larvae, cadelles, grain mites and saw-toothed grain beetles, Angoumois grain moth, chocolate moth, carpet beetle, cigarette beetle, flour grain beetle. Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating equipment. Apply at the rate of 4 to 12 oz. of undiluted concentrate per 1,000 cu. ft. of space. Direct output from aerosol generator upward towards upper corners of room and ceiling. Leave room closed for one half hour. Do not remain in treated areas. Ventilate the area after treatment is completed. Sweep up and destroy fallen insects. Repeat as necessary.

SPACE SPRAY — Use in Barns & Poultry Houses, Warehouses, Storage Bins & Storage Rooms — To control house flies, stable flies, and horn flies, apply as a mist over the livestock in stanchions or holding pens or over the poultry in cages or in pens. Use a fogger or mist generator, so spray is directed and applied above the animals and direct spray to center of the area where animals are housed. Apply 2 oz. per 1,000 cu. ft. so that the spray mist is distributed uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in dairy barns, aerate the barn thoroughly before milking. Repeat daily, morning and afternoon, or as required. Keep sprayer unit reservoir and nozzle clean, to ensure proper delivery of spray mist or fog. Product is to be used in mechanical sprayers which give particle size aerosol range or smaller, such as ULV type.

SPACE SPRAY — For Use in Mechanical Fogging Equipment — Kills flies, mosquitoes, small flying moths & gnats. Cover or remove exposed food and food handling surfaces. Close room, making sure to shut off all air conditioning or ventilation equipment. Dilute with deodorized kerosene at a rate of 1 part concentrate with 4 parts kerosene. Apply at the rate of 50 fl. oz. of spray per 25,000 cu. ft. of space. Make sure to direct the spray upward towards upper corners of room & ceiling. Keep room closed for one half hour after spraying. Do not remain in treated area. Ventilate after treatment is completed.

FOODS SHOULD BE REMOVED OR COVERED DURING TREATMENT. ALL FOOD PROCESSING SURFACES SHOULD BE COVERED DURING TREATMENT OR THOROUGHLY CLEANED BEFORE USING. WHEN USING THIS PRODUCT IN THESE AREAS, APPLY ONLY WHEN THE FACILITY IS NOT IN OPERATION.

"Stamp on Backs"